

Published in final edited form as:

Bone Marrow Transplant. 2019 December ; 54(12): 1995–2003. doi:10.1038/s41409-019-0573-6.

Bone Marrow Harvesting from Paediatric Patients Undergoing Hematopoietic Stem Cell Gene Therapy

Francesca Tucci^{#1}, Marta Frittoli^{#1}, Federica Barzagli^{1,2}, Valeria Calbi¹, Maddalena Migliavacca¹, Francesca Ferrua^{1,3,4}, Francesca Fumagalli^{1,5}, Laura Lorioli¹, Laura Castagnaro³, Marcella Facchini³, Claudia Fossati³, Stefano Zancan³, Paola Massariello⁶, Michele Manfredini⁶, Giulia Consiglieri^{1,4}, Daniele Canarutto^{1,4}, Salvatore Recupero^{1,4}, Francesco Calzadini^{1,4}, Miriam Casiraghi³, Silvia Darin³, Gigliola Antonioli³, Roberto Miniero⁷, Rossana Fiori⁸, Paolo Silvani⁸, Matilde Zambelli⁹, Sarah Markt¹⁰, Salvatore Gattillo⁹, Raffaella Milani⁹, Luca Santoleri⁹, Fabio Ciceri^{4,10}, Alessandra Biffi^{3,11}, Maria Pia Cicalese^{1,3}, Maria Ester Bernardo^{1,3}, Alessandro Aiuti^{1,3,4}

¹Pediatric Immunohematology Unit and BMT Program, IRCCS San Raffaele Scientific Institute, Milan, Italy

²Tor Vergata University, Rome, Italy

³San Raffaele Telethon Institute for Gene Therapy (SR-Tiget), IRCCS San Raffaele Scientific Institute, Milan, Italy

⁴Vita-Salute San Raffaele University, Milan, Italy

⁵Neurology Department, IRCCS San Raffaele Scientific Institute, Milan, Italy

⁶Molmed S.p.A, Milan, Italy

⁷Department of Pediatrics, Pugliese-Ciaccio Hospital Magna Graecia University, Catanzaro, Italy

⁸Department of Anesthesia and Critical Care, IRCCS San Raffaele Scientific Institute, Milan, Italy

⁹Immunohematology and Transfusion Medicine Unit, IRCCS San Raffaele Scientific Institute, Milan, Italy

¹⁰Hematology and Bone Marrow Transplantation Unit, IRCCS San Raffaele Scientific Institute, Milan, Italy

¹¹Gene Therapy Program, Dana-Farber/Boston Children's Cancer and Blood Disorders Center, and Harvard Medical School, Boston, USA

These authors contributed equally to this work.

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

Contact Information: Prof. Alessandro Aiuti, San Raffaele Telethon Institute for Gene Therapy (SR-TIGET), Pediatric Immunohematology and Bone Marrow Transplantation Unit, San Raffaele Scientific Institute, Milan, Italy, Via Olgettina, 60, 20132, Milan, Italy. aiuti.alessandro@hsr.it.

Conflict of interest: Fondazione Telethon and San Raffaele Hospital developed gene therapy for ADA-SCID, WAS and MLD for which GlaxoSmithKline (GSK) acquired their license. AA is the PI of the WAS and MLD clinical trials for gene therapy. ADA-SCID gene therapy (Strimvelis) was licensed to GSK in 2010 and received European marketing authorization in 2016. These licenses were transferred to Orchard Therapeutics (OTL) in April 2018. The other authors declare no conflict of interest.

Abstract

Collection of an adequate amount of autologous hematopoietic stem progenitor cells (HSPC) is required for ex vivo manipulation and successful engraftment for certain inherited disorders. Fifty-seven paediatric patients (age 0.5-11.4 years) underwent a bone marrow harvest for the purpose of HSPC gene therapy (GT), including adenosine deaminase severe combined immunodeficiency (ADA-SCID), Wiskott Aldrich syndrome (WAS) and metachromatic leukodystrophy (MLD) patients. Total nucleated cells and percentage and absolute counts of CD34+ cells were calculated at defined steps of the procedure (harvest, CD34+ cell purification, transduction with the gene transfer vector and infusion of the medicinal product). Minimum CD34+ cell dose for infusion was $2 \times 10^6/\text{kg}$, with optimal target at $5\text{-}10 \times 10^6/\text{kg}$.

Median volume of bone marrow harvested was 34.2 ml/kg (range 14.2-56.6). The number of CD34+ cells collected correlated inversely with weight and age in all patients and particularly in the MLD children group. All patients reached the minimum target dose for infusion: median dose of CD34+ cells/kg infused was $10.3 \times 10^6/\text{kg}$ (3.7-25.9), with no difference among the 3 groups. Bone marrow harvest of volumes >30 ml/kg in infants and children with ADA-SCID, WAS and MLD is well tolerated and allows obtaining an adequate dose of medicinal product for HSPC-GT.

Introduction

Gene therapy (GT) with autologous hematopoietic stem/progenitor cells (HSPC) is a promising treatment for primary immunodeficiency, such as adenosine deaminase-severe combined immunodeficiency [1–4] (ADA-SCID) and Wiskott Aldrich syndrome [5,6] (WAS), and inherited metabolic disorders as metachromatic leukodystrophy [7,8] (MLD).

Collection of high numbers of CD34+ HSPC is therefore crucial for in vivo administration of adequate numbers of gene-modified HSPC.

Bone marrow (BM) harvest from iliac crests represents a suitable modality of stem cell procurement in healthy paediatric subjects donating for allogeneic hematopoietic stem cell transplantations (HSCT) of their siblings [9–10], as well as for autologous HSPC transplantations [11–12], autologous back-up [13] or, more recently, for GT. Indeed it has been used for all our San Raffaele-Telethon Institute for Gene Therapy (SR-TIGET) trials for ADA-SCID, WAS and MLD [1,4,5,7] as well as in other gene therapy clinical trials for primary immune deficiencies [2,3,6,14,15].

A direct correlation between cell-dose and engraftment level is a well-known observation in hematopoietic transplant medicine with recommended minimal dose of $> 4 \times 10^8$ total nucleated cells (TNC) [16] or $> 3 \times 10^6$ CD34+ cells per kg of body weight [17] for allogeneic transplantation. For GT clinical trials, usually a minimum target dose at infusion of 2×10^6 CD34+ cells, with a target of $5\text{-}10 \times 10^6$ CD34+ cells/kg, is recommended. Data from ADA-SCID GT clinical trial suggest that the engraftment of transduced progenitors is dependent on the dose of CD34+ cells administered [15]. In WAS patients the degree of myeloid cell engraftment and of platelet reconstitution correlated with the dose of gene-corrected cells administered [6].

GT requires collecting a higher number of cells than usually recommended for a conventional unmanipulated transplantation due to the expected cell loss occurring during subsequent procedures such as mononuclear cell purification, CD34+ cell positive selection, ex vivo cell culture with transduction and possibly cryopreservation. Therefore, we aimed at collecting at least $5-8 \times 10^6$ /kg before any manipulation to achieve the minimal target dose.

Here we evaluated the BM yield, mainly in terms of tolerability and content of HSPCs harvested in children affected by ADA-SCID, WAS and MLD undergoing GT. We also evaluated the presence of patient and harvest related features that could influence the autologous HSPCs collection.

2 Patients and methods

2.1 Patients' and procedure description

We analysed 57 infants and children (age 0.5-11.4 years) who underwent from October 2002 to November 2017 a BM harvest for the purpose of HSPC-GT, including 23 ADA-SCID, 6 WAS and 28 MLD patients. Details about patients' enrolment are reported as supplementary data. Written informed consent was obtained from the donors' parents or legal guardians in accordance with Italian law.

According to the respective clinical protocol/guideline, a HSPC back-up was harvested and cryopreserved unmanipulated to be used in case of poor engraftment or technical issues with product manufacture. About one month after back-up harvest, patients underwent a main harvest for the preparation of transduced medicinal product. Patients' CD34+ cell manipulation and transduction were performed at MolMed S.p.A., a certified Good Manufacturing Practice (GMP) facility.

On average, BM volume collected was set at 20-30 ml/kg of donor weight according to the clinical trial protocols.

To achieve an optimal dose of $5-10 \times 10^6$ CD34+ transduced cells/Kg at infusion our target of CD34+ cell dose collected was $10-20 \times 10^6$ CD34+ cells/Kg before any manipulation. Initial, mid and final harvest TNC count was used to set the total volume to be harvested. Percentage and absolute counts of CD34+ cells were calculated by cell count and cytofluorimetric analysis measured during the procedure at defined steps of the procedure: 1) at the end of BM harvest; 2) CD34+ cell purification; 3) positive selection; 4) transfer vector; 5) quality controls and 6) infusion of the medicinal product.

2.2 Statistical analysis

Statistical analysis was performed with Prism version 6 (GraphPad Software, San Diego, CA, USA). For continuous variables, the results are expressed as medians and ranges. Comparison among patients' harvest parameters and biological characteristics was performed by linear regression analysis or by two-tailed Student t-test. Comparison among 3 or more groups was performed by ANOVA with Bonferroni correction. A p-value less than 0.05 was considered statistically significant. If a p-value is 0.05, 0.01, 0.001 or

0.0001, graphs are flagged with one (*), two (**), three (***) or four star (****), respectively.

3 Results

We analysed 57 paediatric patients who underwent BM harvest at our Institution for the purpose of HSPCs collection for GT ([4,5,8] and data not shown). Data from 23 ADA-SCID patients (16 males, 7 females), 6 male WAS patients and 28 MLD patients (15 males, 13 females) are summarized in Table 1. In the MLD group, 16 were affected by late infantile (LI) variant and 12 by early juvenile (EJ) variant.

All patients underwent a back-up harvest before the main harvest for GT; 54/57 were collected from BM (median 23.5 days of interval between the two procedures) and 3 from mobilized peripheral blood (MPB). Data from back-up harvests are summarized in Suppl. Table 1.

For the main harvest, a median volume of 303.0 ml, 424.3 ml and 420.0 ml (with anticoagulant citrate dextrose), corresponding to a median of 31.2, 34.2 and 36.9 ml/kg of body weight, was collected in ADA-SCID, WAS and MLD patients, respectively (Table 1). There was no significant differences among volumes collected.

The harvested BM processed at the GMP facility (MolMed) contained a median of 43.3, 55.5 and 77.7 x 10⁸ TNCs, corresponding to 4.2, 5.7 and 6.3 x10⁸ cells/kg (Table 1) respectively in ADA-SCID, WAS and MLD patients. ADA-SCID patients showed a significant lower TNC concentration (Figure 1A, ADA-SCID vs MLD patients, p<0.0001) and TNC count/kg (Figure 1B, ADA-SCID vs MLD patients, p<0.0001).

Evaluation of BM CD34+ cell content was first performed in the operating room in order to guide the collection towards the achievement of the HSPC. CD34+ cell frequency and absolute counts were measured during the procedure and after collection. In the harvested bag ADA-SCID patients had a higher statistically significant amount of CD34+ percentage of cells than WAS and MLD patients (median CD34+ 5.7%, 2.9% and 3.0% in ADA-SCID, WAS and MLD; ADA-SCID vs WAS p= 0.033; ADA-SCID vs MLD p=0.0086) (Figure 2A). In contrast we did not observe differences among the 3 groups after normalizing for the volume collected and body weight in terms of CD34+ x10⁶/ml (one way ANOVA test p 0.48) (Figure 2B) and CD34+ x10⁶ cells/kg (one way ANOVA test p= 0.17) (Figure 2C).

Next we analysed the influence of patient related factors with respect to TNC and CD34+ cell harvest. MLD LI patients showed a higher percentage and concentration compared to the EJ variant (median CD34+ x10⁶/ml in LI 0.76, in EJ 0.33; p= 0.0005).

In order to define factors affecting cell dose, the correlation between the collected HSPCs and weight and age was studied.

Collected TNCs were analysed in relationship with patient's characteristics, age and weight (Figure 3A-B). The cumulative group of all patients showed an inverse correlation only for age (p=0.03). MLD patients showed a significant inverse correlation with respect to the two

parameters ($p=0.0001$ for both), indicating a reduction in TNCs harvested for patients with higher weight and age. On the contrary, a direct correlation was seen for ADA-SCID patients, regarding weight ($p=0.04$) (Fig. 3A) but not age (Fig. 3B). No correlation was seen in the WAS disease setting. Then we investigated the relationship between the CD34+ cell concentration and donor's weight and age in the 3 populations. An inverse correlation was documented in the group of all patients and in the subgroup of MLD patients for weight ($p<0.0001$) (Fig. 3C) and age ($p=0.0001$) (Fig. 3D) with a similar trend in WAS patients.

The harvested volume was used as a correlation parameter for TNCs and CD34+ cells. In the cumulative patient cohort, TNC and CD34+ cell concentration was shown to be inversely correlated to the harvested volume ($p=0.01$ and $p<0.0001$, respectively) (Figure 4A and 4C). We hypothesised that in children, where aspiration surface is limited, a possible hemodilution might occur during harvest procedure. Therefore, volume was normalised to patient's body weight. We confirmed a significant inverse correlation also for TNC concentration with volume/Kg (Figure 4B) and a trend for CD34+ cell concentration (Figure 4D), suggesting that increasing the aspirated volume can result in reduced cell concentration. When analysing these parameters by disease, MLD patients displayed a similar pattern for both TNC and CD34+ cells, whereas the correlation was significant for TNC concentration and normalised blood volume in ADA-SCID patients. There was no correlation in WAS patients, possibly also due to the limited sample size of this group.

CD34+ cell counts after manipulation and transduction are summarized in Table 1. A median of 10.4 selected CD34+ cells $\times 10^6/\text{Kg}$ were obtained after HSPC purification; 13.8 , 9.1 and 9.6 CD34+ $\times 10^6/\text{Kg}$ cells were obtained in ADA-SCID, WAS and MLD children, respectively. After transduction, a median of 10.3 CD34+ $\times 10^6/\text{Kg}$ were infused; 10.7 , 9.0 and 10.3 CD34+ cells/kg were infused in the 3 patients' groups, in line with the target range ($5-10 \times 10^6/\text{Kg}$) of the clinical protocols. No statistical difference was observed within the 3 disease groups (one way ANOVA test $p=0.71$). The infused dose appeared to be directly associated with the amount of CD34+ cells collected and purified after the BM harvest (Figure 5).

We next evaluated the cell yield during the different steps of the production process. The median cell yield after CD34+ cell purification was 58.3% ($39.5-94.1$) and was similar for all the 3 groups (Figure 6A). Median cell recovery after transduction was 92.1% ($53.6-138.5$) with no substantial differences in the 3 groups despite different culture conditions among the ADA-SCID, WAS and MLD protocols (Figure 6B).

We also investigated if donor age has a relationship with selected (Figure 7A) and infused transduced CD34+ cell counts (Figure 7B). Overall, we observed the same trend previously seen for the unmanipulated HSPCs with a significant inverse correlation with respect to age ($p<0.0001$ both for selected and for infused cells). Among the 3 groups of disease, only selected CD34+ cell/kg of MLD patients showed a reduction of CD34+ cell counts in older patients ($p=0.0007$) (Figure 7A). With regards to infused cell, an indirect correlation with age was seen both in ADA-SCID and in MLD patients (ADA-SCID $p=0.0012$; MLD $p=0.0079$) (Figure 7B).

4 Discussion

The amount of harvested HSPCs is a critical parameter for GT because it represents the starting material for the preparation of the medicinal product. Harvested CD34+ cells need to be purified and *ex vivo* cultured to produce the drug product that is infused in the patient. BM historically has been the preferred source for HSPCs collection in the GT setting, although MPB has been increasingly adopted in the last years [9]. To date, few data are available regarding BM harvest in paediatric population, mainly limited to the maximum amount that can be safely collected and the purification yields of CD34+ cells [18–19]. Moreover, scattered data on specific diseases such as primary immunodeficiency or lysosomal storage disorders are reported [13,20].

Here we performed a comprehensive analysis of the tolerability of large volume collection and of the quality of BM harvest in the context of an autologous GT setting. Moreover, we evaluated the impact of patients' and harvest related features on the amount of HSPCs available for cell manipulation and transduction in children enrolled in clinical trials of GT for ADA-SCID, WAS and MLD.

We found that the number of TNCs and HSPCs harvested and selected for GT, depends on patients- and harvest-related factors. In our study, we aimed at infusing at least 2×10^6 CD34+/kg with an optimal target dose of $5-10 \times 10^6$ CD34+/kg. Taking into account the HSPCs loss due to cell manipulation (selection and transduction), a target range of $10-20 \times 10^6$ CD34+ cells/Kg at collection was considered for the BM harvest. A single BM harvest was sufficient for most of the patients to obtain the optimal target dose of hematopoietic progenitor cells for transplantation. Six out 57 patients were below the target range but still above the minimal dose. For 3 of these an additional aliquot of purified CD34+ cells, previously collected during back-up leukapheresis harvest (n=2) or BM (n=1), was available and used to produce a second lot of transduced cells in order to increase the amount of cells infused. Two additional patients received transduced cells obtained from the main BM harvest and previously stored CD34+ cells from BM (n=1) or MPB (n=1) (data not shown).

The BM procedure was overall well tolerated in all the children. No major complications were reported despite the median volume collected was higher than the amount conventionally indicated and reported in previous published studies in paediatric healthy donors (up to 23.8 ml/Kg) [18–19] and in ADA-SCID patients who underwent GT in other centers (up to 20 ml/Kg) [3,15]. Moreover, no significant difference in CD34+ cell concentration was observed in patients who underwent a BM back up before the main GT BM harvest, suggesting that two collection procedures do not preclude harvesting of HSPCs for GT after a relative short time interval (median of 23.5 days).

In a single-center report on 109 paediatric healthy donors, BM volumes exceeded the standard BM volumes per body weight in 65% of the cases with a reported median BM volume/kg of 18 ml which corresponding to half of our median values [21]. The authors did not report severe adverse events and suggested that larger volumes are safe with the exception of the higher chances of being exposed to allogeneic blood products, which anyhow would be administered in the case of GT.

Another recent retrospective study showed large collected volumes harvested for the purpose of both autologous and allogeneic BM transplantations in children. Median amount of BM volume were comparable to our results (37.66 ml/kg; range 20-55.3 ml/kg) with no mention about adverse events related to the procedures [22].

Concerning the harvest characterization, a significant higher CD34+ percentage of cells was found in ADA-SCID BM compared to those of WAS and MLD patients. This difference can be ascribed to the know block in B- cell differentiation in the BM of these patients [23] and/or to the lower amount of nucleated cells intrinsically related to the immunodeficiency status of the disease. Indeed a lower TNCs content was documented in ADA-SCID patients compared to MLD patients. After normalization for harvested volume and body weight CD34+ cell content was similar in ADA-SCID patients with respect to WAS and MLD children. This indicates that the difference among patients was only relative to CD34+ cell proportion and not to absolute count.

The amount of BM CD34+ cell retrieved after harvest (median $0.5 \times 10^6/\text{mL}$) was in line with the data from of Furey et al. with a median of median $0.7 \times 10^6/\text{mL}$ in children <6 years [19], and higher than those reported in two other studies [22,24].

There were minor discrepancies in the range of recoveries of CD34+ cells between the theoretical number at harvest, calculated based on interim analyses in the operating room and the final values counted at the GMP facility before starting the manipulation of the cells. These could be due to the different methods used for counting CD34+ cells. The cell yield after CD34+ cell purification (median 58.3%) was in line with previous report [25] and should take in consideration the fact that the process that included also a density gradient centrifugation in addition to immune-magnetic purification. Following ex vivo cell culture and transduction for a total of 3 days (MLD and WAS) and 4 days (ADA-SCID) the median yield of CD34+ cells was about 92.1%, with one third of patients showing CD34+ cell expansion. A direct positive correlation was observed between selected and infused CD34+ cells per kilogram of body weight for all ADA-SCID, WAS and MLD groups suggesting that, in spite of some variability among the patients, the steps leading to the final product could be considered stable and reproducible.

Donor/recipient characteristics are crucial factors that can influence stem cell yield and directly affect the outcome of gene therapy, which requires patients' selected CD34+ cells as starting product for gene modifications. The correlation between TNC and CD34+ cells collected and the clinical features of the children was evaluated in order to establish if they could be direct indicators of rich or poor cell collection.

A significant difference between LI and EJ variants in MLD children was observed in terms of CD34+ cell content. This could be related to the fact that EJ MLD patients have a later onset and therefore are usually harvested at an older age whereas only pre-symptomatic LI were enrolled in the MLD clinical trial. Indeed, CD34+ cell yield correlates inversely with weight and age at harvest in MLD children while only a tendency was seen in ADA-SCID and WAS children.

Similarly to other reports, we found that significantly higher CD34+ cell counts are observed in younger donors [18,19,24]. Friebert et al., found that the cellularity of pediatric BM, including healthy donors, was highest in patients younger than 2 years (approximately 80%), declined to approximately 60% by the age of 5 years and then remained relative constant in patients aged 5-18 years [26]. Furey et al. showed that the median CD34+ cell count obtained from pediatric sibling donors is higher in the younger donors, suggesting that the volume of product harvested might be decreased without significantly changing the overall CD34+ cell count [19]. These observations were similar to the findings of the present study in MLD patients, whose BM status may be considered closer to healthy donors because of the absence of haematological and immunological defects. The same correlation was not documented in ADA-SCID and WAS diseases, where immunodeficiency is often associated to growth impairment and age increase does not always correspond to weight increase. Yet, a relationship between age of the donor and the dose of infused CD34+ cells after transduction was documented in ADA-SCID patients similarly to the findings of Shaw et al. in their phase I/II trial of GT for ADA-SCID [15].

In summary, the cumulative data indicate that harvesting BM for the purpose of autologous gene therapy is a well-tolerated procedure that allows yielding an adequate stem cell amount for reaching a target cell dose at infusion of the engineered HSPCs. Additional studies are needed to compare the outcome of gene therapy after infusion of gene corrected HSPC from different sources (BM and MPB).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors would like to thank the entire medical and nurse personnel of the Paediatric Immunohematology and Hematology and Bone Marrow Transplant Unit of San Raffaele Hospital (Milan), the personnel of the SR-TIGET Clinical Trial Office, local referring physicians who helped with patient management, and all patients who participated in this study and their families. The authors thank Michela Gabaldo (SR-TIGET) for her continuous support to the projects and Giuliana Tomaselli, Luisella Meroni and Samih El Hossary (all of SR-Tiget) for their support to patients.

Abbreviations

ADA-SCID	Adenosine Deaminase Severe Combined Immunodeficiency
BM	Bone Marrow
CUP	Compassionate Use Program
EJ	Early Juvenile
GMP	Good Manufacturing Practice
GT	Gene Therapy
HSPC	Hematopoietic Stem/Progenitor Cell

HSCT	Hematopoietic Stem Cell Transplant
LI	Late Infantile
MLD	Metachromatic Leukodystrophy
MPB	Mobilized Peripheral Blood
SR-TIGET	San Raffaele-Telethon Institute for Gene Therapy
TNC	Total Nucleated Cells
WAS	Wiskott Aldrich Syndrome

References

1. Aiuti A, Cattaneo F, Galimberti S, Benninghoff U, Cassani B, Callegaro L, et al. Gene therapy for immunodeficiency due to adenosine deaminase deficiency. *N Engl J Med*. 2009; 360(11):447–458. [PubMed: 19179314]
2. Gaspar HB, Cooray S, Gilmour KC, Parsley KL, Zhang F, Adams S, et al. Hematopoietic stem cell gene therapy for adenosine deaminase-deficient severe combined immunodeficiency leads to long-term immunological recovery and metabolic correction. *Sci Transl Med*. 2011; 3(97):97ra80.
3. Candotti F, Shaw KL, Muul L, Carbonaro D, Sokolic R, Choi C, et al. Gene therapy for adenosine deaminase-deficient severe combined immune deficiency: clinical comparison of retroviral vectors and treatment plans. *Blood*. 2012; 120(18):3635–46. [PubMed: 22968453]
4. Cicalese MP, Ferrua F, Castagnaro L, Pajno R, Barzaghi F, Giannelli S, et al. Update on the safety and efficacy of retroviral gene therapy for immunodeficiency due to adenosine deaminase deficiency. *Blood*. 2016; 128(1):45–54. [PubMed: 27129325]
5. Aiuti A, Biasco L, Scaramuzza S, Ferrua F, Cicalese MP, Baricordi C, et al. Lentiviral hematopoietic stem cell gene therapy in patients with Wiskott-Aldrich syndrome. *Science*. 2013 Aug 23.341(6148)
6. Hacein-Bey Abina S, Gaspar HB, Blondeau J, Caccavelli L, Charrier S, Buckland K, et al. Outcomes following gene therapy in patients with severe Wiskott-Aldrich syndrome. *JAMA*. 2015; 313(15):1550–63. [PubMed: 25898053]
7. Biffi A, Montini E, Lorioli L, Cesani M, Fumagalli F, Plati T, et al. Lentiviral hematopoietic stem cell gene therapy benefits metachromatic leukodystrophy. *Science*. 2013 Aug 23.341(6148)
8. Sessa M, Lorioli L, Fumagalli F, Acquati S, Redaelli D, Baldoli C, et al. Lentiviral haemopoietic stem-cell gene therapy in early-onset metachromatic leukodystrophy: an ad-hoc analysis of a non-randomised, open-label, phase 1/2 trial. *Lancet*. 2016; 388(10043):476–87. [PubMed: 27289174]
9. Passweg JR, Baldomero H, Peters C, Gaspar HB, Cesaro S, Dreger P, et al. Hematopoietic SCT in Europe: data and trends in 2012 with special consideration of pediatric transplantation. *Bone Marrow Transplant*. 2014; 49(6):744–50. [PubMed: 24637898]
10. Khandelwal P, Millard HR, Thiel E, Abdel-Azim H, Abraham AA, Auletta JJ, et al. Hematopoietic Stem Cell Transplantation Activity in Pediatric Cancer between 2008 and 2014 in the United States: A Center for International Blood and Marrow Transplant Research Report. *Biol Blood Marrow Transplant*. 2017; 23(8):1342–1349. [PubMed: 28450183]
11. Weisdorf DJ, Verfaillie CM, Miller WJ, Blazar BR, Perry E, Shu XO, et al. Autologous bone marrow versus non-mobilized peripheral blood stem cell transplantation for lymphoid malignancies: a prospective, comparative trial. *Am J Hematol*. 1997; 54(3):202–8. [PubMed: 9067498]
12. Petersen FB, Lynch MH, Clift RA, Appelbaum FR, Sanders JE, Bensinger WI, et al. Autologous marrow transplantation for patients with acute myeloid leukemia in untreated first relapse or in second complete remission. *J Clin Oncol*. 1993; 11(7):1353–60. [PubMed: 8315433]
13. Frittoli MC, Biral E, Cappelli B, Zambelli M, Roncarolo MG, Ferrari G, et al. Bone marrow as a source of hematopoietic stem cells for human gene therapy of β -thalassemia. *Hum Gene Ther*. 2011; 22(4):507–13. [PubMed: 20979441]

14. Hacein-Bey-Abina S, Pai SY, Gaspar HB, Armant M, Berry CC, Blanche S, et al. A modified γ -retrovirus vector for X-linked severe combined immunodeficiency. *N Engl J Med*. 2014; 371(15): 1407–17. [PubMed: 25295500]
15. Shaw KL, Garabedian E, Mishra S, Barman P, Davila A, Carbonaro D, et al. Clinical efficacy of gene-modified stem cells in adenosine deaminase-deficient immunodeficiency. *J Clin Invest*. 2017; 127(5):1689–1699. [PubMed: 28346229]
16. Kalwak K, Porwolik J, Mielcarek M, Gorczy ska E, Owoc-Lempach J, Ussowicz M, et al. Higher CD34(+) and CD3(+) cell doses in the graft promote long-term survival, and have no impact on the incidence of severe acute or chronic graft-versus-host disease after in vivo T cell-depleted unrelated donor hematopoietic stem cell transplantation in children. *Biol Blood Marrow Transplant*. 2010; 16(10):1388–401. [PubMed: 20382248]
17. Bittencourt H, Rocha V, Chevret S, Socié G, Espérou H, Devergie A, et al. Association of CD34 cell dose with hematopoietic recovery, infections, and other outcomes after HLA-identical sibling bone marrow transplantation. *Blood*. 2002; 99(8):2726–33. [PubMed: 11929759]
18. Yabe M, Morimoto T, Shimizu T, Koike T, Takakura H, Ohtsubo K, et al. Feasibility of marrow harvesting from pediatric sibling donors without hematopoietic growth factors and allotransfusion. *Bone Marrow Transplant*. 2014; 49(7):921–6. [PubMed: 24777192]
19. Furey A, Rastogi S, Prince R, Jin Z, Smilow E, Briamonte C, et al. Bone Marrow Harvest in Pediatric Sibling Donors: Role of Granulocyte Colony-Stimulating Factor Priming and CD34+ Cell Dose. *Biol Blood Marrow Transplant*. 2018; 24(2):324–329. [PubMed: 29074373]
20. Biral E, Chiesa R, Cappelli B, Rocca T, Frugnoli I, Noè A, et al. Multiple BM harvests in pediatric donors for thalassemic siblings: safety, efficacy and ethical issues. *Bone Marrow Transplant*. 2008; 42(6):379–84. [PubMed: 18574444]
21. van Walraven SM, Straathof LM, Switzer GE, Lankester A, Korthof ET, Brand A, Ball LM. Immediate and long-term somatic effects, and health-related quality of life of BM donation during early childhood. A single-center report in 210 pediatric donors. *Bone Marrow Transplantation*. 2013; 1:40–45.
22. Witt V, Pichler H, Fritsch G, Peters C. Multiple small versus few large amount aspirations for bone marrow harvesting in autologous and allogeneic bone marrow transplantation. *Transfus Apher Sci*. 2016; 55(2):221–224. [PubMed: 27452644]
23. Brigida I, Sauer AV, Ferrua F, Giannelli S, Scaramuzza S, Pistoia V, et al. B-cell development and functions and therapeutic options in adenosine deaminase-deficient patients. *J Allergy Clin Immunol*. 2014; 133(3):799–806. [PubMed: 24506932]
24. Fettah A, Özbek N, Özgüner M, Azık F, I ık P, Avcı Z, et al. Factors associated with bone marrow stem cell yield for pediatric allogeneic stem cell transplantation: The impact of donor characteristics. *Pediatr Transplant*. 2017; 21(1)
25. Gaipa G, Dassi M, Perseghin P, Venturi N, Corti P, Bonanomi S, et al. Allogeneic bone marrow stem cell transplantation following CD34+ immunomagnetic enrichment in patients with inherited metabolic storage diseases. *Bone Marrow Transplant*. 2003; 31(10):857–60. [PubMed: 12748661]
26. Friebert SE, Shepardson LB, Shurin SB, Rosenthal GE, Rosenthal NS. Pediatric bone marrow cellularity: are we expecting too much? *J Pediatr Hematol Oncol*. 1998; 20(5):439–43. [PubMed: 9787316]




Figure 1. TNC concentration (A) and TNC $\times 10^8/\text{kg}$ (B) in ADA SCID, WAS, MLD. Each box plot displays the distribution of data based on the five number summary: minimum, first quartile, median, third quartile, and maximum.




Figure 2.

CD34+ cell values are expressed as percentages (A), concentration (B) and per kg of body weight (C) in 8 ADA SCID, 5 WAS and 28 MLD patients. Data measured after collection at the GMP facility are shown.




Figure 3. Relationship between the weight (A) and age (B) of the donor and concentration of TNCs harvested in the 3 patients subgroups and in all the patients. Relationship between patients' weight (C) and age (D) with concentration of CD34+ cells in the BM harvest.




Figure 4. Correlation between TNC concentration and volume (A) or volume/Kg (B) and between CD34⁺ cell concentration and Volume (C) or Volume/Kg (D) in the 3 patients subgroups and in all the patients.




Figure 5.
Relationship between selected and infused CD34+ cells in the 3 patients subgroups and in all the patients.




Figure 6. Cell yield during the consecutive steps of the production process. We calculated cell yield between harvest and selection (A) and cell yield after transduction step (B).




Figure 7. relationship between age of the donor and CD34+ cells (as CD34+ x 10⁶/kg) after purification (A) and after cell manipulation and transduction (B).

Table 1
Summary data of bone marrow harvests for GT. Data indicate median and range.

^data available for the last consecutive 8 ADA SCID patients.

Disease	Age at GT (years)	Weight (kg)	Harvested volume (ml)	Volume/kg	Collected TNC (x10 ⁸)/kg	CD34+ (%)	Collected CD34+ cells (x10 ⁶)/kg	Selected CD34+ cells (x10 ⁶)/kg	Infused CD34+ cells (x10 ⁶)/kg
ADA SCID (n=23)	1.7 (0.5-6.0)	10.1 (5.9-26.5)	303.0 (190.0-1167.5)	31.2 (14.2-48.6)	4.2 (2.9-6.8)	5.7 (2.0-6.7)^	28.9 (9.8-43.1)^	13.8 (4.4-22.9)	10.7 (3.7-19.7)
WAS (n=6)	2.1 (0.9-6.0)	12.7 (8.0-26.3)	424.3 (275.0-773.0)	34.2 (29.4-36.5)	5.7 (3.5-6.6)	2.9 (1.8-5.2)	15.8 (10.8-18.7)	9.1 (4.5-14.5)	9.0 (3.7-14.1)
MLD (n=28)	1.3 (0.5-11.4)	11 (7.3-25.3)	420.0 (194.0-1132.0)	36.9 (17.6-56.6)	6.3 (3.4-9.4)	3.0 (1.0-7.7)	16.3 (5.6-65.9)	9.6 (3.9-34.7)	10.3 (3.8-25.9)
Total (n=57)	1.5 (0.5-11.4)	10.6 (5.9-26.5)	348.0 (190.0-1167.5)	34.2 (14.2-56.6)	5.1 (2.9-9.4)	3.1 (1.0-7.7)	17.3 (5.6-65.9)	10.4 (3.9-34.7)	10.3 (3.7-25.9)