

Published in final edited form as:

Osteoporos Int. 2019 November 01; 30(11): 2155–2165. doi:10.1007/s00198-019-05104-5.

Determinants, consequences and potential solutions to poor adherence to anti-osteoporosis treatment: results of an expert group meeting organized by the European Society for Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases (ESCEO) and the International Osteoporosis Foundation (IOF)

Mickael Hiligsmann¹, Dennis Cornelissen¹, Bernard Vrijens², Bo Abrahamsen^{3,4,5}, Nasser Al-Daghri⁶, Emmanuel Biver⁷, Maria-Luisa Brandi⁸, Oliver Bruyère⁹, Nansa Burlet¹⁰, Cyrus Cooper¹¹, Bernard Cortet¹², Elaine Dennison¹³, Adolfo Diez-Perez¹⁴, Andrea Gasparik¹⁵, Alda Grosso¹⁶, Peyman Hadji¹⁷, Philippe Halbout¹⁸, John A Kanis¹⁹, Jean-Marc Kaufman²⁰, Andrea Laslop²¹, Stefania Maggi²², René Rizzoli⁷, Thierry Thomas²³, Sansin Tuzun²⁴, Mila Vlaskovska²⁵, Jean-Yves Reginster^{6,9}

¹Department of Health Services Research, CAPHRI Care and Public Health Research Institute, Maastricht University, Maastricht, The Netherlands ²Research and Development, AARDEX Group, Liège, Belgium and Department of Public Health, University of Liège, Liège, Belgium ³Open Patient Data Explorative Network, University of Southern Denmark, Institute of Clinical Resesarch, Odense Denmark ⁴Holbæk Hospital, Department of Medicine, Holbæk, Denmark ⁵NDORMS, University of Oxford, Oxford, United Kingdom ⁶Chair for Biomarkers of Chronic

Correspondence to: Mickael Hiligsmann, Department of Health Services Research, Maastricht University, P.O. Box 616, 6200 MD Maastricht, the Netherlands, T +31 43 38 82 219 F +31 43 38 841 62; m.hiligsmann@maastrichtuniversity.nl.

Conflicts of interest

MLB has received honoraria from Amgen, Bruno Farmaceutici, Kyowa Kirin; academic grants and/or speaker from Abiogen, Alexion, Amgen, Bruno, Farmaceutici, Eli Lilly, Kyowa Kirin, MSD, NPS, Servier, Shire, SPA and consulting fees from Alexion, Bruno Farmaceutici, Kyowa Kirin, Servier, Shire.

OB received grants research from Biophytis, IBSA, MEDA, Servier, SMB and consulting or lecture fees from Amgen, Biophytis, IBSA, MEDA, Servier, SMB, TRB Chemedica, UCB.

BC has received Fees for occasional interventions as an expert or speaker for Amgen, Expanscience, Ferring, Lilly, Medtronic, MSD, Mylan, Novartis, Roche Diagnostics, and UCB

CC has received lecture fees and honoraria from Amgen, Danone, Eli Lilly, GSK, Kyowa Kirin, Medtronic, Merck, Nestlé, Novartis, Pfizer, Roche, Servier, Shire, Takeda and UCB outside of the submitted work.

ED has received honoraria from Pfizer and UCB.

ADP has received consulting and speaker fees from Amgen, UCB, Eli Lilly, Roche, Gilead, and own shares of Active Life Sci.

JAK reports grants from Amgen, Eli Lilly and Radius Health; consulting fees from Theramex.

PH has received consulting and speaker fees from Amgen, Dr. Kade/Besin, Gedeon Richter, Eli Lilly, Meda/Mylan, Novartis, Pfizer, Roche, Theramex and UCB and has given expert testimony on behalf of Novartis.

RR has been speaker bureau or member of scientific advisory boards for Danone, Echolight, Effryx, Mylan, Nestlé, ObsEva, Pfizer, Radius Health, Sandoz and TEVA/Theramex.

TT reports personal fees for lectures and expertises from Abbvie, Amgen, Arrow, BMS, Chugai, Expanscience, Gilead, HAC-Pharma, LCA, Lilly, Medac, MSD, Pfizer, Theramex, Thuasne, TEVA and UCB and financial support or fees for research activities from Amgen, Bone Therapeutics, Chugai, HAC-Pharma, MSD, Novartis, Pfizer, and UCB

JYR has received consulting fees or paid advisory boards from IBSA-GENEVRIER, Mylan, Radius Health, Pierre Fabre, lecture fees when speaking at the invitation of sponsor for IBSA-GENEVRIER, Mylan, CNIEL, Dairy Research Council and grant support from industry (all through institution) from IBSA-GENEVRIER, Mylan, CNIEL, Radius Health.

The others authors have no conflict of interest to declare relevant to the content of this study.

Diseases, Biochemistry Department, College of Science, King Saud University, Riyadh, Saudi Arabia ⁷Division of Bone Diseases, Geneva University Hospitals and Faculty of Medicine, Geneva, Switzerland ⁸FirmoLab. Fondazione FIRMO e Università di Firenze, Firenze, Italy ⁹Division of Public Health, Epidemiology and Health Economics, University of Liège, Liège, Belgium and WHO Collaborating Centre for Public Health Aspects of Musculoskeletal Health and Aging, Liège, Belgium ¹⁰Global Head of Patient Insights Innovation, Patient Solution Unit, Sanofi ¹¹MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK; NIHR Southampton Biomedical Research Centre, University of Southampton and University Hospital Southampton; NHS Foundation Trust, Southampton, UK; NIHR Oxford Biomedical Research Centre, University of Oxford, Oxford, UK ¹²Department of Rheumatology and EA 4490, University-Hospital of Lille, Lille, France ¹³MRC Lifecourse Epidemiology Unit, University of Southampton, Southampton, UK ¹⁴Musculoskeletal Research Unit, IMIM-Parc Salut Mar, CIBERFES, Universitat Autònoma de Barcelona, Barcelona, Spain ¹⁵Department of Public Health, University of Medicine, Pharmacy, Science and Technology of Tirgu Mures, Tirgu Mure , Romania ¹⁶Patient partner ¹⁷Frankfurt Centre of Bone Health, Frankfurt, Germany & Philips-University of Marburg, Marburg, Germany ¹⁸International Osteoporosis Foundation, Nyon, Switzerland ¹⁹Centre for Metabolic Bone Diseases, University of Sheffield Medical School, Sheffield, UK and Mary McKillop Health Institute, Catholic University of Australia, Melbourne, Australia ²⁰Department of Endocrinology, Ghent University Hospital, Ghent, Belgium ²¹Scientific Office, Austrian Federal Office for Safety in Health Care, Vienna, Austria ²²CNR Aging Branch-NI, Padua, Italy ²³Department of Rheumatology, Hôpital Nord, CHU Saint-Etienne, Saint-Etienne and INSERM U1059, Université de Lyon-Université Jean Monnet, Saint-Etienne, France ²⁴Department of Physical Medicine and Rehabilitation, Cerrahpa a Medical Faculty, stanbul University Cerrahpasa, Istanbul, Turkey ²⁵Medical Faculty, Department of Pharmacology, Medical University Sofia, Sofia, Bulgaria

Abstract

Purpose—Poor adherence to medications is a major challenge in the treatment of osteoporosis. This paper aimed to provide an overview of the consequences, determinants and potential solutions to poor adherence and persistence to osteoporosis medication.

Methods—A working group was organized by the European Society on Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal diseases (ESCEO) to review consequences, determinants and potential solutions to adherence, and to make recommendations for practice and further research. A systematic literature review and a face-to-face experts meeting were undertaken.

Results—Medication non-adherence is associated with increased risk of fractures, leading to a substantial decrease in the clinical and economic benefits of drug therapy. Reasons for non-adherence are numerous and multidimensional for each patient, depending on the interplay of multiple factors, suggesting the need for multi-factorial and individualized solutions. Few interventions have been shown to improve adherence or persistence to osteoporosis treatment. Promising actions include patient education with counselling, adherence monitoring with feedback and dose simplification including flexible dosing regimen. Recommendations for practice and

further research were also provided. To adequately manage adherence, it is important to (1) understand the problem (initiation, implementation and/or persistence), (2) to measure adherence and (3) to identify the reason of non-adherence and fix it.

Conclusion—These recommendations are intended for clinicians to manage adherence of their patients, and to researchers and policy makers to design, facilitate and appropriately use adherence interventions.

Keywords

adherence; determinants; osteoporosis; persistence; solutions

Introduction

Poor or non-adherence to medication remains a major problem in most chronic diseases, including osteoporosis. Effective and safe medications are available to reduce the risk of fractures [1], but numerous patients do not initiate treatment for osteoporosis or do not take it appropriately, resulting in a substantial clinical and economic burden [2]. Initiation of osteoporosis therapy is even poorer than in other disease areas, with a substantial decrease in bisphosphonates initiation observed in the last few years in Europe and the US [3, 4]. Investing in medication adherence could improve health outcomes and health system efficiency [5].

Poor or non-adherence with osteoporosis medications is not a new problem. Several studies have already assessed the consequences and burden of non-adherence with osteoporosis medications being one of the major challenges of successful osteoporosis management [6], and have highlighted the urgency of managing medication adherence. In 2012, a systematic review of interventions to improve adherence to osteoporosis medications suggested few high-quality studies and mixed effects of interventions [7]. In recent years, osteoporosis management has evolved (e.g. new treatments, new diagnostic tools, FLS organizations), more interventions to improve adherence have been tested and a better understanding of the determinants of non-adherence is available.

Considering the burden of non-adherence with osteoporosis medications and the need to provide up-to-date recommendations to manage medication adherence, a working group was convened by the European Society on Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases (ESCEO) to review the consequences and determinants of poor adherence and interventions to manage adherence and improve persistence to osteoporosis medication. In addition, this paper aims to provide recommendations for practice and for further research. Information included in this review could be of value to help clinicians to manage adherence of their patients, but also for researchers and policy makers involved in osteoporosis.

Methods

An international working group was formed to review the consequences and determinants of poor adherence and interventions, to manage adherence and improve persistence to

osteoporosis medication, and to provide recommendations for practice and for further research. The working group comprised clinical scientists, researchers (including experts in adherence) and a patient selected by the Scientific Advisory Board of ESCEO.

In preparation for a working group meeting, a systematic literature review was conducted to identify articles about the determinants and interventions to improve adherence or persistence to osteoporosis medications. The literature search was conducted for articles up to December 2018 in PubMed. Further details about the search strategy and inclusion of articles can be found in Cornelissen et al. [8]

A face-to-face meeting then took place on January 22nd 2019. The meeting started with three short presentations about the clinical and economic consequences of non-adherence to osteoporosis medications (JYR), the determinants of non-adherence to osteoporosis medications (MH) and the potential solutions to improve adherence to anti-osteoporosis medications (DC). A group discussion led by MH and ADP followed. Each participant had the opportunity to comment on the determinants, consequences and potential solutions, and then to provide recommendations for practice and further research. Following this meeting and a symposium that was held at the World Congress of Osteoporosis (WCO-IOF-ESCEO congress, 6th of April 2019), members of the writing team (MH, DC, BV, BA, JYR, JK, AG, PH) reviewed a first version of the article drafted by MH; this was then reviewed and commented on by all members of the working group.

Terminology

In the literature, a number of terms such as ‘adherence,’ ‘compliance’, ‘concordance’ and ‘persistence’ have been used to define how patients take their medicines [9]. In 2012, a collaboration of European research groups in the field of medication adherence funded by the European Commission suggested the ABC taxonomy for describing and defining adherence to medications [9], that will be used in this paper. Adherence to medication is defined as “the process by which patients take their medications as prescribed, composed of A) initiation, B) implementation and C) discontinuation” [9, 10]. *Initiation* occurs when the patient takes the first dose of a prescribed medication, *discontinuation* when the patient stops taking the medication and *implementation* is the extent to which a patient's actual dosing corresponds to the prescribed dosing regimen, from initiation until the last dose. Medication persistence is further defined as the length of time between initiation and the last dose, which immediately precedes discontinuation [9]. Overall, this definition is in agreement with the definition of the *Group for the Respect of Ethics Excellence in Science* in osteoporosis [11], with the exception that implementation has replaced compliance, and initiation is preferred over primary adherence.

Adherence to osteoporosis medications, clinical and economic consequences

In a recent review including 124 studies, the prevalence of medication adherence ranged from 12.9% to 95.4% [12]. Several studies have reported that initiation, implementation and persistence to osteoporosis medications are suboptimal. In line with treatment initiation rates

in other diseases [13], about 20-30% of patients do not initiate a treatment after a prescription for oral bisphosphonates [14]. In addition, patients on bisphosphonates frequently miss doses and therefore do not implement as prescribed [15]. Persistence rates at one year for oral bisphosphonates are commonly estimated between 16 and 60% [16–18], in line with a review of 95 clinical studies from different disease areas suggesting that about 40% of patients discontinued at one year [19]. For example, in a large observational Belgian study, persistence rates to oral bisphosphonates were estimated at 39.5% at one year, without a gap of more of 5 weeks in treatment [20], while only 48% of patients were adherent (defined as a medical possession ratio (MPR) \geq 80%). In another study, van Boven et al. showed that persistence to weekly oral bisphosphonates was about 60% at 1 year and decreased to 25% at 5 years [21]; in line with a Danish study [22]. Other studies have even reported that less than 20% were still on treatment as soon as after 12 months after treatment initiation [23]. Differences in methodologies (e.g. permissible gap to define persistence, adherence measures, incorporation of switchers) could explain the differences between studies and made direct comparison between them difficult.

Further, osteoporosis remains largely underdiagnosed and a substantial number of patients at increased risk of fractures do not even receive any prescription for osteoporotic therapy [24], or only calcium and vitamin D supplementation. The osteoporosis treatment gap was estimated between 25% and 95% in European countries [25]. Even in randomised controlled trials, persistence and adherence with therapy declines over time, and any reduced effectiveness caused by sub-optimal adherence is to some extent already captured in clinical trials.

Different profiles of non-adherent patients could thus be identified [26]. Some patients never initiate a treatment, while others delay initiation of therapy. There are patients who frequently miss doses, and multi-week drug holiday periods have also been observed. Several patients further discontinue treatment earlier than prescribed. In a US study, it was also observed that among discontinuers, about 45% [27] reinitiated therapy, with the majority doing so within 6 months of discontinuation, suggesting that many patients are thus exposed to multiple episodes of starting and stopping of drug actions. Finally, several medication switchers have been reported [28]. Failure to take account of switchers could distort estimates of adherence and persistence.

Poor persistence and adherence to osteoporosis medications reduce the potential benefits of osteoporosis therapy, lowering gains in bone mineral density and resulting in increased risk of fragility fractures [29]. Two systematic reviews and meta-analyses assessed the impact of adherence/persistence to oral bisphosphonates on fracture risk. The meta-analysis of Ross et al. [30] indicated that fracture risk increased by approximately 30% with non-adherence (defined as medical possession ratio $<$ 80%) and by 30% to 40% with non-persistence. Similarly, Imaz et al. [31] reported that low adherence to oral bisphosphonates was significantly associated with increased non-vertebral fracture risk (relative risk (RR) of 1.16), increased hip fracture risk (RR 1.28) and increased vertebral fracture risk (RR 1.43).

Studies reporting adherence to osteoporosis medication and its relationship with fracture risk mainly focused on oral bisphosphonates, although data for other medications (e.g.

teriparatide, raloxifene, denosumab or zoledronic acid) also revealed suboptimal levels [32, 33]. In a large US study, persistence to teriparatide and denosumab at one year were estimated at 69% and 58% [34]. Non-persistence to denosumab could further lead to important health problems as discontinuation with denosumab has recently been shown to be associated with rapid bone loss and increased risk of multiple vertebral fractures [35]. Persistence to yearly intravenous injection of zoledronic acid is also not convincing, as suggested by a study that showed that only one third of patients agreed to a second administration after one year [36]. Furthermore, for some osteoporosis medications, persistence to generic formulations has been shown to be poorer than for branded formulations [37, 38].

A few model-based studies have estimated the economic consequences of non-adherence at a population level [2, 39–42], suggesting important clinical and/or economic implications of poor adherence/persistence with osteoporosis medications. For example, poor adherence with osteoporosis medications resulted in about 50% reduction in the potential benefits observed in clinical trials (in terms of fractures prevented and quality-adjusted life-years (QALY)) and a substantial deterioration of the cost-effectiveness resulting from these medications [2, 40]. Medication adherence has further become an important factor for inclusion in cost-effectiveness analyses in osteoporosis [43–45]. Interestingly, economic studies have also suggested that interventions to improve adherence may likely confer cost-effectiveness benefits [43]. Improving adherence to medication could lead to greater benefits than designing a new more effective drug [46].

Determinants of non-adherence

Adherence is a complex multidimensional phenomenon determined by the interplay of several factors. Numerous determinants of non-adherence have been identified in the literature [12, 47]. The World Health Organization has classified these factors into five main categories, i.e. patient-related, therapy-related, condition-related, health system and socio-economic factors [5]. Recently, Yeam et al. [12] conducted a systematic literature review up to January 2018 to review and identify factors that influence patients' adherence to anti-osteoporotic therapy. A total of 24 factors and 139 sub-factors were identified from 124 relevant studies. The authors presented types and number of studies that presented the case for and against each specific factor and revealed that patient-related factors were the most commonly studied category followed by therapy-related and condition-related category.

Condition-related factors that were associated with poorer medication adherence included polypharmacy and having gastrointestinal diseases. History of falls, fractures and screening for osteoporosis were however not associated with higher adherence. This is important as patients with a recent fracture are at imminent risk of a further fracture [48]. *Patient-related factors* associated with poorer medication adherence included male gender, lower education levels, misconceptions about osteoporosis, lack of perceived benefits of therapy, whereas higher age was associated with higher medication adherence but only in half of the studies. Among *therapy-related factors*, medication side effects, complex instructions for medication administration and complex medication regimens were associated with poorer adherence; lower dosing frequency with higher adherence, while a history of anti-osteoporosis treatment

was not found to be a predictor of adherence. *Health system-based factors* associated with poorer medication adherence included care under different medical specialties and lack of patient education (information sharing, counselling from healthcare professional, etc.). *Socio-economic-related factors* associated with poorer medication adherence included current smoking, lower income level and lack of medical insurance coverage [12].

This review did not classify the factors according to adherence level, i.e. initiation, implementation and persistence. In another review of systematic reviews that covered 19 disease areas [47], a total of 771 individual factors of medication adherence were identified of which most were determinants of implementation, and only 47 determinants of persistence with medication and no determinant were specifically provided for initiation. In the field of osteoporosis, the patient perceived need for treatment, patient knowledge, bone density testing, improved patient-provider relationship, hospitalizations and prescription use have been shown to be positively associated with initiating osteoporotic therapy [49–52]. Reasons for not initiating osteoporotic therapy also include lack of motivation, concern about side-effects and medication costs [52]. Reasons for poor implementation and persistence with osteoporosis therapy include side effects of treatment, difficulties with medication intake (e.g. 30 minutes before breakfast), inconvenient dosing regimen, concern about treatment, no perceived benefits, drug cost, misinformation, insufficient motivation or dissatisfaction with their doctor visits [47, 53–55]. Adherence to osteoporosis medications is simply a matter of patient choice, and is due mostly to deliberative choice [56]. However, reasons for non-adherence can also be unintentional, resulting from forgetfulness, especially in holiday period [53] and some irrational behaviour could also be reported. Factors involved in treatment initiation may further differ than factors involved in persistence or implementation. As example, polypharmacy has been shown to have a positive effect on treatment initiation with patients who have not previously taken medication being less likely to start [14] while persistence and implementation are generally lower in patients on multiple medications [12]. Lau et al. conducted focus groups in Canada to understand reasons for non-adherence and included different patients' quotes regarding belief in the importance of taking medications for osteoporosis, medication-specific factors, beliefs regarding medications and health, relationships with health care providers, information exchange [57].

The patient participating at our working group stressed the importance of giving a positive message to the patient, and not exaggerating the consequences if the patient does not take the drug. Web and social medias nowadays also attract a lot of attention and could further negatively influence adherence to treatment.

Reasons for non-adherence are therefore numerous and multidimensional for the same patient. Each patient's reason(s) for non-adherence is different, depending on the interplay of these multiple factors, and could also change over time, and be different for each key element of adherence. Medication adherence is therefore not predictable, suggesting the need for individualized solutions.

Review of interventions

In 2012, Hiligsmann et al. [7] reviewed and critically appraised interventions to improve adherence and persistence to osteoporosis medications. A total of 20 studies tested a patient adherence intervention and reported quantitative results on adherence. Education programs (e.g. written materials, counselling, motivational interviewing, combination) was the most frequent intervention (n=9). Although patient education improved medication adherence in four studies, two large-scale randomized studies reported no benefits [58, 59]. Monitoring/supervision by bone mineral density or bone turnover markers was not shown to be associated with improved adherence. However, a positive message revealing a good bone turnover marker response was associated with a significant improvement in persistence [60]. Simplification of dosing regimens (with and without patient support program), electronic prescription and a pharmacist intervention were associated with improved adherence but only in couple of studies.

For the purpose of this working group, the previous systematic review was updated. A description of the literature search, data and critical appraisal can be found in the article of Cornelissen et al. [8] A total of 15 studies were identified between June 2012 and December 2018. Very few high-quality studies (such as RCTs) were identified and interventions were mainly single component intervention. Interventions were classified as patient education and/or support (n=9), monitoring and supervision (n=2), flexible dosing regimen and patient support (n=3), and interdisciplinary collaboration (n=1). In each subtype of interventions, mixed results on adherence/persistence were found where some studies were shown to be successful and others not (see next section). Interventions which include counselling with education and/or flexible dosing regimens seem to be most effective, as well as interventions aiming on testing and initiating medication.

Recommendations for practice

Many factors affect medication adherence. Single interventions could work but certainly not for all patients. There is not one single intervention to manage adherence. Some recommendations to improve adherence are made below.

It is of primary importance to emphasize the need for treatment adherence and to talk about adherence with patients. To adequately manage medication adherence, at least three stages are recommended:

1. *What is the adherence problem (initiation, implementation and/or persistence)?* It is key to identify the element(s) of non-adherence problem(s) that you want to address with the patient(s).
2. *How to measure adherence?* Different methods are available to measure the elements of medication adherence. Prescription refills and databases are appropriate methods to measure initiation and discontinuation of therapy [61] while electronic monitoring is the preferred method for implementation, although could be difficult to set up in real-life settings [61]. The use of self-reports or pill counts could be alternative methods.

3. *What is the reason and how to fix it?* Given the numerous potential factors for non-adherence, it is therefore important to understand the reason(s) for non-adherence for each patient and to find an adequate personalized solution.

These three steps are in line with the Six Sigma framework to manage medication adherence developed by Vrijens et al. [62]. This framework includes definition (robust taxonomy), measure (operational definition and performance), analyse (understand the cause), address the cause and implement solutions, and measurement-guided adherence management.

In addition to the updated systematic review of adherence interventions in osteoporosis, a comprehensive list of interventions is provided with evidence about their effectiveness including systematic reviews of adherence interventions in other diseases including depression, cardiovascular diseases and hypercholesterolaemia [63–65] (Table 1). Patient education and counselling, adherence monitoring with feedback, and dose simplification including flexible dosing regimen were associated with higher adherence improvements. We however acknowledge that the efficacy of these actions is still largely unknown and current data could be controversial.

Patient education and counselling were shown to improve post-fracture care and treatment initiation, with more controversial results on medication adherence. As a prominent example for post-fracture care, the PREVOST trial suggested that repeated oral and written information about fragility fractures and osteoporosis management by a case manager increased treatment initiation (53% initiated post-fracture care in the intervention compared to 33% in the control) [66]. In another study [67], patient education and referral to endocrinologist by a nurse was shown to improve the initiation of calcium and vitamin D, although up to 50% of patients with osteoporosis did not complete follow-up visits and/or did not adhere to treatment recommendations for osteoporosis. Assigning a screening coordinator to identify, educate, and follow up with fragility fracture patients and inform their physicians of the need to evaluate bone health was also shown to increase treatment initiation [68]. A systematic screening using FRAX® was also shown to increase use of, and adherence to anti-osteoporosis medications in the UK SCOOP Trial [69]. In addition, an osteoporosis school program (i.e. four classes of 8-12 participants over four weeks), peer-led community education and mentorship program or patient education program were associated with improved knowledge of osteoporosis and initiation of treatment [70]. In a large pragmatic randomized controlled trial, however, telephonic motivational interviewing intervention was not associated with significant improvements in medication adherence [59].

Adherence monitoring with feedback was also associated with improved adherence. Stuurman-Bieze et al. [71] showed that a proactive pharmaceutical care including counseling at baseline and at 2 weeks, and an active monitoring and counseling every 3 months by pharmacists for patients who should have redeemed a new prescription lead to improved adherence. Ducomlombier et al. [72] further suggests that phone calls by medical secretaries every 2 months to motivate patients to maintain good adherence to the treatment, to detect any difficulties in adherence with the prescription using non-incriminating questions was associated with improved adherence.

Dose simplification including flexible dosing regimen was associated with improved adherence. Offering patients a medication with less strict administration instructions such as the use of gastro-resistant risedronate tablets that could be taken after breakfast was associated with improved persistence to treatment [73]. Longer dosing regimen (such as six-month subcutaneous injection of denosumab or yearly intravenous injection of zoledronic acid) can also be interesting to improve adherence, although adherence levels have also been disappointed and far from optimal (see previously). Finally, a flexible dosing regimen (before breakfast; in-between meals; before bedtime) was also shown to be associated with improved persistence, although there was no statistical difference in terms of adherence [74].

Improving *patient interaction and shared-decision making* can also lead to improved treatment initiation, and potentially improve adherence. Several guidelines and international groups recommend that shared decision-making be part of standard treatment. In shared decision-making, both parties share information: the clinician offers options and describes their risks and benefits, and the patient expresses his or her preferences and values [75]. Each participant is thus armed with a better understanding of the relevant factors and shares responsibility in the decision about how to proceed [75]. Decision aids emphasize shared-decision making and include several features to support individualized treatment discussions like those needed for fracture prevention. Decision aids for a range of conditions have been shown to decrease decisional conflict, increase knowledge, and (when probabilities are included in the aid) improve the accuracy of risk perception [76]. Some decision aids aiming to facilitate shared-decision making in osteoporosis have been designed and shown to improve the quality of clinical decisions about bisphosphonate therapy, and may have improved adherence [77–79].

Providing patients with good quality information about osteoporosis, their risk and treatment options is needed as a crucial step to increase the diagnosis and treatment of osteoporosis. The *doctor* is therefore a key actor to improve medication adherence. As an educator and a partner in making treatment decisions, the doctor should be familiar with the medical evidence, able to discuss complex medical information in a manner that is understood by the patient and appropriately consider the patient's expectations, beliefs, and concerns [80, 81]. A doctor should thus facilitate interaction with the patient and him/her, understand the patient's needs and investigate the type of treatment appropriate for the patients (e.g. dosing regimen).

Recommendations for further research

Some areas where further research is needed were identified. First, it is important to improve our understanding on what osteoporotic patients want, need and prefer, and to better understand how and why patients will (not) be adherent. It is particularly important to understand the root causes for each aspect of adherence, i.e. initiation, implementation and persistence. Qualitative research is the best way to understand patients' needs and reasons for non-adherence, and should be a starting point for designing appropriate interventions. Stated-preference methods could also be interesting to reveal patients' preferences and important treatment characteristics [82].

Second, register-based studies can be attractive to investigate the importance of different factors and identify the strongest and most prevalent risk factors for poor adherence as targets for intervention. Big (pharmaco-epidemiologic) data provide new opportunities to help understand the reasons of medication non-adherence, and may also prove helpful in assessing the effects of new initiatives in the area.

Third, with a better understanding of patients' needs and non-adherence factors, it would then be possible to design interventions that fit patients' needs and wishes on a personalized manner. It is indeed recommended to develop complex, individualized and multi-factorial interventions. There is not one single intervention to manage adherence. Involving patients in the preparation, development and assessment of these complex interventions would be crucial [83]. Further research is also warranted on patient involvement in medication adherence research and patient empowerment and its role in promoting adherence. Complex interventions would thus need to be tested in large scale randomized controlled trials.

Fourth, as the cost of an adherence intervention may be high, it would also be important to identify patients that will be receptive to interventions and spend our limited resources on those patients. We further need patient outcome tools to better understand the patient perspective and patient biases in decision-making concerning osteoporosis therapies [84].

In addition, web-based applications could play a crucial role in supporting patients to improve their adherence to medication. Further research focusing on digital adherence interventions is necessary to determine their value and ideal ways to improve osteoporosis medication adherence.

Other areas for further research include the need to adequately communicate about osteoporosis, fracture risk and osteoporosis treatment to patients; development of shared-decision making; assessment of adherence-management model (such as the three stage model proposed previously); innovative methods for monitoring medication adherence and policy/health care systems initiatives to support adherence interventions.

Conclusions

Osteoporosis represents a significant healthcare burden in European countries which, due to increasing life expectancies, is predicted to increase further in the future [24]. Despite the increasing burden and the availability of effective treatments in reducing the risk of fractures, most patients are not taking their medication appropriately, or do not even start an osteoporosis medication. Improving treatment initiation and adherence to therapy is therefore urgently needed to leverage in full the benefits of drug therapy.

Poor and non-adherence with medication is common with a treatment gap estimated between 25% and 95% in European countries [25] and about half of the patients discontinuing therapy within one year. Reasons for non-adherence are numerous, diverse and multidimensional, depending on the interplay of multiple factors and may be different for each key element of adherence. Few single interventions have been shown to improve adherence or persistence to osteoporosis treatment. Potentially promising interventions include patient education with counselling, adherence monitoring with feedback and dose

simplification including flexible dosing regimen. Recommendations for practice and further research were provided, suggesting the need to emphasize the importance of adherence to treatment and to talk about adherence with patients, and the need to implement sound measures of adherence so that the intervention can be individualized. These recommendations are intended for clinicians to manage adherence of their patients, and to researchers and policy makers to design, facilitate and appropriately use adherence interventions, and to advance research in the field.

Acknowledgements

The meeting was funded by the European Society for Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases (ESCEO). The authors are grateful to the Prince Mutaib Chair for Biomarkers of Chronic Disease, King Saud University, Riyadh, KSA and the WHO Collaborating Centre for Public Health Aspects of Musculoskeletal Health and Aging, for its support.

References

1. Kanis JA, Cooper C, Rizzoli R, Reginster JY, Scientific Advisory Board of the European Society for C, Economic Aspects of O. et al. European guidance for the diagnosis and management of osteoporosis in postmenopausal women. *Osteoporos Int.* 2019; 30(1):3–44. [PubMed: 30324412]
2. Hiligsmann M, McGowan B, Bennett K, Barry M, Reginster JY. The clinical and economic burden of poor adherence and persistence with osteoporosis medications in Ireland. *Value Health.* 2012; 15(5):604–12. [PubMed: 22867768]
3. Jha S, Wang Z, Laucis N, Bhattacharyya T. Trends in Media Reports, Oral Bisphosphonate Prescriptions, and Hip Fractures 1996-2012: An Ecological Analysis. *J Bone Miner Res.* 2015; 30(12):2179–87. [PubMed: 26018247]
4. Kanis JA, Cooper C, Rizzoli R, Abrahamsen B, Al-Daghri NM, Brandi ML, et al. Identification and management of patients at increased risk of osteoporotic fracture: outcomes of an ESCEO expert consensus meeting. *Osteoporos Int.* 2017; 28(7):2023–34. [PubMed: 28451733]
5. Sabate, E, editor. *Adherence to Long-Term Therapies: Evidence for Action.* Geneva: World Health Organization; 2003.
6. Gold DT, Silverman S. Review of adherence to medications for the treatment of osteoporosis. *Curr Osteoporos Rep.* 2006; 4(1):21–7. [PubMed: 16527004]
7. Hiligsmann M, Salas M, Hughes DA, Manias E, Gwady-Sridhar FH, Linck P, et al. Interventions to improve osteoporosis medication adherence and persistence: a systematic review and literature appraisal by the ISPOR Medication Adherence & Persistence Special Interest Group. *Osteoporosis Int.* 2013; 24(12):2907–18.
8. Cornelissen D, et al. Interventions to improve osteoporosis medication adherence and persistence: an updated systematic review and literature appraisal.
9. Vrijens B, De Geest S, Hughes DA, Przemyslaw K, Demonceau J, Ruppar T, et al. A new taxonomy for describing and defining adherence to medications. *Br J Clin Pharmacol.* 2012; 73(5):691–705. [PubMed: 22486599]
10. De Geest S, Zullig LL, Dunbar-Jacob J, Helmy R, Hughes DA, Wilson IB, et al. ESPACOMP Medication Adherence Reporting Guideline (EMERGE). *Ann Intern Med.* 2018; 169(1):30–5. [PubMed: 29946690]
11. Lekkerkerker F, Kanis JA, Alsayed N, Bouvenot G, Burlet N, Cahall D, et al. Adherence to treatment of osteoporosis: a need for study. *Osteoporos Int.* 2007; 18(10):1311–7. [PubMed: 17585359]
12. Yeam CT, Chia S, Tan HCC, Kwan YH, Fong W, Seng JJB. A systematic review of factors affecting medication adherence among patients with osteoporosis. *Osteoporosis Int.* 2018; 29(12): 2623–37.

13. Fischer MA, Stedman MR, Lii J, Vogeli C, Shrank WH, Brookhart MA, et al. Primary medication non-adherence: analysis of 195,930 electronic prescriptions. *J Gen Intern Med.* 2010; 25(4):284–90. [PubMed: 20131023]
14. Reynolds K, Muntner P, Cheetham TC, Harrison TN, Morisky DE, Silverman S, et al. Primary non-adherence to bisphosphonates in an integrated healthcare setting. *Osteoporos Int.* 2013; 24(9): 2509–17. [PubMed: 23595561]
15. Rabenda V, Reginster JY. Overcoming problems with adherence to osteoporosis medication. *Expert Rev Pharmacoecon Outcomes Res.* 2010; 10(6):677–89. [PubMed: 21155701]
16. Rabenda V, Hiligsmann M, Reginster JY. Poor adherence to oral bisphosphonate treatment and its consequences: a review of the evidence. *Expert Opin Pharmacother.* 2009; 10(14):2303–15. [PubMed: 19640210]
17. Strom O, Borgstrom F, Kanis JA, Compston J, Cooper C, McCloskey EV, et al. Osteoporosis: burden, health care provision and opportunities in the EU: a report prepared in collaboration with the International Osteoporosis Foundation (IOF) and the European Federation of Pharmaceutical Industry Associations (EFPIA). *Arch Osteoporos.* 2011; 6:59–155. [PubMed: 22886101]
18. Landfeldt E, Strom O, Robbins S, Borgstrom F. Adherence to treatment of primary osteoporosis and its association to fractures—the Swedish Adherence Register Analysis (SARA). *Osteoporos Int.* 2012; 23(2):433–43. [PubMed: 21286686]
19. Blaschke TF, Osterberg L, Vrijens B, Urquhart J. Adherence to medications: insights arising from studies on the unreliable link between prescribed and actual drug dosing histories. *Annu Rev Pharmacol Toxicol.* 2012; 52:275–301. [PubMed: 21942628]
20. Rabenda V, Mertens R, Fabri V, Vanoverloop J, Sumkay F, Vannecke C, et al. Adherence to bisphosphonates therapy and hip fracture risk in osteoporotic women. *Osteoporos Int.* 2008; 19(6): 811–8. [PubMed: 17999022]
21. van Boven JFM, de Boer PT, Postma MJ, Vegter S. Persistence with osteoporosis medication among newly-treated osteoporotic patients. *J Bone Miner Metab.* 2013; 31(5):562–70. [PubMed: 23575910]
22. Roerholt C, Eiken P, Abrahamsen B. Initiation of anti-osteoporotic therapy in patients with recent fractures: a nationwide analysis of prescription rates and persistence. *Osteoporos Int.* 2009; 20(2): 299–307. [PubMed: 18551241]
23. Jacob L, Dreher M, Kostev K, Hadji P. Increased treatment persistence and its determinants in women with osteoporosis with prior fracture compared to those without fracture. *Osteoporos Int.* 2016; 27(3):963–9. [PubMed: 26519418]
24. Svedbom A, Hernlund E, Ivergard M, Compston J, Cooper C, Stenmark J, et al. Osteoporosis in the European Union: a compendium of country-specific reports. *Arch Osteoporos.* 2013; 8:137. [PubMed: 24113838]
25. Kanis JA, Svedbom A, Harvey N, McCloskey EV. The osteoporosis treatment gap. *J Bone Miner Res.* 2014; 29(9):1926–8. [PubMed: 24956507]
26. Blaschke TF, Osterberg L, Vrijens B, Urquhart J. Adherence to Medications: Insights Arising from Studies on the Unreliable Link Between Prescribed and Actual Drug Dosing Histories. *Annu Rev Pharmacol.* 2012; 52:275.
27. Balasubramanian A, Brookhart MA, Goli V, Critchlow CW. Discontinuation and reinitiation patterns of osteoporosis treatment among commercially insured postmenopausal women. *Int J Gen Med.* 2013; 6:839–48. [PubMed: 24235846]
28. Wade SW, Satram-Hoang S, Nadkar A, Macarios D, Tosteson AN. Impact of medication adherence on health care utilization and productivity: self-reported data from a cohort of postmenopausal women on osteoporosis therapy. *Clin Ther.* 2011; 33(12):2006–15. [PubMed: 22099387]
29. Siris ES, Selby PL, Saag KG, Borgstrom F, Herings RM, Silverman SL. Impact of osteoporosis treatment adherence on fracture rates in North America and Europe. *Am J Med.* 2009; 122(2 Suppl):S3–13.
30. Ross S, Samuels E, Gairy K, Iqbal S, Badamgarav E, Siris E. A meta-analysis of osteoporotic fracture risk with medication nonadherence. *Value Health.* 2011; 14(4):571–81. [PubMed: 21669382]

31. Imaz I, Zegarra P, Gonzalez-Enriquez J, Rubio B, Alcazar R, Amate JM. Poor bisphosphonate adherence for treatment of osteoporosis increases fracture risk: systematic review and meta-analysis. *Osteoporos Int.* 2010; 21(11):1943–51. [PubMed: 19967338]
32. Kyvernitakis I, Kostev K, Kurth A, Albert US, Hadji P. Differences in persistency with teriparatide in patients with osteoporosis according to gender and health care provider. *Osteoporos Int.* 2014; 25(12):2721–8. [PubMed: 25011986]
33. Hadji P, Papaioannou N, Gielen E, Feudjo Tepie M, Zhang E, Frieling I, et al. Persistence, adherence, and medication-taking behavior in women with postmenopausal osteoporosis receiving denosumab in routine practice in Germany, Austria, Greece, and Belgium: 12-month results from a European non-interventional study. *Osteoporos Int.* 2015; 26(10):2479–89. [PubMed: 26018090]
34. Cheng LI, Durden E, Limone B, Radbill L, Juneau PL, Spangler L, et al. Persistence and Compliance with Osteoporosis Therapies Among Women in a Commercially Insured Population in the United States. *J Manag Care Spec Ph.* 2015; 21(9):824–U322.
35. McClung MR, Wagman RB, Miller PD, Wang A, Lewiecki EM. Observations following discontinuation of long-term denosumab therapy. *Osteoporos Int.* 2017; 28(5):1723–32. [PubMed: 28144701]
36. Lee YK, Nho JH, Ha YC, Koo KH. Persistence with intravenous zoledronate in elderly patients with osteoporosis. *Osteoporosis Int.* 2012; 23(9):2329–33.
37. Ringe JD, Moller G. Differences in persistence, safety and efficacy of generic and original branded once weekly bisphosphonates in patients with postmenopausal osteoporosis: 1-year results of a retrospective patient chart review analysis. *Rheumatol Int.* 2009; 30(2):213–21. [PubMed: 19430791]
38. Kanis JA, Reginster JY, Kaufman JM, Ringe JD, Adachi JD, Hiligsmann M, et al. A reappraisal of generic bisphosphonates in osteoporosis. *Osteoporos Int.* 2012; 23(1):213–21. [PubMed: 21953472]
39. Hiligsmann M, Rabenda V, Gathon HJ, Ethgen O, Reginster JY. Potential Clinical and Economic Impact of Nonadherence with Osteoporosis Medications. *Calcif Tissue Int.* 2010; 86(3):202–10.
40. Hiligsmann M, Rabenda V, Bruyere O, Reginster JY. The clinical and economic burden of non-adherence with oral bisphosphonates in osteoporotic patients. *Health Policy.* 2010; 96(2):170–7. [PubMed: 20153543]
41. Danese MD, Badamgarav E, Bauer DC. Effect of adherence on lifetime fractures in osteoporotic women treated with daily and weekly bisphosphonates. *J Bone Miner Res.* 2009; 24(11):1819–26. [PubMed: 19419313]
42. Strom O, Borgstrom F, Kanis J, Jonsson B. Incorporating adherence into health economic modelling of osteoporosis. *Osteoporosis Int.* 2009; 20(1):23–34.
43. Hiligsmann M, Boonen A, Rabenda V, Reginster JY. The importance of integrating medication adherence into pharmaco-economic analyses: the example of osteoporosis. *Expert Rev Pharmacoecon Outcomes Res.* 2012; 12(2):159–66. [PubMed: 22458617]
44. Hiligsmann M, Reginster JY, Tosteson ANA, Bukata SV, Saag KG, Gold DT, et al. Recommendations for the conduct of economic evaluations in osteoporosis: outcomes of an experts' consensus meeting organized by the European Society for Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases (ESCEO) and the US branch of the International Osteoporosis Foundation. *Osteoporos Int.* 2019; 30(1):45–57.
45. Landfeldt E, Strom O, Robbins S, Borgstrom F. Adherence to treatment of primary osteoporosis and its association to fractures—the Swedish Adherence Register Analysis (SARA). *Osteoporosis Int.* 2012; 23(2):433–43.
46. Burkhart PV, Sabate E. Adherence to long-term therapies: Evidence for action. *J Nurs Scholarship.* 2003; 35(3):207.
47. Kardas P, Lewek P, Matyjaszczuk M. Determinants of patient adherence: a review of systematic reviews. *Front Pharmacol.* 2013; 4
48. Johansson H, Siggeirsdottir K, Harvey NC, Oden A, Gudnason V, McCloskey E, et al. Imminent risk of fracture after fracture. *Osteoporos Int.* 2017; 28(3):775–80. [PubMed: 28028554]
49. Beaton DE, Dyer S, Jiang D, Sujic R, Slater M, Sale JE, et al. Factors influencing the pharmacological management of osteoporosis after fragility fracture: results from the Ontario

- Osteoporosis Strategy's fracture clinic screening program. *Osteoporos Int.* 2014; 25(1):289–96. [PubMed: 23794044]
50. Majumdar SR, McAlister FA, Johnson JA, Weir DL, Bellerose D, Hanley DA, et al. Critical impact of patient knowledge and bone density testing on starting osteoporosis treatment after fragility fracture: secondary analyses from two controlled trials. *Osteoporos Int.* 2014; 25(9):2173–9. [PubMed: 24803330]
 51. Morell S, Hemmeler C, Amsler F, Gross T. Adherence to osteoporosis pharmacotherapy one year after osteoporotic fracture - a Swiss trauma center secondary prevention project. *Swiss Med Wkly.* 2017; 147:w14451. [PubMed: 28695555]
 52. Yu J, Brenneman SK, Sazonov V, Modi A. Reasons for not initiating osteoporosis therapy among a managed care population. *Patient Prefer Adherence.* 2015; 9:821–30. [PubMed: 26150702]
 53. Tafaro L, Nati G, Leoni E, Baldini R, Cattaruzza MS, Mei M, et al. Adherence to anti-osteoporotic therapies: role and determinants of "spot therapy". *Osteoporos Int.* 2013; 24(8):2319–23. [PubMed: 23404614]
 54. Iversen MD, Vora RR, Servi A, Solomon DH. Factors affecting adherence to osteoporosis medications: a focus group approach examining viewpoints of patients and providers. *J Geriatr Phys Ther.* 2011; 34(2):72–81. [PubMed: 21937896]
 55. Papaioannou A, Kennedy CC, Dolovich L, Lau E, Adachi JD. Patient adherence to osteoporosis medications: problems, consequences and management strategies. *Drugs Aging.* 2007; 24(1):37–55. [PubMed: 17233546]
 56. Silverman SL, Schousboe JT, Gold DT. Oral bisphosphonate compliance and persistence: a matter of choice? *Osteoporos Int.* 2011; 22(1):21–6. [PubMed: 20458571]
 57. Lau E, Papaioannou A, Dolovich L, Adachi J, Sawka AM, Burns S, et al. Patients' adherence to osteoporosis therapy - Exploring the perceptions of postmenopausal women. *Can Fam Physician.* 2008; 54(3):394–402. [PubMed: 18337534]
 58. Shu ADH, Stedman MR, Polinski JM, Jan SA, Patel M, Truppo C, et al. Adherence to Osteoporosis Medications After Patient and Physician Brief Education: Post Hoc Analysis of a Randomized Controlled Trial. *Am J Manag Care.* 2009; 15(7):417–24. [PubMed: 19589009]
 59. Solomon DH, Iversen MD, Avorn J, Gleeson T, Brookhart MA, Patrick AR, et al. Osteoporosis telephonic intervention to improve medication regimen adherence: a large, pragmatic, randomized controlled trial. *Arch Intern Med.* 2012; 172(6):477–83. [PubMed: 22371876]
 60. Delmas PD, Vrijens B, Eastell R, Roux C, Pols HA, Ringe JD, et al. Effect of monitoring bone turnover markers on persistence with risedronate treatment of postmenopausal osteoporosis. *J Clin Endocrinol Metab.* 2007; 92(4):1296–304. [PubMed: 17244788]
 61. Vrijens B, Heidbuchel H. Non-vitamin K antagonist oral anticoagulants: considerations on once- vs. twice-daily regimens and their potential impact on medication adherence. *Europace.* 2015; 17(4):514–23. [PubMed: 25694538]
 62. Vrijens B. A Six Sigma framework to successfully manage medication adherence. *Br J Clin Pharmacol.* 2019
 63. Rash JA, Campbell DJT, Tonelli M, Campbell TS. A systematic review of interventions to improve adherence to statin medication: What do we know about what works? *Prev Med.* 2016; 90:155–69. [PubMed: 27413005]
 64. van Dalem J, Krass I, Aslani P. Interventions promoting adherence to cardiovascular medicines. *Int J Clin Pharm-Net.* 2012; 34(2):295–311.
 65. Chong WW, Aslani P, Chen TF. Effectiveness of interventions to improve antidepressant medication adherence: a systematic review. *Int J Clin Pract.* 2011; 65(9):954–75. [PubMed: 21849010]
 66. Merle B, Chapurlat R, Vignot E, Thomas T, Haesebaert J, Schott AM. Post-fracture care: do we need to educate patients rather than doctors? The PREVOST randomized controlled trial. *Osteoporosis Int.* 2017; 28(5):1549–58.
 67. Seuffert P, Sagebien CA, McDonnell M, O'Hara DA. Evaluation of osteoporosis risk and initiation of a nurse practitioner intervention program in an orthopedic practice. *Archives of Osteoporosis.* 2016; 11(1)

68. Beaton DE, Mamdani M, Zheng H, Jaglal S, Cadarette SM, Bogoch ER, et al. Improvements in osteoporosis testing and care are found following the wide scale implementation of the Ontario Fracture Clinic Screening Program An interrupted time series analysis. *Medicine*. 2017; 96(48)
69. Parsons C, Harvey N, Shepstone L, Kanis J, Lenaghan E, Clarke S, et al. Systematic Screening Using Frax Leads to Increased Use of, and Adherence to, Anti-Osteoporosis Medications: The Uk Scoop Trial. *Osteoporosis Int*. 2018; 29:613.
70. Nielsen D, Ryg J, Nielsen W, Knold B, Nissen N, Brixen K. Patient education in groups increases knowledge of osteoporosis and adherence to treatment: A two-year randomized controlled trial. *Patient Educ Couns*. 2010; 81(2):155–60. [PubMed: 20400258]
71. Stuurman-Bieze AGG, Hiddink EG, van Boven JFM, Vegter S. Proactive pharmaceutical care interventions decrease patients' nonadherence to osteoporosis medication. *Osteoporosis Int*. 2014; 25(6):1807–12.
72. Ducoulombier V, Luraschi H, Forzy G, Vandecandelaere M, Houvenagel E. Contribution of Phone Follow-up to Improved Adherence to Oral Osteoporosis Treatment. *Am J Pharm Benefist*. 2015; 7(3):E81–E9.
73. Hiligsmann M, Reginster JY. Cost-effectiveness of gastro-resistant risedronate tablets for the treatment of postmenopausal women with osteoporosis in France. *Osteoporosis Int*. 2019; 30(3): 649–58.
74. Oral A, Lorenc R, Investigators F-AS. Compliance, persistence, and preference outcomes of postmenopausal osteoporotic women receiving a flexible or fixed regimen of daily risedronate: A multicenter, prospective, parallel group study. *Acta Orthop Traumatol Turc*. 2015; 49(1):67–74. [PubMed: 25803256]
75. Barry MJ, Edgman-Levitan S. Shared decision making--pinnacle of patient-centered care. *N Engl J Med*. 2012; 366(9):780–1. [PubMed: 22375967]
76. Stacey D, Legare F, Lewis K, Barry MJ, Bennett CL, Eden KB, et al. Decision aids for people facing health treatment or screening decisions. *Cochrane Db Syst Rev*. 2017; (4)
77. Montori VM, Shah ND, Pencille LJ, Branda ME, Van Houten HK, Swiglo BA, et al. Use of a Decision Aid to Improve Treatment Decisions in Osteoporosis: The Osteoporosis Choice Randomized Trial. *Am J Med*. 2011; 124(6):549–56. [PubMed: 21605732]
78. Hiligsmann M, Ronda G, van der Weijden T, Boonen A. The development of a personalized patient education tool for decision making for postmenopausal women with osteoporosis. *Osteoporosis Int*. 2016; 27(8):2489–96.
79. Smallwood AJ, Schapira MM, Fedders M, Neuner JM. A pilot randomized controlled trial of a decision aid with tailored fracture risk tool delivered via a patient portal. *Osteoporosis Int*. 2017; 28(2):567–76.
80. Lewiecki EM. Risk Communication and Shared Decision Making in the Care of Patients With Osteoporosis. *J Clin Densitom*. 2010; 13(4):335–45. [PubMed: 20663701]
81. Gasparik AI. Clinical Setting Influences Patterns of Interaction between Osteoporosis Patient and Physician. *Calcified Tissue Int*. 2014; 95(2):122–4.
82. Hiligsmann M, Pinto D, Dennison E, Al-Daghri N, Beaudart C, Branco J, et al. Patients' preferences for osteoarthritis treatment: the value of stated-preference studies. *Aging Clinical and Experimental Research*. 2019; 31(1):1–3. [PubMed: 30607878]
83. de Wit M, Cooper C, Tugwell P, Bere N, Kirwan J, Conaghan PG, et al. Practical guidance for engaging patients in health research, treatment guidelines and regulatory processes: results of an expert group meeting organized by the World Health Organization (WHO) and the European Society for Clinical and Economic Aspects of Osteoporosis, Osteoarthritis and Musculoskeletal Diseases (ESCEO). *Aging Clin Exp Res*. 2019; 31(7):905–915. [PubMed: 30993659]
84. Silverman S, Gold DT. Medication Decision-making in Osteoporosis: Can We Explain Why Patients Do Not Take Their Osteoporosis Medications? *Current Osteoporosis Reports*. 2018; 16(6):772–4. [PubMed: 30374627]

Mini-abstract

Many patients at increased risk of fractures do not take their medication appropriately, resulting in a substantial decrease in the benefits of drug therapy. Improving medication adherence is urgently needed but remains laborious, given the numerous and multidimensional reasons for non-adherence, suggesting the need for measurement-guided, multi-factorial and individualized solutions.

Table 1
Evidence about the effectiveness of adherence-enhancing interventions

Intervention subclass	OP studies	Other conditions*	Conclusion
Education and support			
Patient education program	4	4	Mixed evidence of effectiveness
Patient counselling	1	10	Strong evidence of effectiveness
Patient education combined with counselling	11	15	Strong evidence of effectiveness
Provision of educational material	2	5	No evidence of effectiveness
Monitoring supervision			
Non-adherence monitoring	1	2	Limited evidence of effectiveness
Adherence monitoring combined with counselling	5	10	Strong evidence of effectiveness
Drug regimen combined with counselling	1	1	Limited evidence of effectiveness
Reminders to take the medication	1	7	Strong evidence of effectiveness
Monitoring biomarkers	3	N/a	No evidence of effectiveness
Dose regimen adjustment and simplification			
Dose simplification including flexible dosing regimen	4	8	Strong evidence of effectiveness
Individual medication program	1	1	Limited evidence of effectiveness
Costs covered	N/a	1	Limited evidence of effectiveness
Other subclasses			
Combination of interventions mentioned above	3	2	Mixed evidence of effectiveness
Other interventions	4	9	Low evidence of effectiveness

N/a Not Applicable

* Restricted to depression, cardiovascular diseases and hypercholesterolaemia