



Published in final edited form as:

*J Biol Chem.* 2007 November 2; 282(44): 32208–32221. doi:10.1074/jbc.M704870200.

## REGULATION OF IRF-3 DEPENDENT INNATE IMMUNITY BY THE PAPAINE-LIKE PROTEASE DOMAIN OF THE SARS CORONAVIRUS

Santhana G. Devaraj<sup>1</sup>, Nan Wang<sup>1</sup>, Zhongbin Chen<sup>2</sup>, Zihong Chen<sup>1</sup>, Monica Tseng<sup>1</sup>, Naina Barretto<sup>2</sup>, Rongtuan Lin<sup>4</sup>, Clarence J. Peters<sup>1,3</sup>, Chien-Te K. Tseng<sup>1,3</sup>, Susan C. Baker<sup>2</sup>, and Kui Li<sup>1</sup>

<sup>1</sup> Department of Microbiology and Immunology, University of Texas Medical Branch, Galveston, TX

<sup>3</sup> Center of Biodefense and Emerging Infectious Diseases, Institute for Human Infections and Immunity, University of Texas Medical Branch, Galveston, TX

<sup>2</sup> Department of Microbiology and Immunology, Loyola University of Chicago, Maywood, IL

<sup>4</sup> Lady Davis Institute for Medical Research, Montreal, Canada

### Abstract

Severe acute respiratory syndrome coronavirus (SARS-CoV) is a novel coronavirus that causes a highly contagious respiratory disease, SARS, with significant mortality. While factors contributing to the highly pathogenic nature of SARS-CoV remain poorly understood, it has been reported that SARS-CoV infection does not induce type I interferons (IFNs) in cell culture. However, it is uncertain whether SARS-CoV evades host detection or has evolved mechanisms to counteract innate host defenses. We show here that infection of SARS-CoV triggers a weak IFN response in cultured human lung/bronchial epithelial cells without inducing the phosphorylation of IFN-regulatory factor 3 (IRF-3), a latent cellular transcription factor that is pivotal for type I IFN synthesis. Furthermore, SARS-CoV infection blocked the induction of IFN antiviral activity and the upregulation of protein expression of a subset of IFN-stimulated genes triggered by dsRNA or an unrelated paramyxovirus. In searching for a SARS-CoV protein capable of counteracting innate immunity, we identified the papain-like protease (PLpro) domain as a potent IFN antagonist. The inhibition of the IFN response does not require the protease activity of PLpro. Rather, PLpro interacts with IRF-3, and inhibits the phosphorylation and nuclear translocation of IRF-3, thereby disrupting the activation of type I IFN responses through either Toll-like receptor 3 or retinoic acid inducible gene I/melanoma differentiation-associated gene 5 pathways. Our data suggest that regulation of IRF-3-dependent innate antiviral defenses by PLpro may contribute to the establishment of SARS-CoV infection.

Upon virus infection, the host immediately launches an innate immune defense mechanism characterized by production of type I interferons (IFN- $\alpha$  and  $\beta$ ). IFN induces the expression of hundreds of IFN-stimulated genes (ISGs) that establish an antiviral state, thereby limiting viral replication and spread (1–5). This innate antiviral response is initiated upon host detection of viral pathogen-associated molecular patterns (PAMPs) via a number of cellular pattern recognition receptors (PRRs). Viral nucleic acids comprising viral genomes or generated during viral replication present major PAMPs that can be recognized by two different classes of PRRs, i.e., the membrane-bound Toll-like receptors (TLRs) or the caspase recruitment domain (CARD)-containing, cytoplasmic RNA helicases, retinoic acid-inducible gene I (RIG-I) or melanoma differentiation-associated gene 5 (MDA5) (6–8). Among these, TLR3 and

MDA5 detect viral double-stranded (ds) RNA in the endosomes and cytoplasm, respectively (9–11). Intracellular viral RNAs bearing 5' triphosphates are recognized by RIG-I (12,13). Upon engagement of their respective ligands, these PRRs recruit different adaptor molecules, relaying signals to downstream kinases that activate IFN regulatory factor-3 (IRF-3), nuclear factor kappa B (NF- $\kappa$ B), and ATF/C-Jun, transcription factors that coordinately regulate IFN- $\beta$  transcription (1,7). IRF-3 is a constitutively expressed, latent transcription factor that plays a pivotal role in type I IFN responses. Its activation requires specific C-terminal phosphorylation mediated by two non-canonical I $\kappa$ B kinases, TBK1 or IKK $\epsilon$  (14,15). Phosphorylation of IRF-3 leads to its homodimerization, nuclear translocation, and association with CBP/p300, whereupon it collaborates with activated NF- $\kappa$ B to induce IFN- $\beta$  synthesis (16,17). In addition, although it is not utilized by most parenchymal cell types, TLR7 constitutes the major pathway in plasmacytoid dendritic cells and triggers potent IFN- $\alpha$  production in circulation upon engagement of single-stranded (ss) viral RNAs (18,19). During co-evolution with their hosts, many viruses have acquired mechanisms to circumvent cellular IFN responses by encoding proteins that disrupt the IFN signaling pathways. It is thought that the early virus-host interactions may significantly impact the course and/or outcome of the infection (1,5).

Severe acute respiratory syndrome coronavirus (SARS-CoV) is a novel coronavirus that causes a highly contagious respiratory disease, SARS, with a significant mortality rate of about 10% (20). The first 3/4 of the positive-sense, single-stranded, 29.7-kb genome of SARS-CoV is translated into two large replicase polyproteins (22–24) called as pp1a and pp1ab (25,26). Papain-like protease (PLpro) and 3C-like protease domains present within these polyproteins direct their processing into 16 non-structural proteins (nsp1–16) that assemble to generate a multifunctional, membrane-associated replicase complex. Unlike other coronaviruses that encode two different PLpros, SARS-CoV encodes only one. Residing within the 213-kDa, membrane-associated replicase product nsp3, the SARS-CoV PLpro is responsible for cleaving junctions spanning nsp1 through nsp4 (27). Surprisingly, the crystal structure of SARS-CoV PLpro resembles that of known cellular deubiquitinating (DUB) enzymes USP14 and HAUSP (28). SARS-CoV PLpro has also been demonstrated to have both DUB and deISGylation activities *in vitro*, consistent with the fact that it shares the consensus recognition motif LXGG with cellular DUBs (29,30). Based on these findings, it has been proposed that SARS-CoV PLpro may have important but still uncharacterized functions in regulating cellular processes, in addition to its role as a viral protease (29,30).

While factors contributing to the highly pathogenic nature of SARS-CoV remain poorly understood, infection of HEK293 cells does not induce production of type I IFNs (21). However, pre-treatment of permissive cells with IFN prevents SARS-CoV replication (31). These are intriguing observations, as the ability to disrupt IFN responses seems to correlate with *in vivo* virulence in some viral infections, e.g., equine, swine, and H5N1 avian influenza viruses which express NS1, a well-characterized IFN antagonist (32–34). The absence of IRF-3 activation and IFN- $\beta$  mRNA induction in SARS-CoV-infected HEK293 cells has led to the proposal that SARS-CoV is able to disrupt the activation of IRF-3-mediated defenses (21). However, contradictory results have been published regarding whether SARS-CoV evades host detection, or alternatively, has developed strategies to actively block host innate defense mechanisms. Three SARS-CoV proteins, ORF3b, ORF6, and nucleocapsid (N), were recently identified as IFN antagonists, based on their ability to inhibit IRF-3 activation (35). SARS-CoV nsp1 was also shown to inhibit host IFN responses by inducing the degradation of host mRNAs (36). However, others have reported that infection of mouse hepatitis virus (MHV), a closely related type 2 coronavirus, or SARS-CoV does not activate IFN- $\beta$  synthesis in murine fibroblasts nor inhibit its induction by poly (I-C) or Sendai virus (SeV) (37,38). It was proposed from these studies that group 2 coronaviruses are “invisible” to host cells and thus, avoid innate immunity by somehow avoiding host recognition (37,38).

Here, we show that SARS-CoV triggers a weak, yet demonstrable IFN response in cultured human lung/bronchial epithelial cells and fibroblasts derived from fetal rhesus monkey kidney, both of which are naturally permissive to SARS-CoV. Furthermore, SARS-CoV infection blocks the induction of IFN antiviral activity and the upregulation of a subset of ISG proteins in response to dsRNA or SeV. We also identify the PLpro of SARS-CoV as a potent IFN antagonist. PLpro interacts with IRF-3, and inhibits its phosphorylation and nuclear translocation, thereby disrupting multiple signaling pathways leading to IFN induction. Finally, we demonstrate a role of IRF-3 signaling in control of SARS-CoV infection, by showing that disruption of IRF-3 function significantly upregulates SARS-CoV replication. Our data suggest that SARS-CoV is detected by the innate immune surveillance mechanism of host cells, but that active regulation of IRF-3-mediated host defenses limits the IFN responses and likely contributes to establishment of SARS-CoV infection.

## EXPERIMENTAL PROCEDURES

### Cells

Human bronchial epithelial cells Calu-3, African green monkey kidney cells MA104 and Vero, Human embryonic kidney (HEK) 293, 293T, 293-TLR3 (a gift from Kate Fitzgerald), HeLa, human hepatoma cells Huh7 were maintained by conventional techniques. HeLa Tet-Off cell lines (Clontech) were cultured following the manufacturer's instructions.

### Plasmids

To obtain high-level expression of SARS-CoV PLpro in eukaryotic cells, the codon usage of the SARS-CoV PLpro core domain (aa 1541–1855 of pp1a of SARS-CoV Urbani strain, GenBank ID AY278741) was optimized based on the human codon usage frequency, and the potential splicing sites and polyadenylation signal sequences were removed from the coding regions of PLpro. The optimized PLpro was named SARS PLpro Sol (See supporting online information), and cloned into pcDNA3.1-V5/HisB (Invitrogen) between the BamHI and EcoRI sites with in-frame fusion with C-terminal V5 and 6XHis tags, to result in the expression plasmid pcDNA3.1-SARS-PLpro-Sol. To generate pcDNA3.1-SARS-PLpro-TM (aa 1841–2425), the coding sequence for the TM domains downstream of PLpro in SARS-CoV nsp3 was amplified from the plasmid pPLpro-HD (27) by PCR with primers P1, 5'-TTAGAATTC ACCTGTGTCGTAATC-3' and P2, 5'-TAACTCGAGGTGAGTCTTGCAGAAGC-3', and the product was digested with EcoRI/XhoI and cloned downstream of PLpro Sol with C-terminal in-frame fusion of V5-6XHis tags. The catalytically inactive mutants of PLpro Sol and PLpro-TM (C1651A and D1826A) were generated by QuikChange site-directed mutagenesis (Stratagene) using specific primers containing the desired mutations (see supporting online information). The fidelity of the cDNA inserts in all constructs was validated by DNA sequencing. To construct a tetracycline (tet)-regulated expression plasmid for SARS-CoV PLpro-TM, cDNA fragment encoding PLpro-TM along with its downstream V5-6XHis tags in pcDNA3.1-SARS-PLpro-TM were excised by restriction digestion with HindIII and PmeI, and ligated into pTRE2Bla (39) that was digested with Hind III and EcoRV, to result in the plasmid pTRE2Bla-SARS-PLpro-TM. To construct a tet-regulated expression plasmid for a dominant negative (DN) form of IRF-3, cDNA encoding the aa 134–427 of IRF-3 was amplified from Huh7 cDNA by PCR and cloned into pTRE2Bla (39) between HindIII and XbaI sites, to generate pTRE2Bla-IRF3DN133.

The cDNA expression plasmids for TRIF, RIG-I, MAVS and bovine viral diarrhea virus (BVDV) Npro have been described (39–41,43). The following plasmids were kind gifts from the respectively indicated contributors: pcDNA3-A20-myc (from Nancy Raab-Traub) (42); p55C1Bluc, pEFBos N-RIG and pEFBos N-MDA5 (from Takashi Fujita) (6); pcDNA3-Flag TBK1 and pcDNA3-Flag IKK $\epsilon$  (from Kate Fitzgerald) (15); pIFN- $\beta$ -luc. IRF3-5D, GFP-IRF3

and GFP-IRF3 5D (from Rongtuan Lin) (17); PRDII-Luc (from Michael Gale); (PRDIII-I)4-Luc (from Christina Ehrhardt) (44).

### Generation of tet-regulated cell lines

The detailed procedures for establishment of HeLa-and Huh7-inducible cells were similar to those previously described (39,45). Briefly, to generate HeLa PLpro-inducible cells, HeLa Tet-Off cells were transfected with pTRE2Bla-SARS-PLpro-TM using Fugene 6 (Roche) and double-selected in complete medium supplemented with 100µg/ml G418, 1 µg/ml Blasticidin and 2 µg/ml tet. Approximately three weeks later, individual cell colonies were selected, expanded, and examined for SARS-CoV PLpro-TM expression by indirect immunofluorescence staining using a mouse monoclonal antibody (mAb) against the V5 epitope (Invitrogen) (1:500) that is present at the C-terminus of PLpro-TM, after being cultured in the absence of tet for 4 days. Two independent cell clones, designated HeLa PLpro- 4 and -10, respectively, allowed tight regulation of PLpro-TM expression under tet control, and were selected for further characterization. Inducible expression of PLpro-TM in these cells was also confirmed by immunoblot analysis using both the V5-tag mAb and a rabbit antiserum against SARS-CoV nsp3 (27). To generate Huh7 cells with conditional expression of DN-IRF-3, cells were cotransfected with pTet-Off (Clontech) and pTRE2Bla-IRF3DN133 at a ratio of 1:10, and double-selected in complete medium supplemented with 400 µg/ml G418, 2 µg/ml Blasticidin and 2 µg/ml tet. Expansion and screening of positive clones were conducted by procedures similar to those used for HeLa PLpro-inducible cells, except that a rabbit polyclonal IRF-3 antiserum (kindly provided by Michael David) was used for immunostaining and immunoblot analysis. Three individual cell clones, designated Huh7 DN-6, -12, and -18, respectively, allowed tightly regulated expression of DN-IRF3 at various levels. However, under fully induced conditions (-tet), only DN-18 cells expressed DN-IRF-3 at a level higher than that of the endogenous IRF-3, and thus allowed a demonstrable DN effect on viral induction of IRF-3 target genes (Fig. 9A and data not shown). Therefore, we selected DN-18 cells for further analysis.

### SeV infection and poly (I-C) treatment

Where indicated, cells were infected with 100 hemagglutinin units (HAU)/ml of SeV (Cantell strain, Charles River Laboratories) for 8–16 h prior to cell lysis for luciferase/β-gal reporter assays and/or immunoprecipitation/immunoblot analysis as previously described (39–41,46). For poly (I-C) treatment, poly (I-C) (Sigma) was added directly to the culture medium at 50µg/ml (M-pIC) or complexed with lipofectamine 2000 (at 1:1 ratio) (T-pIC) and loaded onto the cells for the indicated time period.

### SARS-CoV infection and titration

Where indicated, cells were infected with the Urbani strain of SARS-CoV (kindly provided by T. G. Ksiazek, Centers for Disease Control and Prevention, Atlanta, GA) at the indicated MOI as previously described (47). The cell-free SARS-CoV stock with a titer of  $1 \times 10^7$  TCID<sub>50</sub>/ml (50% tissue culture infectious dose) was generated following 2 passages of the original stock in Vero E6 cells and stored in aliquots at –80 °C. All experiments involving infectious viruses were conducted in an approved biosafety level 3 laboratory at the University of Texas Medical Branch (UTMB). The infectious viral titers in the cell-free supernatants collected at different time points postinfection were determined by a standard TCID<sub>50</sub> assay on permissive Vero E6 cell monolayers in 96-well plates with a series of 10-fold-diluted samples. The titer was expressed as TCID<sub>50</sub> per ml of individual samples.

### IFN bioactivity assay

IFN bioactivity in gamma-irradiated cell culture supernatants was determined by a standard microtiter plaque reduction assay using vesicular stomatitis virus on Vero cells, as described previously (48).

### Transfection and reporter gene assay

Cells grown in 24-well plates ( $5 \times 10^4$ /well) were transfected in triplicate with the indicated reporter plasmid (100 ng), pCMV $\beta$ gal (100 ng), and the indicated amount of an expression vector (50–400 ng and supplied with an empty vector to keep the total amount of DNA transfected constant) using Fugene 6 transfection reagent (Roche), as per the manufacturer's instructions. Twenty-four hours later, transfected cells were mock-treated, treated with poly (I-C) for 6 h, or infected with SeV for 16 h before cell lysis and assay for both firefly luciferase and  $\beta$ -galactosidase activities. Data was expressed as mean relative luciferase activity (luciferase activity divided by  $\beta$ -galactosidase activity) with Standard Deviation from a representative experiment carried out in triplicate. A minimum of three independent experiments were performed to confirm the results of each experiment. The fold change of promoter activity was calculated by dividing the relative luciferase activity of stimulated cells by that of mock-treated cells.

### Indirect immunofluorescence staining

Cells grown in chamber slides (LabTek) following the indicated treatments were washed with PBS and fixed with 4% paraformaldehyde in PBS for 30 min at room temperature. Cells were then permeabilized with 0.2% Triton X-100 in PBS for 15 min, and blocked in 3% BSA at RT for 30 min. After a PBS rinse, the slides were incubated with a rabbit polyclonal antiserum against human IRF3 (1:500, kindly provided by Michael David) (49), or with a V5 tag mAb (1:500, Invitrogen), or with a rabbit polyclonal antiserum (R3) against SARS-CoV nsp3 (27) (1:1000) in 3% BSA for 1hr at RT. Following 3X wash in PBS, slides were incubated with FITC or Texas Red-conjugated, goat anti-rabbit or goat anti-mouse IgG secondary antibodies (Southern biotech) at 37°C for 30 min. Slides were washed, mounted in Vectashield mounting medium containing 4',6-diamidino-2-phenylindole (DAPI, to view the nuclei) (Vector labs), and sealed with a coverslip. Stained slides were examined under a Zeiss Axioplan II fluorescence microscope or an LSM-510 META laser scanning confocal microscope in the UTMB Optimal Imaging Core. For calculation of the percentage of cells demonstrating IRF-3 nuclear translocation, the numbers of cells showing IRF-3 nuclear translocation were counted in four random fields at 20X magnification and/or ten random fields at 40X magnification, and divided by the total numbers of cells (DAPI-positive) in the total counted fields.

### RT-PCR

Following the indicated treatments/viral infections, total cellular RNA was extracted from Calu-3, MA104, or HeLa PLpro-inducible cells using TRIzol reagent (Invitrogen), treated with DNaseI to remove genomic DNA contamination and subjected to clean up using an RNeasy Mini Kit (Qiagen). The abundance of cellular mRNAs specific to human IFN- $\beta$ , IL-6 and A20 genes was determined by quantitative real-time RT-PCR (q-RT-PCR) using commercially available primers and Taqman probes (Applied Biosystems) at the UTMB Sealy Center for Cancer Cell Biology Real-time PCR Core. We confirmed that the human IFN- $\beta$  primers and TaqMan probe worked well for MA104 cells of green monkey kidney origin. In some experiments, semiquantitative RT-PCR was also used to compare the mRNA abundance of IFN- $\beta$ , ISG56 and IL-6, and that of viral RNAs, SeV (cantell strain) and SARS-CoV (Urbani strain) as previously described (40). The following forward and reverse primers were used: IFN- $\beta$  (works for both human and rhesus monkey); IFN-B con107+, tcctgtgtgctctccac, and IFN-B con383-, gtctcattccagccagtct. The primers for ISG56 (works for both human and rhesus

monkey) (40) and IL-6 (human) (50) have been described. GAPDH primers were purchased from Clontech. For semiquantitative detection of viral RNAs, the following primers were used: SARS CoV, described in reference (51); SeV; SeV P890+, aatagggaccgcgtctgtct and SeV P1226-, ttccacgctctcttgatct.

### Phosphatase inhibitor treatment

For *in vivo* experiments, HeLa PLpro-4 cells repressed or induced for PLpro expression or HeLa cells were mock-infected or infected with SeV for 12 h, followed by an additional 4-hour incubation with or without 0.05 µg/ml of okadaic acid (OA, Calbiochem) prior to cell lysis and immunoblot analysis. For *in vitro* experiments, forty micrograms of whole cell lysate of HeLa cells infected with SeV were mock-treated, or incubated with 5 units of calf intestine alkaline phosphatase (CIP, New England Biolabs) in CIP buffer, or incubated with 0.05 µg/ml of OA, or treated with CIP in the presence of OA, respectively, at 30°C for 2 h.

### Protein analyses

Cellular extracts were prepared and subjected to immunoblot analysis and/or immunoprecipitation as previously described (39–41). The following mAbs or polyclonal (pAb) antibodies were used: anti-FLAG M2 mAb and pAb, and anti-actin mAb (Sigma); anti-V5 mAb (Invitrogen); anti-GFP mAb (Roche); anti-IRF-3 mAb and rabbit anti-CBP pAb (Santa Cruz); rabbit anti-IRF-3 pAb (from Michael David); rabbit anti-IRF-3-p396 pAb (from John Hiscott); rabbit anti-RIG-I pAb (from Zhijian Chen); rabbit anti-ISG56 pAb (from Ganes Sen); rabbit anti-MxA and anti-Sendai virus pAbs (from Ilkka Julkunen); rabbit anti-SARS-CoV N pAb (Imgenex); rabbit anti-SARS-CoV nsp3 pAb (27); peroxidase-conjugated secondary goat anti-rabbit, and goat anti-mouse pAbs (Southern Biotech). Protein bands were visualized using ECL plus Western Blotting detection reagents (GE Healthcare), followed by exposure to X-ray film. For detection of IRF-3 dimer, native PAGE of protein samples was performed as previously described (52).

## RESULTS

### SARS-CoV weakly activates an IFN response in cultured cells without inducing the phosphorylation of IRF-3

To investigate whether SARS-CoV triggers an IFN response in cell culture, we determined the cellular responses to infection with the Urbani strain of SARS-CoV in a human lung/bronchial epithelial cell line, Calu-3, which serves as an excellent *in vitro* culture model for studying the pathogenesis of SARS-CoV in the respiratory system (47). As a control, cells were also infected with SeV, a paramyxovirus that is a well-characterized inducer of IFN production via the RIG-I pathway (6,40). In Calu-3 cells, SeV infection rapidly triggered phosphorylation and degradation of IRF-3, which was accompanied by potent induction of ISG56, a well-characterized IRF-3 target gene (53,54) (Fig. 1A, left panel, compare lanes 2 vs. 1). However, infection of Calu-3 cells with SARS-CoV did not induce detectable phosphorylation or degradation of IRF-3. Nonetheless, there was a weak, yet reproducible induction of ISG56 expression, starting 24 h postinfection and peaking at 72 h (Fig. 1A, right panels). The weak induction of an IFN response by SARS-CoV in Calu-3 cells was also confirmed by q-RT-PCR analysis of IFN-β mRNA (data not shown). To ensure that these findings generally reflect SARS-CoV infection and were not a cell-type specific phenomenon, we conducted similar experiments in an African green monkey kidney cell line, MA104. These cells have been reported to be permissive for SARS-CoV infection (55), and we have found that MA104 cells supported SARS-CoV replication with faster kinetics than Calu-3 cells, with cytopathic effect (CPE) seen as early as 28 h postinfection (C.T.K. Tseng and K. Li, unpublished observations). In contrast to another African green monkey kidney cell line, Vero, which is widely used for *in vitro* propagation of SARS-CoV and defective in IFN synthesis (56), MA104 cells are IFN-

competent (57). We found that infection of MA104 cells with SARS-CoV upregulated IFN- $\beta$  mRNA by 12- and 49-fold, respectively, at 20 and 30 h postinfection, although the induction of IFN mRNA was significantly weaker than that (409 fold) triggered by transfection of a synthetic dsRNA analog, poly (I-C) (Fig.1B, right panel). Immunoblot analysis (Fig.1B, left panels) demonstrated that SeV triggered the phosphorylation and degradation of IRF-3 in MA104 cells early after infection (8 h), accompanied by the appearance of serine 396-phosphorylated species of IRF-3 (p396-IRF3). However, due to the proteasomal degradation of IRF-3 (17), the abundance of p396-IRF3 detected in SeV-infected MA104 cells was markedly reduced when compared with that induced by transfected poly (I-C) (Supplemental Fig.1, compare lane 5 vs. 3). The weak band reacted with p396-IRF3 antibody was not IRF-7, which is not basally expressed in MA104 cells and only minimally induced during this short infection period (8 h), and has a higher molecular weight than that of p396-IRF-3 (Supplemental Fig.1). SeV also significantly induced the expression of ISG56, as well as MxA, an ISG whose expression is dependent on functional JAK-STAT signaling downstream of the IFN receptor (58). In contrast, SARS-CoV did not induce the phosphorylation or degradation of IRF-3, nor did it trigger IRF-3 phosphorylation at Ser396 site in MA104 cells during the observation period (18 and 28 h, respectively) (Fig.1B, left panels). Nonetheless, we reproducibly observed that SARS-CoV infection of MA104 cells induced a shift in the two IRF-3 basal forms from the lower to the higher mass form, a phenomenon that is more evident late after SARS-CoV infection (26 h) (Fig. 1B, IRF-3 panel, compare lane 4 vs. 1). This is reminiscent of HSV-1 infected A549 cells (59). Furthermore, there was a weak induction of ISG56 protein expression at 26 h postinfection (Fig. 1B, lane 4). Interestingly, despite these findings, induction of MxA occurred early (at 18 h postinfection) and increased with time (Fig. 1B, MxA panel, lanes 3 and 4) in SARS-CoV infected MA104 cells. Collectively, these data suggest that SARS-CoV is recognized by host cells of lung epithelium and kidney origin and able to elicit a weak IFN response, although this does not involve demonstrable phosphorylation and degradation of IRF-3. We conclude that SARS-CoV is subject to the surveillance of cellular innate immunity; however, as previously reported (21), SARS-CoV has likely evolved strategies to evade or counteract these protective host responses.

### **The PLpro domain of SARS-CoV inhibits activation of IFN- $\beta$ promoter following engagement of TLR3 or RIG-I pathways independent of its protease activity**

In searching for SARS-CoV protein(s) that inhibit IFN responses, the PLpro domain of SARS-CoV nsp3 caught our attention. In addition to its role in directing the processing of viral polyproteins, SARS-CoV PLpro was recently shown to have deubiquitination activity (29, 30). As ubiquitination plays a pivotal role in many cellular processes, including innate immunity signaling (60,61), we conducted experiments to determine whether SARS-CoV PLpro could regulate induction of cellular IFN responses. Engagement of the TLR3 pathway by addition of synthetic dsRNA analog, poly (I-C), to culture medium of HEK293 cells that stably express TLR3 (293-TLR3), induced the IFN- $\beta$  promoter by 16 fold. However, poly (I-C)-induced IFN- $\beta$  promoter activity was reduced by 70% in cells ectopically expressing the catalytic core domain of SARS-CoV PLpro (Sol), or PLpro-TM that encodes PLpro-Sol in conjunction with its downstream hydrophobic transmembrane domains (Fig.2A, left panel). Similar inhibition of the activation of IFN- $\beta$  promoter by PLpro-Sol and PLpro-TM was observed when IFN response was triggered by SeV, which activates the RIG-I pathway (6, 11,40) (Fig.2A, right panel). As positive controls, ectopic expression of known viral or mammalian inhibitors of the IFN response, i.e., the papain-like protease of bovine viral diarrhea virus, Npro (43), or a human deubiquitination enzyme, A20 (62), respectively, also strongly suppressed activation of the IFN- $\beta$  promoter by poly (I-C) or SeV (Fig. 2A, left and right panels). Thus, these data suggest that expression of the catalytic core domain of SARS-CoV PLpro is sufficient for inhibiting activation of the IFN- $\beta$  promoter via both TLR3 and RIG-I pathways.

Next, we determined whether the protease activity of PLpro was required for inhibiting the cellular IFN response, a property that is shared by the NS3/4A serine proteases of two flaviviruses, hepatitis C virus and GB virus B (39,46), and the 3ABC protease precursor of a picornavirus, hepatitis A virus (HAV) (63). SARS-CoV PLpro consists of a triad-active site, i.e., Cys1651-His1812-Asp1826. Mutation at any of the three sites is known to abrogate the protease activity of PLpro [(29) and Supplemental Fig. 2]. We found that when PLpro was expressed in conjunction with its downstream hydrophobic transmembrane domain (PLpro-TM) (27), the wild-type, C1651A mutant and D1826A mutant were all able to inhibit activation of the IFN- $\beta$  promoter by SeV (Fig.2B and C) or poly (I-C) (data not shown). The D1826A mutant PLpro suppressed activation of IFN- $\beta$  promoter as efficiently as the WT protein, whereas the C1651A mutant was less efficient than wild-type, but suppressed in a dose-dependent manner. These data suggest that the protease activity of PLpro is not responsible for its inhibition of activation of IFN responses. To further investigate the role of the catalytic residues, we expressed the wild-type and catalytic mutants of PLpro (PLpro-Sol). We found that ectopic expression of the wild-type (WT) PLpro-inhibited the activation of IFN- $\beta$  promoter by SeV (Fig.2D) or medium poly (I-C) (data not shown) in a dose-dependent fashion. In addition, the PLpro-Sol D1826A mutant inhibited the IFN response as efficiently as wild-type PLpro-sol (Fig. 2E). However, there was no appreciable effect with overexpression of the related, catalytically inactive C1651A PLpro-Sol mutant. We hypothesize that the C1651A substitution in PLpro-Sol may alter the folding/conformation of the protein in addition to its disruption of the protease active site. Consistent with this hypothesis, we noticed a significant reduction in the yield of PLpro-Sol containing the C1651A mutation when purifying the protein from *E. Coli* (29), presumably as a result of altered protein folding which renders PLpro mutants insoluble in *E. Coli*. Taken collectively, these data show that the PLpro domain of SARS-CoV is capable of antagonizing the induction of IFN- $\beta$  promoter via both TLR3 and RIG-I pathways, and this occurs independently of the protease activity of PLpro.

### **SARS-CoV PLpro inhibits induction of IRF-3 dependent genes by acting at a level downstream of the IRF-3 kinases and proximal to IRF-3**

The latent cellular transcription factor IRF-3 plays a central role in transcriptional control of the IFN- $\beta$  promoter (17,64). To determine whether PLpro inhibition of the IFN- $\beta$  promoter is related to its suppressive effect on IRF-3 activation, we conducted luciferase reporter assays utilizing an IRF-3-dependent synthetic promoter, 55C1B (6), following poly (I-C) treatment or SeV challenge. As shown in Fig. 3A, we found that activation of 55C1B promoter by either poly (I-C) or SeV was almost completely ablated by ectopic expression of PLpro-TM. Similar results were obtained when another IRF-3-dependent synthetic promoter, PRDIII-I (44), was used in lieu of 55C1B for probing the signaling (data not shown). The two noncanonical I $\kappa$ B kinases, TBK1 and IKK $\epsilon$ , mediate the C-terminal phosphorylation of IRF-3, a prerequisite prior to its activation. They are linked to the distinct viral PRRs, RIG-I/MDA5 and TLR3, by the adaptor proteins MAVS and TRIF, respectively (65–69). To determine the level of the PLpro blockade of IRF-3 activation, we determined the effect of PLpro on induction of IRF-3-dependent promoters, following ectopic expression of signaling proteins known to participate in RIG-I/MDA5 and TLR3 pathways upstream of IRF-3. We found that PLpro-TM strongly inhibited the activation of IFN- $\beta$  promoter by overexpression of the constitutive active CARD domain of RIG-I (N-RIG) or MDA5 (N-MDA5), or by ectopic expression of MAVS, TRIF, TBK1, or IKK $\epsilon$  (Fig.3B). In contrast, the constitutive active, phosphomimetic IRF-3 mutant, IRF-3-5D, activated the IFN- $\beta$  promoter in cells expressing PLpro-TM to a level that was comparable to that in cells transfected with a control vector (Fig.3B). Similar results were obtained when p55C1B-Luc was used in place of IFN- $\beta$ -Luc to probe the signaling (Fig.3C). We conclude from these experiments that PLpro disrupts TLR3 and RIG-I/MDA5 signaling by acting at a level that is downstream of the IRF-3 kinases and proximal to IRF-3.



To better characterize the mechanisms underlying the SARS-PLpro disruption of cellular IFN responses, we developed tet-regulated HeLa cells that conditionally express PLpro-TM under the control of the Tet-Off promoter. Two clonal cell lines, designated PLpro-4 and PLpro-10, respectively, were selected for further analysis. In these cells, expression of PLpro-TM was tightly regulated by tet, and only occurred when tet was removed from the culture medium (Fig.4A and 4B). The expression level of PLpro-TM could also be conveniently titrated by culturing cells at different concentrations of tet (Fig. 4B, right panels). Under confocal microscopy, PLpro-TM showed a perinuclear, cytoplasmic localization that was similar to that reported for nsp3 in SARS-CoV infected cells (27) (Fig.4A, right panel). Confirming the data obtained from promoter-based reporter assays (Figs.2 and 3), we found that de-repression of PLpro-TM expression significantly inhibited the SeV induction of endogenous ISG56 expression in both cell clones (Fig.4B, left panels), and this inhibitory effect was dose-dependent on the expression level of PLpro-TM (Fig.4B, right panels). This could not be attributed to inhibition of SeV replication by PLpro, as similar levels of SeV proteins were produced, regardless of PLpro-TM induction status (Fig.4B, SeV panel). A similar inhibition of poly (I-C) induced ISG56 expression by PLpro-TM was also observed in these cells, regardless of the delivery route, i.e., poly (I-C) being added to culture medium or introduced into cells by transfection, which engage the TLR3 and MDA5 pathways (10,11), respectively (Fig.5A, ISG56 panel, compare lane 6 vs 2, and data not shown). Real-time RT-PCR analysis showed that PLpro-TM expression ablated the upregulation of IFN- $\beta$  mRNA by poly (I-C) or SeV challenge, while leaving the induction of the NF- $\kappa$ B-dependent A20 mRNA transcripts intact (70) (Fig.4C, left and central panels). Furthermore, PLpro-TM potentiated the induction of IL-6 mRNA by poly (I-C) or SeV (Fig.4C, right panel). Taken collectively, these data suggest that SARS-CoV PLpro specifically targets the IRF-3 arm of IFN responses for inhibition.

### **SARS-CoV PLpro interacts with IRF-3, and inhibits virus-induced IRF-3 phosphorylation, dimerization and nuclear translocation**

To determine how SARS-CoV PLpro inhibits IRF-3 activation, we conducted immunoblot analysis of the activation status of IRF-3 in PLpro-inducible cells, repressed or induced for PLpro-TM expression and mock-treated or treated with poly (I-C) or infected with SeV. We found that de-repression of PLpro-TM significantly downregulated the phosphorylation (IRF-3 panel) and dimerization (native gel panel) of IRF-3 (Fig.5A). PLpro-TM induction also ablated SeV-induced phosphorylation of IRF-3 at Ser396 (Fig.5B, compare lane 7 vs 3), a minimal phospho-acceptor site required for *in vivo* activation of IRF-3 in response to virus and dsRNA (71). This could not be attributed to de-phosphorylation of IRF-3, as treatment of PLpro-expressing cells with a potent PP2A phosphatase inhibitor, okadaic acid (72), failed to restore SeV-induced IRF-3 phosphorylation (Fig.5B, compare lane 8 vs. 7 and 4). As positive controls, treatment of SeV-infected HeLa cell lysates with CIP effectively de-phosphorylated IRF-3 (Fig.5C, compare lane 5 vs. 6). The action of CIP, however, was completely blocked by OA (Fig.5C, compare lane 7 vs. 6). Consistent with the inhibition on IRF-3 phosphorylation and dimerization, SeV induced IRF-3 nuclear translocation (Fig.5D), and its association with the transcriptional co-activator CBP (Fig.5E, compare lane 4 vs 2) was greatly reduced in PLpro-expressing cells.

Interestingly, in co-immunoprecipitation (co-IP) experiments, we reproducibly observed that PLpro-TM interacted with IRF-3 (Fig.6A, right panels). The association of PLpro with IRF-3 was specific, as it was detected only when IRF-3 antiserum, but not a control serum, was used for the co-IP experiments (Fig. 6A, right panels, compare lanes 3 and 4 vs. 7 and 8). Furthermore, Co-IP experiments did not reveal an interaction of PLpro with RIG-I, MAVS, TBK1 or IKK $\epsilon$  (data not shown). To determine whether the inhibition of IRF-3 activation by PLpro is related to its interaction with IRF-3, we conducted co-IP experiments determining the ability of different PLpro mutants to interact with IRF-3. We found that WT and the protease-

deficient mutant forms (C1651A and D1826A) of PLpro-TM all interacted with IRF-3 (Fig. 6B), consistent with their ability to inhibit IRF-3 activation (Fig.2). In the case of PLpro-Sol, while D1826A substitution had no effect on its association with IRF-3, the C1651A Sol mutant lost its ability to interact with IRF-3 (Fig.6C). This is in agreement with the fact that D1826A PLpro Sol acts similarly as the WT protein to inhibit IFN induction (Fig.2E), while C1651A PLpro Sol is no longer able to do so (Fig.2D). We speculate that addition of the downstream TM domains to PLpro Sol may render the conformation/folding of PLpro less liable to the C1651A substitution, thereby allowing C1651A PLpro-TM to retain its ability to interact with IRF-3. Next, we investigated whether PLpro can also interact with IRF-3 5D, a constitutively active mutant that mimics the C-terminal phosphorylated IRF-3. This was an intriguing question, as PLpro did not inhibit the activation of IFN response by IRF-3 5D (Fig.3). We found that both WT and 5D versions of GFP-tagged IRF-3 interacted with PLpro-TM, while the GFP control did not (Fig.6D). We conclude from these experiments that SARS-CoV PLpro physically interacts in a complex with IRF-3, leading to inhibition of IRF-3 phosphorylation and its subsequent dimerization, nuclear translocation, and activation of the expression of IRF-3 target genes. However, we note that PLpro does not seem to inhibit the downstream IFN induction once IRF-3 is phosphorylated and activated.

### **Association of nsp3 with IRF-3 and regulation of IFN responses in cells infected with SARS-CoV**

Next, we investigated whether the IRF-3 pathway is regulated in SARS-CoV-infected cells, and whether nsp3, which embraces the PLpro domain, interacts with IRF-3 in the context of SARS-CoV replication. In MA104 cells, SeV infection induced the degradation of IRF-3 and significantly upregulated the expression of ISG56, MxA and RIG-I proteins early after infection (10 h postinfection) (Fig.7, upper panels, compare lanes 2 vs. 1). Again, as described in Fig.1, infection of SARS-CoV triggered a weak ISG response late after infection (at 28 h) without inducing the phosphorylation and degradation of IRF-3 (Fig.7, upper panels, compare lanes 4 and 3 vs. 1). Importantly, pre-infection of SARS-CoV for 18 h ablated the further upregulation of multiple ISG proteins, ISG56, MxA and RIG-I by SeV superinfection (Fig.7, upper panels, compare lanes 5 vs. 2 and 4). Although SARS-CoV infection did not ablate the phosphorylation of IRF-3 induced by SeV challenge, it caused a marked, reproducible reduction in IRF-3 degradation following SeV superinfection (Fig.7, IRF-3 panel, compare lanes 5 vs. 2, and Fig.8A, IRF-3 panel, compare lanes 5 vs. 3). Similar abrogation of the upregulation of ISG proteins by SARS-CoV was observed when poly (I-C) transfection was used in lieu of SeV to trigger the responses (Fig.8A, compare lanes 6 vs. 1 and 4). Consistent with the immunoblot analysis data, while SeV infection or poly (I-C) transfection induced considerable amounts of IFN that were released to the culture supernatant, as measured by IFN bioactivity assay, there was little or no detectable IFN antiviral activity in SeV-challenged or poly (I-C)-treated MA104 cells that were pre-infected with SARS-CoV (Fig. 8B).

Remarkably, we were able to demonstrate a reproducible association of nsp3 with IRF-3 in SARS-CoV-infected MA104 cells by co-IP, and the interaction seemed to be strengthened with time after infection (Fig.7, lower panels, lanes 3 and 4). SeV superinfection did not have a demonstrable effect on the association between nsp3 and IRF-3 (lane 5). Surprisingly and not currently understood, although SARS-CoV infection ablated the induction of IFN and multiple ISG proteins by SeV superinfection or dsRNA transfection (Fig. 7, 8A, and 8B), this did not correlate with an inhibition on upregulation of IFN- $\beta$  mRNA transcript (Fig.8C). While the role of SARS-CoV nsp3 in regulation of IRF-3-dependent innate immunity remains to be further investigated, these data suggest that SARS-CoV has developed additional strategies to inactivate innate cellular defenses by disrupting the expression of a subset of cellular antiviral proteins from their mRNAs.

## The IRF-3 pathway contributes to control of SARS-CoV replication in cultured cells

While this study was in process, three other SARS-CoV proteins, ORF3b, ORF6, and N, were reported to function as IFN antagonists by inhibiting IRF-3 activation (35). The fact that multiple SARS-CoV proteins have evolved the ability to disrupt IRF-3 function prompted us to hypothesize that IRF-3-dependent innate defenses may contribute significantly to cellular control of SARS-CoV infection. We thus sought to determine whether IRF-3 signaling regulates cellular permissiveness for SARS-CoV replication. To answer this question, we developed Huh7 cells (designated Huh7/DN-18) that conditionally express a dominant negative form of IRF-3 (DN-IRF-3) (17) under the control of the Tet-Off promoter. Huh7 cells were reported to be susceptible for SARS-CoV infection, and we have confirmed this independently in our laboratories (K. Li and C.T.K. Tseng, unpublished observations). As seen in Fig. 9A, expression of DN-IRF-3 in Huh7/DN-18 cells was tightly regulated by tet, and only detectable when tet was withdrawn from the culture medium (IRF-3 panel, compare lanes 3 and 4 vs 1 and 2). Confirming its function, DN-IRF-3 induction significantly diminished SeV-induced expression of ISG56 (ISG56 panel, compare lanes 4 vs. 2). We then manipulated DN-18 cells to repress and induce DN-IRF-3 expression, respectively, and challenged them with a low dose of SARS-CoV (MOI=0.01). The yields of infectious progeny SARS-CoV in culture supernatants were determined dynamically during the three-day follow-up period. In three independently conducted experiments, we reproducibly observed that SARS-CoV replicated at a significantly faster rate in cells induced for DN-IRF-3 expression than in those under continuous repression, releasing up to 2-log higher titers of infectious progeny viruses (Fig. 9B). Consistent with this, CPE was observed in as many as 70% of infected cells induced for DN-IRF-3 expression, while in only 30% of cells without DN-IRF-3 expression at day 3 postinfection (data not shown). Taken together, these data suggest that IRF-3 signaling contributes to the cellular control of SARS-CoV infection.

## DISCUSSION

Data regarding the interactions of SARS-CoV with the innate immune system of host cells have been controversial. While it is conceivable that professional IFN-producing, plasmacytoid DCs may sense ssRNAs derived from SARS-CoV in endosome compartments by TLR7 and initiate an IFN response, as recently demonstrated by Cervantes-Barragan et al. (73), it is currently not known whether parenchymal cells, including those from the lung epithelium, which represent the major target for SARS-CoV infection *in vivo*, can sense components or products of the invading SARS-CoV and initiate an innate defense response. Furthermore, it has been controversial whether SARS-CoV evades host innate immunity by simply avoiding detection by the host or has acquired mechanisms to actively block host antiviral responses (35,37,38).

Using a highly relevant human lung/bronchial epithelial cell line (Calu-3) for SARS-CoV infection (47) and a fetal green monkey kidney-derived fibroblast cell line (MA104) competent for IFN production, we have demonstrated in this study that SARS-CoV is sensed by the innate immune system in parenchymal cells that actively support the replication of SARS-CoV. This conclusion is supported by the fact that infection of SARS-CoV triggers a weak, yet reproducible induction of IFN- $\beta$  mRNA as well as a low level expression of a subset of ISGs in both cell types (Figs. 1, 7 and 8). Among these ISGs, induction of MxA and RIG-I is known to depend on functional JAK-STAT signaling downstream of type I IFN receptors (58,74). Moreover, disruption of IRF-3 function by conditional expression of a dominant negative IRF-3 mutant significantly upregulated SARS-CoV replication in cell culture (Fig. 9), indicating that IRF-3 dependent innate defenses are likely being triggered by SARS-CoV and have a restrictive effect on viral replication. In agreement with a previously published study in HEK293 cells (21), infection of SARS-CoV did not trigger demonstrable phosphorylation

of IRF-3 in both Calu-3 and MA104 cells, despite the latter being readily induced by SeV (Fig. 1 and Fig.7) or poly (I-C) transfection (Fig.8A and data not shown). Nonetheless, we were able to reproducibly observe a shift in the two IRF-3 basal forms from the lower to the higher mass in MA104 cells late postinfection (Fig.1B and Fig.7), which represents a low level of IRF-3 activation pattern similar to that previously reported in HSV-1 infected A549 cells (59). The reasons Calu-3 cells do not show such a shift in the two IRF-3 forms are currently not known. While cell type-specific differences could be involved, this seems less likely as both cell types respond to SeV well by rapidly inducing the phosphorylation and degradation of IRF-3 (Fig. 1). It is reasonable to speculate that this difference could be related to an earlier, more robust replication of SARS-CoV in MA104 cells (Fig.1 and data not shown), as the degree of IRF-3 phosphorylation has been shown to correlate with the level of viral replication (59).

While we have shown that SARS-CoV infection is sensed by and triggers innate immune responses of host cells, the IFN response induced by SARS-CoV shows delayed kinetics and the overall magnitude is much lower than that triggered by SeV or transfected dsRNA analog, poly (I-C) (Fig.1). Replication of SARS-CoV occurs within double membrane vesicles (DMVs) derived from ER membranes (75–78), which may prevent the recognition of viral PAMPs generated during SARS-CoV replication by host PRRs (37,38). While this evasion mechanism may contribute to the absence of IRF-3 phosphorylation early after infection, it does not explain the blockade of induction of IFN antiviral activity and the upregulation of multiple ISG proteins that we have observed in SARS-CoV infected cells that were superinfected with SeV or transfected with poly (I-C) (Fig.8B,7 and 8A). Instead, our data suggest that SARS-CoV has evolved strategies to actively block host innate defenses. Consistent with this notion, multiple SARS-CoV proteins have recently been suggested to antagonize various aspects of the host innate immunity, as those reported with ORF3b, ORF6, N and nsp1 (35,36). We have shown in this study that the PLpro domain of nsp3 also serves as an IFN antagonist. However, it is important to consider the possibility that both “stealth” and “counteraction” mechanisms may be involved in virus-host interactions in SARS-CoV-infected cells and regulated in a coordinated manner at different stages of viral life cycle, in order to circumvent host responses and offer SARS-CoV a maximal survival advantage.

In searching for an IFN antagonist encoded by SARS-CoV, we identified the PLpro domain of nsp3 as a specific, potent inhibitor for IRF-3 activation. As one of two major SARS-CoV proteases, PLpro comprises part of the nsp3 protein and is responsible for processing the polyprotein junctions spanning nsp1 through nsp4. Intriguingly, PLpro was recently shown to have unexpected DUB/deISGylation activities (29,30), which prompted us to explore its effect on innate immune signaling. We found that PLpro disrupts IFN induction by acting downstream of the IRF-3 kinases, TBK1 and IKK $\epsilon$ , and inhibits the induction of IRF-3-dependent genes via both TLR3 and RIG-I/MDA5 pathways (Figs. 2 through 5). Importantly, we demonstrated this upon the transient constitutive expression of PLpro or its stable, conditional expression in a tet-regulated manner. Furthermore, we show the inhibitory effect was dose-dependent on the expression of PLpro (Fig.2B, 2C and 4B). Unlike NS3/4A of HCV and GBV-B or 3ABC of HAV (39,46,63), SARS-CoV PLpro does not rely on its protease activity to disrupt IFN responses (Fig.2). Instead, it involves an interaction between PLpro and IRF-3 (Fig.6) and inhibits the phosphorylation, dimerization and nuclear translocation of IRF-3 (Fig.5), prerequisites for its activation as a transcription factor to turn on the type I IFN response (17). The involvement of PLpro-IRF-3 interaction in PLpro’s IFN antagonist function is supported by the fact that the ability to interact with IRF-3 by various PLpro mutants (Fig.6B and 6C) seems to correlate with their capability of inhibiting IFN response (Fig.2). Specifically, the C1651A PLpro-Sol mutant that loses its inhibitory effect on IFN response (Fig.2D) is also unable to interact with IRF-3 (Fig.6C). However, precisely how this interaction contributes to PLpro’s function is not known. We speculate that the association of PLpro with IRF-3 may affect the recognition of IRF-3 by the IRF-3 kinases and/or otherwise prevent its interactions

with the IRF-3 kinases and/or other cellular interaction partners essential for IRF-3 activation. Moreover, our data suggest that PLpro has to act at a step before the phosphorylation of IRF-3, as PLpro does not inhibit the activation of IFN response by the phosphomimetic IRF-3 5D (Fig.3B and 3C), despite of its capability of interacting with IRF-3 5D (Fig.6D).

At present, we do not know whether the DUB/deISGylation activity contributes to PLpro's function in disrupting IRF-3 activation. Although we have shown that its protease activity is dispensable, thus far we have not been able to identify specific mutation(s) in PLpro that allow us to dissect the protease activity of PLpro from its DUB activity. The D1826A protease-inactive mutant PLpro loses ~99% of the DUB activity *in vitro* (29), yet still acts as good as the WT protein to inhibit the IFN response (Fig.2C and 2E). While this would argue against the involvement of the DUB activity of PLpro in mediating the IRF-3 blockade, it does not exclude the possibility that only 1% of the DUB activity is needed to fully execute the inhibitory function of PLpro on IRF-3 activation. Intriguingly, it has been reported that several cellular DUB/deISGylation enzymes inhibit innate immune signaling independent of their DUB/deISGylation function. For instance, the ubiquitin editing enzyme A20 acts through its C-terminal zinc finger domain of the ubiquitin ligase to inhibit IRF-3-dependent gene expression, as deletion of the N-terminal DUB domain has no significant effect on the inhibitory effect of A20 (62). Similarly, the ubiquitin-specific protease Ubp43 negatively regulates IFN signaling through its interaction with the IFNAR2 subunit and independent of its isopeptidase activity towards ISG15 (79).

Ubiquitination is known to play an important role in NF- $\kappa$ B signaling (60). However, we did not find a specific inhibitory effect of PLpro on the induction of endogenous NF- $\kappa$ B dependent genes, A20 (70) and IL-6 (80), following SeV challenge or pIC treatment (Fig.4C), although we did observe some inhibition of activation of the NF- $\kappa$ B-dependent PRDII promoter in cells with transient, ectopic expression of PLpro (S.G. Devaraj and K. Li, data not shown). Future reverse genetics approaches engineering viable SARS-CoV mutants, which carry substitutions in PLpro that disrupt its DUB/deISGylation activities while retaining the protease activity (assuming that this is possible), would help to address the specific biological roles of the DUB/De-ISGylation activities of PLpro in virus-host interactions and pathogenesis of SARS-CoV infection.

For reasons that remain poorly understood, we did not observe a blockade of SARS-CoV replication on phosphorylation of IRF-3 by SeV or transfected poly (I-C) in MA104 cells, although we demonstrate that SeV-induced degradation of IRF-3 was dramatically reduced (Fig.7 and 8A). We were also unable to demonstrate an inhibition on SeV induction of IFN- $\beta$  mRNA transcript by prior SARS-CoV infection (Fig.8C). This is consistent with two recent reports that MHV and SARS-CoV do not block poly (I-C) or SeV-induced IRF-3 nuclear translocation (37,38), but is contradictory to the fact that multiple SARS-CoV proteins, including PLpro, ORF3b, ORF6, and N (35), can efficiently inhibit IRF-3 activation. The treatments used to trigger IFN response [SeV or poly (I-C) transfection] were administered at 18 h postinfection of SARS-CoV (MOI=10) when SARS-CoV growth kinetics had reached its plateau and about 50% cells were infected as determined by immunofluorescence staining for SARS-CoV nsp3 antigen (data not shown). In contrast, in the study conducted by Versteeg et al (38), poly (I-C) was transfected at 1 h postinfection of SARS-CoV and SeV was coinfecting with SARS-CoV. It is possible that uninfected cells still responded to SeV or poly (IC) which may have diluted the inhibitory effect of PLpro and/or other viral IFN antagonists. However, it would be needed to determine whether PLpro can inhibit IRF-3 activation when expressed in the context of the entire nsp3 protein. However, thus far we have not been able to clone the full-length nsp3 cDNA in a mammalian expression vector. The reasons for this remain unclear, but could be due to a toxic effect of nsp3 in *E. Coli*. One may also speculate that PLpro, as an essential viral protease involved in viral replication, is sequestered within the replication

complex in SARS-CoV-infected cells, preventing its binding to IRF-3. Although less likely, it is also possible that SeV replication or poly (I-C) transfection may disrupt the DMVs, resulting in the release of additional SARS-CoV PAMPs that trigger IFN responses via as yet unknown mechanisms that are not sensitive to PLpro's inhibition. Nevertheless, we were able to reproducibly observe that nsp3 co-precipitates with IRF-3 in SARS-infected cells (Fig. 7, lower panels), although further studies are needed to determine whether such an interaction is direct or through another protein partner(s). Furthermore, the PLP2 of NL63, another closely related coronavirus, also interacts with IRF-3 and has a similar inhibitory effect on IFN responses (Z.B. Chen and S.C. Baker, unpublished data). Therefore, it seems that inhibition of IRF-3 activation by PLpro is a shared feature of, at least some other coronaviruses. Future reverse genetics approaches engineering viable SARS-CoV mutants, which carry substitutions in PLpro that disrupt its interaction with IRF-3 while retaining the protease activity, would help us to fully understand the contribution of PLpro-IRF-3 interaction in regulation of host innate immunity in SARS-CoV infected cells.

Our data demonstrated that SARS-CoV infection ablates the upregulation of protein levels of several ISGs, e.g., ISG56, RIG-I, etc. and the secretion of detectable IFN antiviral activity (Fig. 7, 8A, and 8B). This indicates that SARS-CoV has developed additional strategies to inactivate innate cellular defenses by disrupting the expression of at least a subset of cellular antiviral proteins from their mRNAs. Whether this reflects a selective disruption of certain cellular mRNA translation, nuclear export or turnover, or is due to a nonspecific degradation of mRNAs that was recently reported with nsp1 (36), requires further investigation.

Our data suggest that IRF-3-dependent host defenses contribute to cellular control of SARS-CoV infection, as disruption of IRF-3 function significantly enhanced the replication of SARS-CoV (Fig. 9). This is consistent with the finding that SARS-CoV triggers a weak IFN response in cell culture (Fig. 1), and that multiple SARS-CoV proteins, including PLpro identified in this study, as well as several others, ORF3b, ORF6 and N (35), have the ability to inhibit IRF-3 activation. Although the biological relevance of this newly discovered function for these SARS-CoV proteins remains to be carefully determined in the context of SARS-CoV infection, it suggests that limiting IRF-3-dependent innate immunity is important for establishment of SARS-CoV infection. Further studies are needed, however, regarding the nature of the cellular effectors downstream of IRF-3 that negatively regulate SARS-CoV replication. Specifically, whether this involves cellular genes of direct IRF-3 target or requires a soluble factor to be secreted by infected cells (e.g. IFN). Moreover, it remains to be elucidated regarding the innate signaling pathways that are triggered by SARS-CoV and lead to the IRF-3-dependent innate defenses against SARS-CoV infection.

## Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

## Acknowledgments

This work was supported in part by NIH grants AI069285 and DA018054 (to K.L.), AI45798 and AI060915 (to S.C.B), AI30039 and a WRCE Career Development grant (AI057156) (to C.T.K.T). K.L. is the Cain Foundation Investigator in Innate Immunity.

We thank Stanley Lemon for support and critical review of the manuscript; Samuel Baron for advice with the IFN bioassay; Kate Fitzgerald, Takashi Fujita, John Hiscott, Zhijian Chen, Ilkka Julkunen, Michael David, Ganes Sen, Michael Gale, Nancy Raab-Traub, Christina Ehrhardt, Thomas Ksiazek, and Mohammad Jamaluddin for sharing reagents. We are grateful to Mardelle Susman for careful reading of the manuscript, the UTMB Optimal Imaging Core for assistance with confocal microscopy and the UTMB Sealy Center for Cancer Biology Real Time PCR core for assistance with real-time RT-PCR assays.

## The abbreviations used are

<b>SARS-CoV</b>	severe acute respiratory syndrome coronavirus
<b>PLpro</b>	Papain-like protease
<b>PAMPs</b>	pathogen associated molecular patterns
<b>IFN</b>	interferon
<b>IRF-3</b>	interferon regulatory factor-3
<b>ISG</b>	interferon-stimulated gene
<b>dsRNA</b>	double-stranded RNA
<b>TLR</b>	Toll-like receptor
<b>TRIF</b>	Toll-IL1 receptor domain-containing adaptor inducing IFN- $\beta$
<b>RIG-I</b>	retinoic acid-inducible gene I
<b>MDA5</b>	melanoma differentiation associated gene-5
<b>MAVS</b>	mitochondrial antiviral signaling protein
<b>TBK1</b>	Tank binding kinase 1
<b>DUB</b>	deubiquitination
<b>SeV</b>	Sendai virus
<b>CPE</b>	cytopathic effect
<b>DAPI</b>	4',6-diamidino-2-phenylindole

## References


1. Sen GC. Annu Rev Microbiol 2001;55:255–281. [PubMed: 11544356]
2. Goutagny N, Severa M, Fitzgerald KA. Nat Immunol 2006;7(6):555–557. [PubMed: 16715065]
3. Stetson DB, Medzhitov R. Immunity 2006;25(3):373–381. [PubMed: 16979569]

4. Garcia-Sastre A, Biron CA. *Science* 2006;312:879–882. [PubMed: 16690858]
5. Kawai T, Akira S. *Nat Immunol* 2006;7(2):131–137. [PubMed: 16424890]
6. Yoneyama M, Kikuchi M, Natsukawa T, Shinobu N, Imaizumi T, Miyagishi M, Taira K, Akira S, Fujita T. *Nat Immunol* 2004;5(7):730–737. [PubMed: 15208624]
7. Seth RB, Sun L, Chen ZJ. *Cell Res* 2006;16(2):141–147. [PubMed: 16474426]
8. Meylan E, Tschopp J. *Mol Cell* 2006;22(5):561–569. [PubMed: 16762830]
9. Sen GC, Sarkar SN. *Cytokine Growth Factor Rev* 2005;16(1):1–14. [PubMed: 15733829]
10. Alexopoulou L, Holt AC, Medzhitov R, Flavell RA. *Nature* 2001;413(6857):732–738. [PubMed: 11607032]
11. Kato H, Takeuchi O, Sato S, Yoneyama M, Yamamoto M, Matsui K, Uematsu S, Jung A, Kawai T, Ishii KJ, Yamaguchi O, Otsu K, Tsujimura T, Koh CS, Reis e Sousa C, Matsuura Y, Fujita T, Akira S. *Nature* 2006;441(7089):101–105. [PubMed: 16625202]
12. Hornung V, Ellegast J, Kim S, Brzozka K, Jung A, Kato H, Poeck H, Akira S, Conzelmann KK, Schlee M, Endres S, Hartmann G. *Science* 2006;314(5801):994–997. [PubMed: 17038590]
13. Pichlmair A, Schulz O, Tan CP, Naslund TI, Liljestrom P, Weber F, Reis e Sousa C. *Science* 2006;314:997–1001. [PubMed: 17038589]
14. Sharma S, tenOever BR, Grandvaux N, Zhou GP, Lin R, Hiscott J. *Science* 2003;300(5622):1148–1151. [PubMed: 12702806]
15. Fitzgerald KA, McWhirter SM, Faia KL, Rowe DC, Latz E, Golenbock DT, Coyle AJ, Liao SM, Maniatis T. *Nat Immunol* 2003;4:491–496. [PubMed: 12692549]
16. Yoneyama M, Suhara W, Fujita T. *J Interferon Cytokine Res* 2002;22(1):73–76. [PubMed: 11846977]
17. Lin R, Heylbroeck C, Pitha PM, Hiscott J. *Mol Cell Biol* 1998;18:2986–2996. [PubMed: 9566918]
18. Diebold SS, Kaisho T, Hemmi H, Akira S, Reis e Sousa C. *Science* 2004;303(5663):1529–1531. [PubMed: 14976261]
19. Heil F, Hemmi H, Hochrein H, Ampenberger F, Kirschning C, Akira S, Lipford G, Wagner H, Bauer S. *Science* 2004;303(5663):1526–1529. [PubMed: 14976262]
20. Peiris JS, Guan Y, Yuen KY. *Nat Med* 2004;10(12 Suppl):S88–97. [PubMed: 15577937]
21. Spiegel M, Pichlmair A, Martinez-Sobrido L, Cros J, Garcia-Sastre A, Haller O, Weber F. *J Virol* 2005;79:2079–2086. [PubMed: 15681410]
22. Yount B, Roberts RS, Sims AC, Deming D, Frieman MB, Sparks J, Denison MR, Davis N, Baric RS. *J Virol* 2005;79:14909–14922. [PubMed: 16282490]
23. Ziebuhr J. *Adv Exp Med Biol* 2006;581:3–11. [PubMed: 17037497]
24. Ziebuhr J. *Curr Top Microbiol Immunol* 2005;287:57–94. [PubMed: 15609509]
25. Marra MA, Jones SJ, Astell CR, Holt RA, Brooks-Wilson A, Butterfield YS, Khattri J, Asano JK, Barber SA, Chan SY, Cloutier A, Coughlin SM, Freeman D, Girn N, Griffith OL, Leach SR, Mayo M, McDonald H, Montgomery SB, Pandoh PK, Petrescu AS, Robertson AG, Schein JE, Siddiqui A, Smailus DE, Stott JM, Yang GS, Plummer F, Andonov A, Artsob H, Bastien N, Bernard K, Booth TF, Bowness D, Drebot M, Fernando L, Flick R, Garbutt M, Gray M, Grolla A, Jones S, Feldmann H, Meyers A, Kabani A, Li Y, Normand S, Stroher U, Tipples GA, Tyler S, Vogrig R, Ward D, Watson B, Brunham RC, Krajden M, Petric M, Skowronski DM, Upton C, Roper RL. *Science* 2003;300:1399–1404. [PubMed: 12730501]
26. Rota PA, Oberste MS, Monroe SS, Nix WA, Campagnoli R, Icenogle JP, Penaranda S, Bankamp B, Maher K, Chen MH, Tong S, Tamin A, Lowe L, Frace M, DeRisi JL, Chen Q, Wang D, Erdman DD, Peret TC, Burns C, Ksiazek TG, Rollin PE, Sanchez A, Liffick S, Holloway B, Limor J, McCaustland K, Olsen-Rasmussen M, Fouchier R, Gunther S, Osterhaus AD, Drosten C, Pallansch MA, Anderson LJ, Bellini WJ. *Science* 2003;300(5624):1394–1399. [PubMed: 12730500]
27. Harcourt BH, Jukneliene D, Kanjanahaluethai A, Bechill J, Severson KM, Smith CM, Rota PA, Baker SC. *J Virol* 2004;78(24):13600–13612. [PubMed: 15564471]
28. Ratia K, Saikatendu KS, Santarsiero BD, Barretto N, Baker SC, Stevens RC, Mesecar AD. *Proc Natl Acad Sci U S A* 2006;103(15):5717–5722. [PubMed: 16581910]
29. Barretto N, Jukneliene D, Ratia K, Chen Z, Mesecar AD, Baker SC. *J Virol* 2005;79 (24):15189–15198. [PubMed: 16306590]



30. Lindner HA, Fotouhi-Ardakani N, Lytvyn V, Lachance P, Sulea T, Menard R. *J Virol* 2005;79(24):15199–15208. [PubMed: 16306591]
31. Zheng L, Baumann U, Reymond JL. *Nucleic Acids Res* 2004;32(14):e115. [PubMed: 15304544]
32. Solorzano A, Webby RJ, Lager KM, Janke BH, Garcia-Sastre A, Richt JA. *J Virol* 2005;79(12):7535–7543. [PubMed: 15919908]
33. Quinlivan M, Zamarin D, Garcia-Sastre A, Cullinane A, Chambers T, Palese P. *J Virol* 2005;79(13):8431–8439. [PubMed: 15956587]
34. Li Z, Jiang Y, Jiao P, Wang A, Zhao F, Tian G, Wang X, Yu K, Bu Z, Chen H. *J Virol* 2006;80(22):11115–11123. [PubMed: 16971424]
35. Kopecky-Bromberg SA, Martinez-Sobrido L, Frieman M, Baric RA, Palese P. *J Virol* 2007;81(2):548–557. [PubMed: 17108024]
36. Kamitani W, Narayanan K, Huang C, Lokugamage K, Ikegami T, Ito N, Kubo H, Makino S. *Proc Natl Acad Sci U S A* 2006;103(34):12885–12890. [PubMed: 16912115]
37. Zhou H, Perlman S. *J Virol* 2007;81(2):568–574. [PubMed: 17079305]
38. Versteeg GA, Bredenbeek PJ, van den Worm SH, Spaan WJ. *Virology* 2007;361(1):18–26. [PubMed: 17316733]
39. Chen Z, Benureau Y, Rijnbrand R, Yi J, Wang T, Warter L, Lanford RE, Weinman SA, Lemon SM, Martin A, Li K. *J Virol* 2007;81:964–976. [PubMed: 17093192]
40. Li K, Chen Z, Kato N, Gale M Jr, Lemon SM. *J Biol Chem* 2005;280(17):16739–16747. [PubMed: 15737993]
41. Li K, Foy E, Ferreon JC, Nakamura M, Ferreon AC, Ikeda M, Ray SC, Gale M Jr, Lemon SM. *Proc Natl Acad Sci U S A* 2005;102(8):2992–2997. [PubMed: 15710891]
42. Fries KL, Miller WE, Raab-Traub N. *Virology* 1999;264(1):159–166. [PubMed: 10544141]
43. Chen Z, Rijnbrand R, Jangra RK, Devaraj SG, Qu L, Ma Y, Lemon SM, Li K. *Virology* 2007;366:277–292. [PubMed: 17531282]
44. Ehrhardt C, Kardinal C, Wurzer WJ, Wolff T, von Eichel-Streiber C, Pleschka S, Planz O, Ludwig S. *FEBS Lett* 2004;567(2–3):230–238. [PubMed: 15178328]
45. Li K, Prow T, Lemon SM, Beard MR. *Hepatology* 2002;35(5):1237–1246. [PubMed: 11981774]
46. Foy E, Li K, Wang C, Sumpter R Jr, Ikeda M, Lemon SM, Gale M Jr. *Science* 2003;300(5622):1145–1148. [PubMed: 12702807]
47. Tseng CT, Tseng J, Perrone L, Worthy M, Popov V, Peters CJ. *J Virol* 2005;79 (15):9470–9479. [PubMed: 16014910]
48. Langford MP, Weigent DA, Stanton GJ, Baron S. *Methods Enzymol* 1981;78(Pt A):339–346. [PubMed: 6173610]
49. Navarro L, David M. *J Biol Chem* 1999;274(50):35535–35538. [PubMed: 10585427]
50. Eickelberg O, Pansky A, Mussmann R, Bihl M, Tamm M, Hildebrand P, Perruchoud AP, Roth M. *J Biol Chem* 1999;274(18):12933–12938. [PubMed: 10212284]
51. Ksiazek TG, Erdman D, Goldsmith CS, Zaki SR, Peret T, Emery S, Tong S, Urbani C, Comer JA, Lim W, Rollin PE, Dowell SF, Ling AE, Humphrey CD, Shieh WJ, Guarner J, Paddock CD, Rota P, Fields B, DeRisi J, Yang JY, Cox N, Hughes JM, LeDuc JW, Bellini WJ, Anderson LJ. *N Engl J Med* 2003;348(20):1953–1966. [PubMed: 12690092]
52. Iwamura T, Yoneyama M, Yamaguchi K, Suhara W, Mori W, Shiota K, Okabe Y, Namiki H, Fujita T. *Genes Cells* 2001;6(4):375–388. [PubMed: 11318879]
53. Grandvaux N, Servant MJ, tenOever B, Sen GC, Balachandran S, Barber GN, Lin R, Hiscott J. *J Virol* 2002;76(11):5532–5539. [PubMed: 11991981]
54. Peters KL, Smith HL, Stark GR, Sen GC. *Proc Natl Acad Sci U S A* 2002;99(9):6322–6327. [PubMed: 11972054]
55. Mossel EC, Huang C, Narayanan K, Makino S, Tesh RB, Peters CJ. *J Virol* 2005;79 (6):3846–3850. [PubMed: 15731278]
56. Emeny JM, Morgan MJ. *J Gen Virol* 1979;43(1):247–252. [PubMed: 113494]
57. McKimm-Breschkin JL, Holmes IH. *Infect Immun* 1982;36(3):857–863. [PubMed: 6178689]


58. Ronni T, Matikainen S, Lehtonen A, Palvimo J, Dellis J, Van Eylen F, Goetschy JF, Horisberger M, Content J, Julkunen I. *J Interferon Cytokine Res* 1998;18(9):773–781. [PubMed: 9781817]
59. Collins SE, Noyce RS, Mossman KL. *J Virol* 2004;78(4):1706–1717. [PubMed: 14747536]
60. Chen ZJ. *Nat Cell Biol* 2005;7(8):758–765. [PubMed: 16056267]
61. Liu YC, Penninger J, Karin M. *Nat Rev Immunol* 2005;5:941–952. [PubMed: 16322747]
62. Lin R, Yang L, Nakhaei P, Sun Q, Sharif-Askari E, Julkunen I, Hiscott J. *J Biol Chem* 2006;281(4):2095–2103. [PubMed: 16306043]
63. Yang Y, Liang Y, Qu L, Chen Z, Yi M, Li K, Lemon SM. *Proc Natl Acad Sci U S A* 2007;104(17):7253–7258. [PubMed: 17438296]
64. Yoneyama M, Suhara W, Fukuhara Y, Fukuda M, Nishida E, Fujita T. *Embo J* 1998;17 (4):1087–1095. [PubMed: 9463386]
65. Kawai T, Takahashi K, Sato S, Coban C, Kumar H, Kato H, Ishii KJ, Takeuchi O, Akira S. *Nat Immunol* 2005;6:981–988. [PubMed: 16127453]
66. Meylan E, Curran J, Hofmann K, Moradpour D, Binder M, Bartenschlager R, Tschopp J. *Nature* 2005;437(7062):1167–1172. [PubMed: 16177806]
67. Seth RB, Sun L, Ea CK, Chen ZJ. *Cell* 2005;122(5):669–682. [PubMed: 16125763]
68. Xu LG, Wang YY, Han KJ, Li LY, Zhai Z, Shu HB. *Mol Cell* 2005;19:727–740. [PubMed: 16153868]
69. Yamamoto M, Sato S, Mori K, Hoshino K, Takeuchi O, Takeda K, Akira S. *J Immunol* 2002;169 (12):6668–6672. [PubMed: 12471095]
70. Krikos A, Laherty CD, Dixit VM. *J Biol Chem* 1992;267(25):17971–17976. [PubMed: 1381359]
71. Servant MJ, Grandvaux N, tenOever BR, Duguay D, Lin R, Hiscott J. *J Biol Chem* 2003;278(11):9441–9447. [PubMed: 12524442]
72. Weaver BK, Kumar KP, Reich NC. *Mol Cell Biol* 1998;18:1359–1368. [PubMed: 9488451]
73. Cervantes-Barragan L, Zust R, Weber F, Spiegel M, Lang KS, Akira S, Thiel V, Ludewig B. *Blood* 2007;109(3):1131–1137. [PubMed: 16985170]
74. Akazawa T, Ebihara T, Okuno M, Okuda Y, Shingai M, Tsujimura K, Takahashi T, Ikawa M, Okabe M, Inoue N, Okamoto-Tanaka M, Ishizaki H, Miyoshi J, Matsumoto M, Seya T. *Proc Natl Acad Sci U S A* 2007;104(1):252–257. [PubMed: 17190817]
75. Stertz S, Reichelt M, Spiegel M, Kuri T, Martinez-Sobrido L, Garcia-Sastre A, Weber F, Kochs G. *Virology* 2007;361(2):304–315. [PubMed: 17210170]
76. Gosert R, Kanjanahaluethai A, Egger D, Bienz K, Baker SC. *J Virol* 2002;76(8):3697–3708. [PubMed: 11907209]
77. Goldsmith CS, Tatti KM, Ksiazek TG, Rollin PE, Comer JA, Lee WW, Rota PA, Bankamp B, Bellini WJ, Zaki SR. *Emerg Infect Dis* 2004;10(2):320–326. [PubMed: 15030705]
78. Snijder EJ, van der Meer Y, Zevenhoven-Dobbe J, Onderwater JJ, van der Meulen J, Koerten HK, Mommaas AM. *J Virol* 2006;80(12):5927–5940. [PubMed: 16731931]
79. Malakhov MP, Malakhova OA, Kim KI, Ritchie KJ, Zhang DE. *J Biol Chem* 2002;277(12):9976–9981. [PubMed: 11788588]
80. Richmond A. *Nat Rev Immunol* 2002;2(9):664–674. [PubMed: 12209135]



**Figure 1. SARS-CoV weakly activates an IFN response without inducing the hyperphosphorylation of IRF-3**

A. Human lung/bronchial epithelial Calu-3 cells were mock-infected, or infected with 100 HAU/ml SeV (left panels) or SARS-CoV (right panels) at an MOI of 5 for the indicated times prior to cell lysis and immunoblot analysis of IRF-3, ISG56, and SARS-CoV nucleocapsid (NC) protein. B. The left panels show African green monkey kidney MA104 cells that were mock-infected or infected with SeV (100 HAU/ml) or with SARS-CoV (MOI=10) for the indicated times prior to cell lysis and immunoblot analysis of phospho-Ser396 IRF-3, total IRF-3, MxA, ISG56, SARS-CoV NC, and SeV proteins. A nonspecific band (\*) detected by the anti-ISG56 antibody indicates equal loading. The arrowhead and hatch marks in the total


IRF-3 panel denote phosphorylated (virus-activated) and inactive IRF-3 forms, respectively. In the right panel, MA104 cells grown in 35-mm dishes were mock-treated, infected with SARS-CoV (MOI=10) for 20 or 30 h, or transfected with 5  $\mu$ g poly (I-C) for 10 h prior to total RNA extraction and real-time RT-PCR analysis of IFN- $\beta$  mRNA abundance, which was subsequently normalized to cellular 18S ribosomal RNA. Bars show fold change in IFN- $\beta$  mRNA levels (relative to mock-treated cells).



**Figure 2. SARS-CoV PLpro domain inhibits activation of IFN- $\beta$  promoter following engagement of TLR3 or RIG-I pathways independent of its protease activity**


A. HEK293-TLR3 cells were cotransfected with pIFN- $\beta$ -Luc and pCMV $\beta$ gal plasmids, and a vector encoding SARS PLpro-Sol or SARS PLpro-TM, BVDV Npro, or A20, or an empty vector. Twenty-four hours later, cells were either mock-treated (empty bars), or incubated with 50  $\mu$ g/ml poly (I-C) in culture medium for 6h (hatched bars, left panel), or infected with SeV at 100 HAU/ml for 16 h (solid bars, right panel) prior to cell lysis for both luciferase and  $\beta$ -galactosidase assays. Bars show relative luciferase activity normalized to  $\beta$ -galactosidase activity, i.e, IFN- $\beta$  promoter activity. B and C. IFN- $\beta$  promoter activity in cells transfected with increasing amounts of WT or C1651A mutant PLpro-TM-expressing plasmid (B) or

equivalent amounts of WT or D1826A mutant PLpro-TM plasmid (C). D and E. IFN- $\beta$  promoter activity in cells transfected with increasing amounts of WT or C1651A mutant PLpro-Sol-expressing plasmid (D) or equivalent amounts of WT, C1651A or D1826A mutant PLpro Sol plasmids (E).



**Figure 3. SARS-CoV PLpro inhibits activation of IRF-3-dependent promoters by acting at a level downstream of the IRF-3 kinases and proximal to IRF-3**

A. Activation of p55C1B, an IRF-3-dependent synthetic promoter, by medium poly (I-C) (hatched bars) or SeV infection (solid bars) in HEK293-TLR3 cells with expression of SARS-CoV PLpro-TM (TM) or a control vector (Vect). B and C. Activation of IFN- $\beta$  (B) and p55C1B (C) promoters by ectopic expression of various signaling molecules within TLR3 and RIG-I/MDA5 pathways at or above the level of IRF-3 in HEK293 cells with (solid bars) or without (empty bars) expression of SARS-CoV PLpro-TM. N-RIG and N-MDA5 denote the CARD domain of RIG-I and MDA5, respectively.




**Figure 4. Conditional expression of SARS-CoV PLpro-TM disrupts endogenous, IRF-3-dependent gene expression following dsRNA stimulation or SeV infection**

A. Immunofluorescence staining of PLpro-TM in HeLa PLpro-4 cells cultured in the presence (left) or absence (right) of 2 μg/ml tetracycline for 3 days followed by confocal microscopy. PLpro-TM showed green fluorescence staining (detected with a V5 tag antibody), while nuclei were counterstained blue with DAPI. B. The left panels show HeLa PLpro-4 and PLpro-10 cells that were cultured with and without tetracycline for 3 days and subsequently mock-infected or challenged with SeV for 16 h prior to cell lysis and immunoblot analysis of PLpro (using a V5 tag antibody), ISG56, actin and SeV. The right panels show HeLa PLpro-4 cells that were cultured in the indicated concentrations of tetracycline for 3 days prior to mock




infection (lanes 9 through 13) or infection of SeV (lanes 14 through 17) for 16 h. Expression of PLpro-TM, ISG56, and actin were determined by immunoblot analysis. C. Real-time RT-PCR analysis of IFN- $\beta$  (left), A20 (middle), and IL-6 (right) mRNA transcripts in HeLa PLpro-4 cells repressed or induced for PLpro-TM expression, and mock treated (empty bars), treated with 50  $\mu$ g/ml poly(I-C) in culture medium (hatched bars), or infected with SeV (solid bars) for 16 h. mRNA abundance was normalized to cellular 18S ribosomal RNA. Fold changes were calculated by dividing the normalized mRNA abundance under various treatment conditions by that of the mock-treated cells without PLpro expression.



**Figure 5. SARS-CoV PLpro inhibits virus-induced IRF-3 phosphorylation, dimerization and nuclear translocation**

A. HeLa PLpro-4 cells were manipulated to repress or induce PLpro-TM expression for 3 days, and were subsequently mock-treated, or incubated with 50  $\mu\text{g/ml}$  poly (I-C) in culture medium for 6 h, or challenged with SeV for 16 h. Cell lysates were separated on native PAGE followed by immunoblot analysis to detect IRF-3 monomer and dimer forms (top panel, empty arrow indicates IRF-3 dimer), or subjected to SDS-PAGE and immunoblot analysis of IRF-3, ISG56, PLpro, SeV and Actin (lower panels). B. HeLa PLpro-4 cells were cultured to repress or induce PLpro-TM expression and subsequently mock-infected or infected with SeV for 16 h. Where indicated, cells were incubated with 0.05  $\mu\text{g/ml}$  of okadaic acid (OA) for the last 4 h of SeV


infection/mock infection. Cells were then harvested for immunoblot analysis of IRF-3, p396-IRF3, PLpro (anti-V5), and SeV. \* denotes a nonspecific band. C. Lanes 1 through 4, HeLa cells were mock-infected or infected with SeV for 16h. Where indicated, OA was included in the last 4-h duration of infection/mock infection. Lanes 5 through 8, Cell lysates of HeLa cells infected with SeV were mock-treated, treated with CIP, or OA, or CIP in the presence of OA, as indicated in the Experimental Procedures. IRF-3 and p396-IRF3 were detected by immunoblot analysis. \* denotes a nonspecific band. D. Immunofluorescence staining of IRF-3 subcellular localization in HeLa PLpro-4 cells repressed (upper panels) or induced (lower panels) for PLpro-TM expression, and mock-infected (left) or infected (right) with SeV for 16h. Numbers in SeV-infected cells were the averages of the percentage of cells that had IRF-3 nuclear translocation and were counted from ten random fields (40 X magnification). E. HeLa PLpro-4 cells were cultured to repress or induce PLpro-TM expression and subsequently mock-infected or infected with SeV. Cell lysates were subjected to immunoprecipitation with a rabbit anti-CBP antibody (upper panels) or with a rabbit anti-IRF-3 antiserum (lower panels). The immunoprecipitates were separated on SDS-PAGE, followed by immunoblot analysis of IRF-3 (using an mAb anti-IRF-3) or CBP (using a rabbit anti-CBP antibody).




**Figure 6. SARS-CoV PLpro interacts with IRF-3**

A. HeLa PLpro-4 cells were cultured to repress or induce PLpro-TM expression and subsequently mock-infected or infected with SeV for 16 h. In left panels, expression of p396-IRF3, total IRF-3, PLpro and actin proteins in cell lysates were determined by immunoblot analysis. In right panels, cell lysates were subjected to immunoprecipitation with a rabbit anti-IRF3 antibody (lanes 1 through 4) or a control rabbit serum (lanes 5 through 8). The immunoprecipitates were separated on SDS-PAGE, followed by immunoblot analysis of IRF-3 (using an mAb anti-IRF-3) and PLpro (using an mAb anti-V5 tag). B. HEK293 cells were transfected with WT (lanes 1 through 3), C1651A (lanes 4 through 6), or D1826A (lanes 7 through 9) PLpro-TM, respectively, and where indicated, along with an empty vector (lanes 1, 4, and 7), or a vector expressing Flag-IRF-3 (lanes 2, 5, and 8) or untagged IRF-3 (lanes 3, 6, and 9). Cell lysates were immunoprecipitated with anti-V5 (PLpro-TM), followed by

immunodetection of Flag-IRF-3 (anti-Flag) or PLpro-TM (anti-V5) (right panels). \* denotes IgG heavy chain. Immunoblot analysis of input for Flag-IRF-3 and PLpro-TM is shown in left panels. All 3 forms of PLpro-TM interacted with IRF-3. C. HEK293 cells were transfected with WT (lanes 1 and 2), C1651A (lanes 3 and 4), or D1826A (lanes 5 and 6) PLpro-Sol, respectively, and where indicated, along with an empty vector (lanes 1, 3, and 5) or a vector expressing Flag-IRF-3 (lanes 2, 4, and 6). Cell lysates were immunoprecipitated with anti-V5 (PLpro-Sol), followed by immunodetection of Flag-IRF-3 (anti-Flag) or PLpro-Sol (anti-V5) (lower panels). Immunoblot analysis of input for Flag-IRF-3 and PLpro-Sol is shown in upper panels. Note that WT and D1826A PLpro-Sol interacted with IRF-3, while C1651A PLpro-Sol did not. D. HeLa PLpro-4 cells induced for PLpro-TM expression were transfected with GFP, GFP-IRF-3, or GFP-IRF3-5D, respectively. Cell lysates were subjected to Co-IP experiments using either GFP antibody or V5 antibody for immunoprecipitation, followed by immunoblot analysis of the immunoprecipitates using anti-GFP or anti-V5 antibodies. Note that both GFP-IRF3 and GFP-IRF3-5D were associated with PLpro-TM (detected by anti-V5), while free GFP was not.



**Figure 7. SARS-CoV nsp3 is associated with IRF-3 in cells infected with SARS-CoV**  
 MA104 cells were mock-infected (lane 1), infected with SeV for 10 h (lane 2), infected with SARS-CoV (MOI=10) for 18 and 28 h (lane 3 and 4), respectively, or infected with SARS-CoV for 18 h then superinfected with SeV for an additional 10 h (lane 5). Expression of IRF-3, RIG-I, MxA, ISG56, SARS-CoV NC and nsp3, and SeV proteins in cell lysates was determined by immunoblot analysis (upper panels). A nonspecific band detected by the rabbit anti-MxA antiserum (bottom panel) indicates equal loading. Data shown are representative of three independently conducted experiments. In lower panels, the cell lysates were subjected to immunoprecipitation with a rabbit anti-IRF3 antiserum, followed by immunoblot analysis of SARS-CoV nsp3, and IRF-3 (using an mAb anti-IRF3 antibody). Data shown are representative of two independently conducted experiments.




**Figure 8. Regulation of IFN responses in cells infected with SARS-CoV**

A. MA104 cells were mock-infected (lanes 1 through 3) or infected with SARS-CoV (MOI=10) for 18 h (lanes 4 through 6), followed by mock-treatment (lanes 2 and 4), transfection of poly (I-C) (lanes 1 and 6) or superinfection of SeV (lanes 3 and 5) for 10 h, respectively (a total of 28 h after mock or SARS-CoV infection under these conditions). Immunoblot analysis of cell lysates for expression of ISG56, IRF-3, SeV and actin is shown. Asterisks (\*) in ISG56 panels denote a nonspecific band. B. MA104 cells were mock-infected, infected with SeV for 10 h, infected with SARS-CoV (MOI=10) for 18 or 28 h, or transfected with poly (I-C) for 10 h, respectively, or pre-infected with SARS-CoV for 18 h and then superinfected with SeV or transfected with poly (I-C) for additional 10 h (a total of 28 h after SARS-CoV infection under

these conditions). Cell-free culture supernatants were collected, irradiated, and subjected to IFN bioactivity assay against VSV as described in Experimental Procedures. Bars indicate IFN bioactivity (U/ml) in culture supernatants under the indicated conditions. C. MA104 cells were mock-infected, infected with SeV for 8 h, infected with SARS-CoV (MOI=10) for 16 or 24 h, or pre-infected with SARS-CoV for 16 h and then superinfected with SeV for additional 8 h. Cells were harvested for total RNA extraction, cDNA synthesis, and subsequent real-time RT-PCR detection of IFN- $\beta$  mRNA. Data shown are representative of two independently conducted experiments.





**Figure 9. The IRF-3 pathway regulates SARS-CoV replication in cell culture**

A. Huh7 DN-18 cells with tet-regulated, conditional expression of a dominant negative form of IRF-3 (DN IRF-3) were cultured in the presence or absence of tet for 3 days, to repress or induce DN IRF-3 expression, respectively, and were subsequently mock-infected, or challenged with SeV for 12 h. Expression of IRF-3, ISG56, SeV and actin proteins were detected in cell lysates by immunoblot analysis. The endogenous IRF-3 (FL) and the DN IRF-3 (DN) bands were marked as filled and empty circles, respectively. The asterisk (\*) in ISG56 panel denotes a nonspecific band. B. Huh7 DN-18 cells repressed or induced for DN IRF-3 expression were infected with SARS-CoV at an MOI of 0.01 at 37 °C for 1h. The virus inoculum was then removed and cells were washed extensively and refed with complete medium. Cell-free culture supernatants were collected at the indicated times postinfection and stored at -80 °C until they were subjected to assessment of infectious viral titers using a standard TCID<sub>50</sub> assay in Vero E6 cells, as described in Experimental Procedures. The experiments were terminated at day 3 postinfection as CPE was observed in ~30% and ~70% of infected cells without and with DN IRF-3 expression, respectively. The growth curve of SARS-CoV shown was representative of three independently conducted experiments.