



Published in final edited form as:

*J Neurosci.* 2009 September 23; 29(38): 11982–11992. doi:10.1523/JNEUROSCI.3158-09.2009.

## CD14 and Toll-like receptors 2 and 4 are required for fibrillar A $\beta$ -stimulated microglial activation

Erin G. Reed-Geaghan<sup>1</sup>, Julie C. Savage<sup>1</sup>, Amy G. Hise<sup>2</sup>, and Gary E. Landreth<sup>1</sup>

<sup>1</sup>Alzheimer Research Laboratory, Department of Neurosciences, Case Western Reserve University, School of Medicine, Cleveland, OH 44106

<sup>2</sup>Center for Global Health and Diseases, Case Western Reserve University, Cleveland, OH 44106

### Abstract

Microglia are the brain's tissue macrophages and are found in an activated state surrounding  $\beta$ -amyloid plaques in the Alzheimer's disease brain. Microglia interact with fibrillar  $\beta$ -amyloid (fA $\beta$ ) through an ensemble of surface receptors composed of the  $\alpha_6\beta_1$  integrin, CD36, CD47, and the class A scavenger receptor. These receptors act in concert to initiate intracellular signaling cascades and phenotypic activation of these cells. However, it is unclear how engagement of this receptor complex is linked to the induction of an activated microglial phenotype. We report that the response of microglial cells to fibrillar forms of A $\beta$  requires the participation of Toll like receptors (TLRs) and the co-receptor CD14. The response of microglia to fA $\beta$  is reliant upon CD14, which act together with TLR4 and TLR2 to bind fA $\beta$  and to activate intracellular signaling. We find that cells lacking these receptors could not initiate a Src-Vav-Rac signaling cascade leading to reactive oxygen species production and phagocytosis. The fA $\beta$ -mediated activation of p38 MAPK also required CD14, TLR4, and TLR2. Inhibition of p38 abrogated fA $\beta$ -induced reactive oxygen species production and attenuated the induction of phagocytosis. Microglia lacking CD14, TLR4, and TLR2 showed no induction of phosphorylated I $\kappa$ B $\alpha$  following fA $\beta$ . These data indicate these innate immune receptors function as members of the microglial fA $\beta$  receptor complex and identify the signaling mechanisms whereby they contribute to microglial activation.

### Keywords

microglia; fibrillar A $\beta$ ; Alzheimer's disease; CD14; TLR4; TLR2

### Introduction

Alzheimer's disease (AD) is the most common cause of dementia in the elderly, and is characterized by extensive extracellular deposits of insoluble  $\beta$ -amyloid (A $\beta$ ) in the brain. The most widely held hypothesis for AD pathogenesis posits that a chronic imbalance in the production and clearance of A $\beta$  results in a persistent increase in its steady state levels, ultimately leading to the complex molecular and cellular changes within the brain that typify AD (Selkoe, 2000). A $\beta$  deposition is accompanied by a robust microglial-mediated inflammatory response within the brain (Akiyama et al., 2000). Microglia surrounding A $\beta$  plaques show an activated phenotype and extend processes that envelop the plaque (Itagaki et al., 1989; Perlmutter et al., 1990; Bornemann et al., 2001).

Corresponding Author: Gary E. Landreth, Case Western Reserve University, School of Medicine, 10900 Euclid Avenue, Cleveland, OH 44106-4928, gel2@case.edu.

Section and Senior Editor: Neurobiology of Disease, Dr. Serge Przedborski

Microglia are the brain's tissue macrophage, and continuously survey their immediate environment (Davalos et al., 2005; Nimmerjahn et al., 2005). Following insult, microglia shift from a surveillance to a reactive mode, engaging their immune effector roles to respond to the injury or pathogens. Myeloid lineage cells use ensembles of cell surface receptors to bind complex entities, including fibrillar proteins (Ishibashi et al., 1994; Bornstein, 1995; Wong et al., 1996). Bamberger et al. identified a complex of microglial membrane receptors for fibrillar A $\beta$  (fA $\beta$ ) consisting of the  $\alpha_6\beta_1$  integrin, CD36, CD47, and the class A scavenger receptor (SRA) (Bamberger et al., 2003). Engagement of this receptor complex activates tyrosine kinase-based signaling cascades (McDonald et al., 1997; Combs et al., 1999; Bamberger et al., 2003) resulting in reactive oxygen species (ROS) production, secretion of cytokines, and phagocytosis (Koenigsknecht and Landreth, 2004; Wilkinson et al., 2006).

Toll like receptors (TLRs) are expressed on cells of the innate immune system, and act to mobilize a robust immune reaction in response to pathogens (Akira, 2001; Medzhitov, 2001). TLRs function as dimers and often use co-receptors such as CD14 to assist in pathogen recognition. Specifically, CD14 interacts with TLR4 and TLR2-containing dimeric complexes to transduce activation signals in response to bacterial pathogens (Kielian, 2006). Importantly, TLRs have been found to associate with other receptors, such as CD36 and CD47 (Pfeiffer et al., 2001; Triantafilou et al., 2006) to elicit monocyte/macrophage activation (Triantafilou et al., 2004). A number of studies have reported an association between TLRs and AD, but have not defined a mechanistic link for how they participate in the microglial response to fA $\beta$  (Fassbender et al., 2004; Liu et al., 2005; Walter et al., 2007; Jana et al., 2008; Richard et al., 2008; Udan et al., 2008).

We report that microglia cultured from CD14-, TLR4-, or TLR2-null mice were unable to initiate signaling through a Src-Vav-Rac signaling cascade necessary for the induction of ROS and of phagocytosis following exposure to fA $\beta$ . Furthermore, fA $\beta$ -induced activation of p38 MAP kinase was abrogated in microglia deficient in CD14, TLR4, or TLR2, preventing subsequent ROS and phagocytosis. Together, these data indicate CD14, TLR4, and TLR2 functionally interact with other members of the microglial fA $\beta$  receptor complex to initiate intracellular signaling cascades resulting in an activated phenotype.

## Materials and Methods

### Reagents

A $\beta_{1-42}$  peptide was purchased from American Peptide Company (Sunnyvale, CA), dissolved in endotoxin-free sterile water and incubated at 37°C for 24 h to induce fibril formation. The initial concentration reflects that of the monomeric peptides used to generate the fibrils. This preparation has been well described (Burdick et al., 1992; Lorenzo and Yankner, 1994). *E. coli* K12 lipopolysaccharide (LPS) was purchased from List Biological Laboratories (Campbell, CA) and dissolved in endotoxin-free sterile water at 1 mg/ml. Pam3CSK4 purchased from InvivoGen (San Diego, CA) was dissolved in endotoxin-free sterile water. Phorbol 12-myristate 13-acetate (PMA) was purchased from Sigma (St. Louis, MO) and dissolved in ethanol. Immune IgG was prepared from goat anti-rabbit IgG from Sigma and rabbit anti-mouse IgG from Cappel (Aurora, OH) in a 5:1 ratio as described previously (Koenigsknecht and Landreth, 2004). Nitroblue tetrazolium chloride (NBT) was purchased from Roche (Basel, Switzerland). FluoSpheres (1  $\mu$ m microspheres) were purchased from Molecular Probes (Eugene, OR). The CD14 antibody MY4 was purchased from Beckman Coulter (Miami, FL). Function blocking antibodies to TLR4 (HTA-125), TLR2 (T2.5), CD47 (B6H12), and isotype control antibodies (IgG<sub>1</sub>, IgG<sub>2a</sub>, IgG<sub>2b</sub>) were purchased from eBioscience (San Diego, CA). The CD36 antibody FA6-152 was from Abcam (Cambridge, MA). Antibodies detecting phospho-Src, and phospho-p38, were purchased from Cell Signaling Technology (Danvers, MA). The anti-phospho-tyrosine antibody 4G10 and Rac

antibody were purchased from Upstate Biotechnologies (Lake Placid, NY). Antibodies to Phospho-I $\kappa$ B $\alpha$ , c-Src, Vav, p38, and actin were purchased from Santa Cruz Biotechnology Inc. (Santa Cruz, CA). The antibody to flotillin was purchased from BD Transduction Laboratories (San Jose, CA). Polymixin B was purchased from Sigma, and SB203580 from Invivogen.

### Cell culture

CD14<sup>-/-</sup> and C57BL/6J mice were from Jackson Labs (Bar Harbor, ME), TLR4<sup>-/-</sup> and TLR2<sup>-/-</sup> mice were generated by Shizuo Akira (Osaka University, Japan). Primary microglia were cultured from post-natal day 0-3 CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, TLR2<sup>-/-</sup>, or C57BL/6J mice as previously described by us (McDonald et al., 1997). Briefly, pups were decapitated, and meninges and blood vessels were completely removed from the cortices, which were minced and trypsinized with 0.05% Trypsin-EDTA (Gibco; Grand Island, NY) for 20 min at 37°C. DMEM/F12 (Gibco) containing 10% heat-inactivated fetal bovine serum (HI-FBS; Atlanta Biologicals; Lawrenceville, GA) and 1% penicillin-streptomycin (Gibco) was used to stop the digestion. Cells were then triturated and plated on 150 mm dishes (Corning; Corning, NY). Media was changed the following day to fresh DMEM/F12 with 10% HI-FBS for 14-21 days at 37°C, 5% CO<sub>2</sub>. Microglia were isolated as described by Saura et al. (Saura et al., 2003) by shaking the tissue culture plates for 30 min at room temperature to remove loosely adherent microglia. Astrocytes were next removed by 0.25% trypsin/HBSS diluted 1:4 in serum free DMEM/F12 for 1h at 37°C. Firmly attached microglia were harvested using 0.25% trypsin/PBS. Loosely and firmly adherent microglia were combined, and cells were counted and plated for experiments at the appropriate density in DMEM/F12 containing 2% HI-FBS overnight. Media was changed to serum free DMEM/F12 overnight prior to beginning the experiment.

The immortalized human monocyte cell line THP-1 from American Type Culture Collection (Manassas, VA) was maintained in RPMI 1640 (Gibco) containing 10% HI-FBS and gentamycin. The immortalized murine microglial cell line BV2 was maintained in DMEM (Gibco) containing 2% HI-FBS and gentamycin (Blasi et al., 1990).

### Cellular Adhesion Assay

Cellular binding to fA $\beta$  was performed as previously described (Bamberger et al., 2003). THP-1 monocytes (5 $\times$ 10<sup>5</sup> cells/condition) were incubated at 37°C for 30 min in serum free RPMI 1640 in the presence or absence of function blocking antibodies to CD14 (MY4, 10  $\mu$ g/ml), TLR4 (HTA125, 10  $\mu$ g/ml), TLR2 (T2.5, 10  $\mu$ g/ml), CD36 (FA6-152, 10  $\mu$ g/ml), CD47 (B6H12, 20  $\mu$ g/ml), their isotype controls (IgG<sub>1</sub>, IgG<sub>2a</sub>, or IgG<sub>2b</sub>), or polymixin B (0.1  $\mu$ g/ml). Fibrillar A $\beta$ <sub>1-42</sub> (2  $\mu$ g) was applied to a glass slide and allowed to dry. Cells (20,000) were applied to the slide and allowed to adhere for 5 min. Following washing with RPMI, the number of adherent cells was counted in three independent fields on an inverted microscope.

### Phagocytosis assay

Measurement of fA $\beta$ -stimulated microglial phagocytosis was performed as previously described (Koenigsnecht and Landreth, 2004). Microglia were plated at a density of 100,000 cells/well of a 24-well plate. SB203580 (0.5  $\mu$ M) or function blocking antibodies were added to serum free media for 30 min at 37°C. Media was collected, and fA $\beta$ <sub>1-42</sub> (2.5  $\mu$ M) was added for 30 min at 37°C. Fluorescent microspheres washed in PBS containing 1 mg/ml BSA were added to the cells for an additional 30 min. Cells were then fixed with 2% paraformaldehyde and the three random fields totaling at least 100 cells were counted for the fraction of cells containing microspheres on an inverted microscope.

### Reactive oxygen species assay

Measurement of intracellular superoxide radical generation was measured by the formation of a dark blue formazan deposit resulting from superoxide-mediated reduction of NBT (Pick, 1986; McDonald et al., 1997). Briefly, cells were plated at a density of 100,000 cells/well of a 24-well plate. SB203580 (0.5  $\mu$ M) was added to serum free media for 30 min at 37°C, 5% CO<sub>2</sub>. Media was collected and fA $\beta$ <sub>1-42</sub> (2.5  $\mu$ M), LPS (5  $\mu$ g/ml), Pam3CSK4 (1  $\mu$ M), or PMA (390 nM) was added to the cells for an additional 30 min at 37°C in the presence of 1 mg/ml NBT. Cells were fixed with 2% paraformaldehyde at 4°C for 30 min. Three random fields totaling at least 100 cells were counted on an inverted microscope and the fraction of ROS-positive cells was determined.

### Cell Stimulation, Immunoprecipitations and Western Blots

Cells were plated at  $2.5 \times 10^6$  cells/60 mm dish, and stimulated with fA $\beta$ <sub>1-42</sub> (2.5  $\mu$ M), LPS (5  $\mu$ g/ml), Pam3CSK4 (1  $\mu$ M), or immune IgG (250  $\mu$ g/ml) in serum free DMEM/F12 for 3 or 5 min at room temperature. Cells were collected and lysed in Triton buffer (1% Triton X-100, 20 mM Tris, pH 7.5, 100 mM NaCl, 40 mM NaF, 1 mM EDTA, 1 mM EGTA) containing the protease inhibitors leupeptin (1  $\mu$ g/ml), aprotinin (1  $\mu$ g/ml), PMSF (1 mM), and Na<sub>3</sub>VO<sub>4</sub> (1 mM). Lysates were cleared by centrifugation (15 min, 4°C, 500  $\times$  g). Protein concentration was determined by the BCA method (Pierce). Cellular lysates (0.5 mg) were added to 30  $\mu$ l Protein A agarose beads (Santa Cruz) and 2  $\mu$ g of the anti-Vav antibody per mg of lysate and rotated at 4°C for at least 2h. Samples were washed three times with Triton buffer, followed by the addition of sample buffer. Immunoprecipitations or aliquots of lysate (30  $\mu$ g protein) were boiled for 5 min, loaded onto 9% SDS-PAGE gels, and transferred to polyvinylidenedifluoride membranes (Millipore). Blots were probed with anti-phosphotyrosine (1:1000), -phospho-Src (1:1000), -phospho-p38 (1:1000), or -phospho-I $\kappa$ B $\alpha$  (1:1000) antibodies overnight at 4°C. The antigens were detected by anti-mouse or anti-rabbit HRP-conjugated secondary antibodies and visualized by enhanced chemiluminescence (Pierce, Rockford, IL). Blots were stripped and re-probed with antibodies to Vav (1:1000), c-Src (1:200), p38 (1:1000), or actin (1:1000) for evaluation of loading and immunoprecipitation efficiency. Band intensities were quantified using NIH Image J software (Bethesda, MD).

### Cellular Fractionation

The isolation of cytoplasmic and membrane fractions were performed as previously described (Wilkinson et al., 2006). Primary microglia were plated at  $2.5 \times 10^6$  cells/60 mm dish. Cells were stimulated with fA $\beta$ <sub>1-42</sub> (2.5  $\mu$ M) in serum free DMEM/F12 for 10 min at room temperature, collected and lysed in Relaxation buffer (100 mM KCl, 3 mM NaCl, 3.5 mM MgCl<sub>2</sub>, 1.25 mM EGTA, and 10 mM PIPES, pH 7.3), and incubated on ice for 15 min. Following a 10 s sonication, lysates were cleared by centrifugation at 500  $\times$  g, 4°C for 5 min. The supernatants were centrifuged for an hour at 110,000  $\times$  g, 4°C using a Beckman Coulter SW50.1 rotor. The supernatant (cytosolic fraction) was collected, and the resulting membrane pellet then resuspended in Relaxation buffer. The samples (30  $\mu$ g protein) were loaded onto a 12% SDS-PAGE resolving gel and the resulting blots were probed with an anti-Rac antibody (1:1000) and an anti-flotillin antibody (1:1000) to determine the relative amount of membrane-associated Rac. Band intensities were quantified using NIH Image J software.

### Statistical analysis

All replications of each experiment were pooled to generate graphical representations. GraphPad Prism 3.02 software was used to calculate mean values  $\pm$  SEM, and statistical differences were determined using a one way ANOVA with a Tukey post test to determine *p* values.

## Results

### CD14, TLR4, and TLR2 mediate binding of fA $\beta$

We have investigated the participation of TLRs and the co-receptor CD14 to stimulate microglial activation by testing whether CD14 physically interacted with fA $\beta$  and mediated the binding of the fibrils to the cell. THP-1 monocytes were employed as they are normally non-adherent and provide a sensitive system for assessing fA $\beta$ -binding. Monocytes avidly bind to fA $\beta$ ; however, pre-treatment of THP-1 monocytes with function blocking antibodies to CD14 (MY4), TLR4 (HTA125), or TLR2 (T2.5) dramatically inhibited the binding of monocytes to fA $\beta$  (Fig. 1A). Isotype control antibodies did not affect binding to fA $\beta$ , demonstrating the specificity of this effect. The association of these cells with fA $\beta$  was also blocked by function blocking antibodies to CD36 (FA6-152) and CD47 (B6H12) (Figure 1B), receptors that have been previously reported to be required for fA $\beta$  binding (Bamberger et al., 2003). Polymixin B, which binds the lipid A component of LPS to neutralize endotoxicity, had no effect on fA $\beta$  binding (Figure 1C), indicating that there was no trace contamination of CD14 ligands in the A $\beta$  or antibody preparations.

### CD14, TLR4, and TLR2 are required for a fA $\beta$ -induced phagocytic response

Phagocytosis is the process by which cells ingest large particles (>1  $\mu$ m) in an actin-dependent manner (Aderem and Underhill, 1999), and is important for both the degradation of infectious agents (Aderem, 2002) and tissue maintenance. Numerous studies have demonstrated a requirement for TLRs in initiating phagocytosis of target pathogens (Heine et al., 1999; Underhill et al., 1999; Tabora and Casadevall, 2002; Blander and Medzhitov, 2004; Sendide et al., 2005; Neal et al., 2006). Fibrillar A $\beta$  stimulates microglial phagocytosis in a time- and dose-dependent manner (Kopeck and Carroll, 1998). This response involves the microglial fA $\beta$  receptor complex via a  $\beta_1$ -integrin-dependent process (Koenigsnecht and Landreth, 2004) and requires Vav-mediated activation of Rac1 (Wilkinson et al., 2006). We wished to assess whether CD14, TLR4, and TLR2 participated in the initiation of this fA $\beta$ -stimulated response. BV2 microglia were pretreated with function blocking antibodies to CD14, TLR4, or TLR2 prior to stimulation with fA $\beta$ . Phagocytosis was monitored by the uptake of 1  $\mu$ m fluorescent beads (Figure 2A, B). While fA $\beta$  resulted in increased bead uptake in control cells, CD14, TLR4, or TLR2 function blocking antibodies abrogated this response (Figure 2A). Similarly, in primary microglial cultures prepared from wild-type, CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, or TLR2<sup>-/-</sup> animals, wild-type microglia demonstrated a robust phagocytic response following fA $\beta$  stimulation (Fig. 2C). In contrast, fA $\beta$  was unable to stimulate an increase in the phagocytic uptake of microspheres in CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, or TLR2<sup>-/-</sup> microglia (Figure 2C). These data indicate CD14, TLR4, and TLR2 are required at the cell surface to recognize fA $\beta$  and to initiate a phagocytic response.

### ROS production stimulated by fA $\beta$ requires CD14, TLR4, and TLR2

In order to kill pathogens, microglia use the NADPH oxidase to generate superoxide, as well as the more potent ROS which arise from this free radical, including hydrogen peroxide, hydroxyl radical, peroxynitrite, and other oxidants (Bergendi et al., 1999). Microglia exposed to fA $\beta$  generate a respiratory burst and subsequent superoxide anion release via the microglial fA $\beta$  cell surface receptor complex (Meda et al., 1996; McDonald et al., 1998; Bianca et al., 1999; Combs et al., 1999; Van Muiswinkel et al., 1999; Bamberger et al., 2003). We assessed the participation of CD14, TLR4, and TLR2 in this fA $\beta$ -stimulated response. Microglial cultures prepared from wild-type, CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, or TLR2<sup>-/-</sup> animals were treated with fA $\beta$ , and the reduction of NBT to an insoluble formazan precipitant was used as a measure of intracellular superoxide radical generation. While fA $\beta$  resulted in a 2-fold increase in ROS production in wild-type microglia, it failed to stimulate superoxide radical formation in CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, or TLR2<sup>-/-</sup> microglia (Fig. 3A). LPS and Pam3CSK4 were also able to drive

ROS production in wild-type microglia but not in CD14<sup>-/-</sup> (Fig. 3B), TLR4<sup>-/-</sup> (Fig. 3C), or TLR2<sup>-/-</sup> microglia (Fig. 3D). The ability of phorbol ester PMA to activate PKC leading to NADPH oxidase assembly (McDonald et al., 1997; Bianca et al., 1999; Bamberger et al., 2003) provides a positive control for these studies. CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, or TLR2<sup>-/-</sup> microglia show a robust induction in ROS production following PMA, in a manner comparable to their wild-type counterparts, indicating the NADPH oxidase machinery is intact and functional in the microglia derived from knock-out animals (Fig. 3B-D). These data suggest CD14, TLR4, and TLR2 function as members of the microglial fA $\beta$  cell surface receptor complex that is functionally linked to ROS production.

### Initiation of the Src-Vav-Rac intracellular signaling cascade requires CD14, TLR4, and TLR2

Engagement of the microglial fA $\beta$  receptor complex activates the Src family kinases Lyn and Fyn, and the Syk tyrosine kinase, which are required for subsequent signaling events (McDonald et al., 1997; Combs et al., 1999; Bamberger et al., 2003; Koenigsnecht and Landreth, 2004; Wilkinson et al., 2006). Interestingly, Src and Syk activation also occur following TLR engagement (English et al., 1993; Stefanova et al., 1993; Beaty et al., 1994). Therefore, we sought to determine whether CD14, TLR4, and TLR2 were required for the activation of Src following fA $\beta$  stimulation using an antibody that cross-reacts with activated forms of the Src family members Lyn, Fyn, Lck, Yes, and Hck. While fA $\beta$  stimulated a two-fold increase in level of phosphorylated Src kinases in wild-type microglia, it had no effect on phospho-Src levels in CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, or TLR2<sup>-/-</sup> microglia (Fig. 4A). Lipopolysaccharide (LPS), the canonical CD14 and TLR4 ligand, and Pam3CSK4, a synthetic TLR2-specific ligand, were also able to activate the Src kinases in wild-type microglia, but not in knock-out microglia (Fig. 4B-D) and provides a positive control for these studies. In contrast, immune IgG, which activates the Src kinase family through the Fc receptors (Sanchez-Mejorada and Rosales, 1998), was able to drive Src phosphorylation in CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, and TLR2<sup>-/-</sup> microglia in a manner comparable to that of wild-type microglia (Fig. 4B-D), indicating Src is functionally linked to these receptors and the lack of a response to fA $\beta$  is reflective of the requirement for these receptors in mediating microglial signaling in response to this specific ligand.

Activation of these tyrosine kinases by fA $\beta$  results in their association with the Rac-specific guanine nucleotide exchange factor Vav, and its subsequent phosphorylation (Wilkinson et al., 2006). To determine whether the downstream target of these kinases was affected, we examined Vav phosphorylation following fA $\beta$  treatment. Wild-type microglia respond to fA $\beta$  by a two-fold increase in Vav phosphorylation; however, microglia deficient in CD14, TLR4, or TLR2 did not exhibit Vav phosphorylation after fA $\beta$  exposure (Fig. 5A).

The downstream target of Vav, the small GTPase Rac1, is GTP-loaded in a fA $\beta$ -dependent manner and is translocated to the membrane where it mediates both actin reorganization necessary for phagocytosis and NADPH oxidase complex formation leading to ROS production (Wilkinson et al., 2006). We found that while fA $\beta$  stimulated a two-fold increase in membrane-associated Rac in wild-type microglia, it was unable to do so in CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, or TLR2<sup>-/-</sup> microglia (Fig. 5B). These data demonstrate that the absence of CD14, TLR4, or TLR2 from the microglial cell surface prevents the receptor complex from initiating the Src-Vav-Rac intracellular signaling cascade.

### Activation of p38 MAPK by fA $\beta$ is reduced in CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, and TLR2<sup>-/-</sup> microglia

The MAP kinase p38 is activated in microglia in response to fA $\beta$  *in vitro* (McDonald et al., 1997; Pyo et al., 1998). Activated forms of p38 are found in the brains of AD mouse models and AD patients (Hensley et al., 1999; Zhu et al., 2000; Giovannini et al., 2002; Savage et al., 2002; Sun et al., 2003; Ferrer, 2004; Jin et al., 2005). Moreover, p38 is activated following

engagement of the TLRs in a manner similar to that elicited by fA $\beta$  exposure (Bhat et al., 1998; Pyo et al., 1998; Kim et al., 2004). We assessed levels of phosphorylated p38 in CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, and TLR2<sup>-/-</sup> microglia following fA $\beta$  treatment, and found that, compared to their wild-type counterparts, these microglia failed to show an increase in phospho-p38 (Fig. 6A). Similarly, CD14/TLR4 and TLR2 ligands LPS (Fig. 6B, C) and Pam3CSK4 (Fig. 6D) increased the levels of phosphorylated p38 in wild-type microglia, but not in CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, and TLR2<sup>-/-</sup> microglia. However, immune IgG was able to induce levels of phospho-p38 in the receptor-deficient cells (Fig. 6B-D) as previously reported (Yamamori et al., 2000), indicating p38 MAP kinase signaling pathways are intact in these cells and the failure to activate p38 is specifically abrogated in response to fA $\beta$ .

### Inhibition of p38 attenuates fA $\beta$ -stimulated ROS production and phagocytosis

We used a p38-specific inhibitor, SB203580, to elucidate the role of p38 in fA $\beta$ -induced ROS production. Previous work has shown that p38 phosphorylates the p47<sup>phox</sup> subunit of the NADPH oxidase complex (El Benna et al., 1996; Dang et al., 2006). Phosphorylation of p47<sup>phox</sup> serine residues exposes binding domains that interact with other cytoplasmic NADPH oxidase subunits and phosphoinositide lipid products, resulting in p47<sup>phox</sup> translocation to the membrane, where it participates in NADPH oxidase complex assembly (DeLeo and Quinn, 1996; Sumimoto et al., 1996; Ago et al., 1999; Babior, 1999; Ago et al., 2003). Significantly, this occurs following LPS engagement of CD14 (Yan et al., 2002; Laroux et al., 2005). The p38 inhibitor SB203580 prevented fA $\beta$ -induced ROS production (Fig. 7A), consistent with a previous report using a different p38 inhibitor (Smits et al., 2001).

Next, the role of p38 in fA $\beta$ -induced phagocytosis was assessed. Activation of p38 MAPK is required for cytoskeleton re-arrangement and subsequent autophagy (Park et al., 2008; Tang et al., 2008). The small GTPase Rac has been implicated in membrane ruffling and phagocytosis (Ridley et al., 1992; Cox et al., 1997), and its activation by GTP-loading and cytosol to membrane translocation is regulated by p38 activity (El Bekay et al., 2007; Zuluaga et al., 2007; Osada et al., 2009). To assess whether fA $\beta$ -stimulated p38 activation via the fA $\beta$  receptor complex modulated phagocytosis, we treated microglia with the p38 inhibitor SB203580 prior to fA $\beta$  stimulation. Consistent with work that showed inhibition of p38 attenuated LPS-driven phagocytosis (Kong and Ge, 2008), we found that SB203580 reduced fA $\beta$ -stimulated microsphere uptake (Fig. 7B).

### CD14, TLR4, and TLR2 in fA $\beta$ -induced NF $\kappa$ B activation

Engagement of CD14 and Toll like receptors are known to stimulate NF $\kappa$ B-dependent transcription (Akira, 2001; Medzhitov, 2001). Fibrillar A $\beta$  has also been shown to activate NF $\kappa$ B in monocytes and microglia (Combs et al., 2001). Inhibition of CD14, TLR4, and TLR2 prevents A $\beta$ -stimulated NF $\kappa$ B-dependent gene induction (Fassbender et al., 2004; Walter et al., 2007; Jana et al., 2008; Udan et al., 2008). NF $\kappa$ B is held in an inactive state in the cytosol by its participation in a complex with the inhibitory I $\kappa$ B proteins (Baeuerle and Baltimore, 1988; Beg and Baldwin, 1993; Finco et al., 1994). I $\kappa$ B $\alpha$  is phosphorylated at Ser32 and Ser36, resulting in dissolution of the complex and targeting I $\kappa$ B $\alpha$  for proteasome-mediated degradation, and thereby permitting NF $\kappa$ B nuclear translocation for transcription (Finco et al., 1994; Brockman et al., 1995; Brown et al., 1995; Traenckner et al., 1995; Chen et al., 1996). We assessed NF $\kappa$ B activation by measuring I $\kappa$ B $\alpha$  phosphorylation following fA $\beta$  stimulation. Wild-type microglia responded to fA $\beta$  with a two-fold increase in phospho-I $\kappa$ B $\alpha$  levels; however, CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, or TLR2<sup>-/-</sup> deficient microglia did not show increased I $\kappa$ B $\alpha$  phosphorylation following fA $\beta$  exposure. These data are consistent with previous reports demonstrating a requirement for CD14, TLR4, and TLR2 in fA $\beta$ -stimulated NF $\kappa$ B-dependent transcription (Fassbender et al., 2004; Walter et al., 2007; Jana et al., 2008; Udan et al., 2008).

## Discussion

The inflammatory component of AD reflects the response of microglial cells to deposits of fA $\beta$ . The current study demonstrates an essential role for canonical innate immune receptors in the induction of the microglial inflammatory response. These results are consistent with findings of how immune cells recognize pathogens and other complex entities, namely through the use of receptor ensembles. Immune cells often use ensembles of receptors to bind fibrillar proteins and other complex macromolecules (Ishibashi et al., 1994; Bornstein, 1995; Wong et al., 1996). TLRs act in concert with other immune receptors to form large receptor complexes whose composition is governed by the ligand that is engaged (Pfeiffer et al., 2001; Triantafilou et al., 2006). This combinatorial use of cell surface receptors determines the character of the immune response (Triantafilou et al., 2004). We have previously described a multi-receptor complex on the surface of microglia consisting of the  $\alpha_6\beta_1$  integrin, the integrin associated protein CD47, the B-class scavenger receptor CD36, and SRA (Bamberger et al., 2003). We demonstrate an obligatory role for CD14 in microglial recognition of fA $\beta$  and mounting a cellular response to this ligand. The roles of CD14 and scavenger receptors have been confounded by the use of fucoidan, which has historically been used to inhibit the binding of various ligands to scavenger receptors. It is well appreciated that it also inhibits the ability of CD14 to bind its ligands (Heinzelmann et al., 1998; Gregory and Devitt, 1999; van Oosten et al., 2001). Many of the reports of SRA-dependent signaling have been misinterpreted and are due to CD14 interacting with other scavenger receptor ligands (Kim et al., 2003). Importantly, CD14 has been shown to associate with CD36 and CD47 in response to various ligands (Pfeiffer et al., 2001) and bind fibrillar forms of A $\beta$  (Fassbender et al., 2004), observations consistent with our postulate that these receptors act in concert to bind fA $\beta$  and initiate intracellular signaling. The well documented functional interactions of CD14 with TLR4 or TLR2 led us to investigate their participation in a receptor complex mediating the microglial response to fA $\beta$ .

We report that CD14, TLR4, and TLR2 are necessary for binding fA $\beta$  at the cell surface, and are required for phenotypic activation of microglia and induction of phagocytosis. Microglia are competent phagocytes, reflective of their roles as the brain's principal immune effector cell (Streit et al., 2004). We have previously demonstrated the fA $\beta$ -stimulated induction of phagocytosis via the fA $\beta$  receptor complex that was reliant upon the action of the  $\beta_1$  integrin (Koenigsknecht and Landreth, 2004). Similarly, microglial activation by TLR ligands was reported to stimulate A $\beta$  uptake (Tahara et al., 2006), although it is unclear if this is through macropinocytosis (Mandrekar et al., 2009) or phagocytosis of A $\beta$ . One caveat with this and other in vitro studies is that they employ microglia derived from neonatal animals. A significant issue in the interpretation of these data is that they might not accurately reflect the activities of these cells in vivo in aged animals. Moreover, there is evidence that microglia from aged animals have reduced phagocytic capacity and blunted induction of cytokine expression (Floden and Combs, 2006; Sierra et al., 2007). Indeed, work by Streit and colleagues has demonstrated that microglia can undergo senescence in aged animals (Flanary et al., 2007). However, a recent study by Sierra et al. suggests that the effect of aging is principally to reduce the magnitude of microglial responses to inflammatory stimuli, without changing the nature or specificity of the response (Sierra et al., 2007).

Fibrillar A $\beta$  and TLR ligands both stimulate NADPH oxidase to form superoxide and more potent ROS. We demonstrated that CD14, TLR4, and TLR2 are necessary for fA $\beta$  to stimulate this response. These findings differ from those reported by Qin et al. who found LPS-induced superoxide production by microglia was independent of TLR4 (Qin et al., 2005). There are two significant differences in the studies that may explain this discrepancy. Qin et al. used C3H/HeJ mice, which express a mutated and functionally impaired form of TLR4, while the mice used here were TLR4-null and on a B6 background. Moreover, these two studies



employed different forms of LPS, and LPS structure can influence the immune pathways activated by the various species (Gangloff et al., 1999). Different LPS analogues trigger the recruitment of different receptor ensembles within membrane microdomains, and the composition of each receptor cluster determines the nature of the immune response (Triantafilou et al., 2004).

Signal transduction cascades activated by TLR engagement are indistinguishable from those activated by fA $\beta$ . (English et al., 1993; Stefanova et al., 1993; Beaty et al., 1994; Geng et al., 1994; Arbibe et al., 2000; Lee et al., 2000; Sanlioglu et al., 2001; Wilkinson et al., 2006). Therefore, we tested if CD14, TLR4, and TLR2 participated in fA $\beta$ -stimulated signaling. Significantly, we found that each of these receptors were required. Thus, these innate immune receptors not only function in binding fA $\beta$ , but are essential for the fA $\beta$ -stimulated activation of the Src-Vav-Rac signaling cascade, just as are other members of the fA $\beta$  receptor complex. A primary downstream target of both fA $\beta$  receptor signaling and TLRs is the p38 MAPK (McDonald et al., 1997; Pyo et al., 1998). We demonstrate CD14, TLR4, and TLR2 were required for fA $\beta$ -induced activation of p38 which is required for fA $\beta$ -induced reactive oxygen species production. We also demonstrate that inhibition of p38 activity reduces fA $\beta$ -stimulated phagocytosis. These observations are consistent with the roles of p38 in the immune response.

We report a requirement for CD14, TLR4, and TLR2 in NF $\kappa$ B activation following fA $\beta$ . NF $\kappa$ B becomes transcriptionally active following phosphorylation of its inhibitor I $\kappa$ B $\alpha$ , which targets it for degradation, permitting nuclear translocation of NF $\kappa$ B (Bauerle and Baltimore, 1988; Beg and Baldwin, 1993; Finco et al., 1994; Brockman et al., 1995; Brown et al., 1995; Traenckner et al., 1995; Chen et al., 1996). Fibrillar A $\beta$  has been shown to stimulate NF $\kappa$ B-dependent gene expression via CD14, TLR4, and TLR2 (Fassbender et al., 2004; Walter et al., 2007; Jana et al., 2008; Udan et al., 2008). We extend these studies by showing these receptors are also required for the activation of I $\kappa$ B $\alpha$ , which is required for NF $\kappa$ B nuclear translocation and subsequent target gene transcription.

A number of recent studies have assessed whether the innate immune system is involved in AD. There is increased expression of CD14, TLR2, and TLR4 in the brains both animal models of AD as well as AD patients (Fassbender et al., 2004; Liu et al., 2005; Letiembre et al., 2007; Walter et al., 2007). Moreover, a polymorphism in the TLR4 gene results in a blunted signaling response and is correlated with a 2.7-fold reduction in risk for late onset AD (Minoretti et al., 2006). CD14, when over-expressed in CHO cells, has been shown to bind A $\beta$  (Fassbender et al., 2004). This co-receptor has also been shown to be necessary for release of nitrite and pro-inflammatory gene expression from microglia (Fassbender et al., 2004; Liu et al., 2005). There is evidence that TLRs are involved in the microglial response to A $\beta$  as inhibition of TLR4 or TLR2 through function blocking antibodies or siRNA knock-down also prevented fA $\beta$ -induced nitrite, IL-6, and TNF- $\alpha$  production (Walter et al., 2007; Jana et al., 2008; Udan et al., 2008). These effects were dependent on the fibrillar structure of A $\beta$  (Fassbender et al., 2004; Walter et al., 2007). The role of TLRs in A $\beta$  clearance *in vivo* was examined by two separate studies with seemingly different outcomes. Fourteen to 16 month old APP<sup>swe</sup>/PSEN<sup>dE9</sup> mice with inactive TLR4 showed increased cortical and hippocampal A $\beta$  burden, without changing steady state APP or presenilin1 levels, arguing that the change in A $\beta$  load was due to a change in microglial-mediated A $\beta$  clearance that is reliant upon TLR4 function (Tahara et al., 2006). In contrast, APP<sup>swe</sup>/PSEN<sup>dE9</sup> mice lacking TLR2 had delayed plaque formation through 6 months of age, but by 9 months the plaque deposition was comparable to their wild-type counterparts (Richard et al., 2008). This suggests that during early stages of AD, TLR-independent A $\beta$  clearance mechanisms are effective in removing A $\beta$  from the brain, and potentially, the absence of TLR-mediated inflammation makes these processes more efficient. However, over the course of time, the rate of A $\beta$  production overwhelms these processes and the lack of TLRs enhances A $\beta$  deposition. A recent study by

Scholtzova et al. reported that intraperitoneal injections of TLR9 ligands into a murine model of AD resulted in behavioral improvement and reduction of amyloid burden in the brain, however, the mechanism through which the effects are achieved are unknown since the TLR9 ligands are not likely to have entered the brain (Scholtzova et al., 2009).

The present study places the innate immune receptors CD14, TLR4, and TLR2 at the interface of microglia and fA $\beta$ , consistent with previous reports. Importantly, this study extends what is known about the mechanisms behind these receptors in microglial activation. CD14, TLR4, and TLR2 function along with CD36, the  $\alpha_6\beta_1$  integrin, and CD47 as members of the fA $\beta$  receptor complex whose engagement initiates the parallel signaling cascades of Src-Vav-Rac and the p38 MAP kinase for the induction of ROS, phagocytosis, and NF $\kappa$ B activation. While an acute inflammatory response eliminates pathogens and promotes tissue repair and is therefore neuroprotective, chronic inflammation results in progressive damage and is associated with neurotoxicity (Akiyama et al., 2000). These findings provide an understanding for the role of the innate immune system in the acute inflammatory reaction to fA $\beta$ , allowing for the development of effective therapeutic strategies to control the inflammatory aspect of AD.

## Acknowledgments

This work was supported by the NIH (AG16047) and the Neuroscience Training Grant (5 T32 AG0071), the Blanchette Hooker Rockefeller Foundation, and the American Health Assistance Foundation. Erin Reed-Geaghan is supported by a pre-doctoral Ruth L. Kirschstein National Research Service Award (F31NS057867) from the NINDS. We would like to thank Dr. Shizuo Akira for generously sharing TLR2<sup>-/-</sup> and TLR4<sup>-/-</sup> mice.

## References

- Aderem A. How to eat something bigger than your head. *Cell* 2002;110:5–8. [PubMed: 12150991]
- Aderem A, Underhill DM. Mechanisms of phagocytosis in macrophages. *Annu Rev Immunol* 1999;17:593–623. [PubMed: 10358769]
- Ago T, Nunoi H, Ito T, Sumimoto H. Mechanism for phosphorylation-induced activation of the phagocyte NADPH oxidase protein p47(phox). Triple replacement of serines 303, 304, and 328 with aspartates disrupts the SH3 domain-mediated intramolecular interaction in p47(phox), thereby activating the oxidase. *J Biol Chem* 1999;274:33644–33653. [PubMed: 10559253]
- Ago T, Kuribayashi F, Hiroaki H, Takeya R, Ito T, Kohda D, Sumimoto H. Phosphorylation of p47phox directs phox homology domain from SH3 domain toward phosphoinositides, leading to phagocyte NADPH oxidase activation. *Proc Natl Acad Sci U S A* 2003;100:4474–4479. [PubMed: 12672956]
- Akira S. Toll-like receptors and innate immunity. *Adv Immunol* 2001;78:1–56. [PubMed: 11432202]
- Akiyama H, Barger S, Barnum S, Bradt B, Bauer J, Cole GM, Cooper NR, Eikelenboom P, Emmerling M, Fiebich BL, Finch CE, Frautschy S, Griffin WS, Hampel H, Hull M, Landreth G, Lue L, Mrak R, Mackenzie IR, McGeer PL, O'Banion MK, Pachter J, Pasinetti G, Plata-Salaman C, Rogers J, Rydel R, Shen Y, Streit W, Strohmeyer R, Tooyoma I, Van Muiswinkel FL, Veerhuis R, Walker D, Webster S, Wegrzyniak B, Wenk G, Wyss-Coray T. Inflammation and Alzheimer's disease. *Neurobiol Aging* 2000;21:383–421. [PubMed: 10858586]
- Arbibe L, Mira JP, Teusch N, Kline L, Guha M, Mackman N, Godowski PJ, Ulevitch RJ, Knaus UG. Toll-like receptor 2-mediated NF-kappa B activation requires a Rac1-dependent pathway. *Nat Immunol* 2000;1:533–540. [PubMed: 11101877]
- Babior BM. NADPH oxidase: an update. *Blood* 1999;93:1464–1476. [PubMed: 10029572]
- Baeuerle PA, Baltimore D. I kappa B: a specific inhibitor of the NF-kappa B transcription factor. *Science* 1988;242:540–546. [PubMed: 3140380]
- Bamberger ME, Harris ME, McDonald DR, Husemann J, Landreth GE. A cell surface receptor complex for fibrillar beta-amyloid mediates microglial activation. *J Neurosci* 2003;23:2665–2674. [PubMed: 12684452]

- Beaty CD, Franklin TL, Uehara Y, Wilson CB. Lipopolysaccharide-induced cytokine production in human monocytes: role of tyrosine phosphorylation in transmembrane signal transduction. *Eur J Immunol* 1994;24:1278–1284. [PubMed: 7515809]
- Beg AA, Baldwin AS Jr. The I kappa B proteins: multifunctional regulators of Rel/NF-kappa B transcription factors. *Genes Dev* 1993;7:2064–2070. [PubMed: 8224838]
- Bergendi L, Benes L, Durackova Z, Ferencik M. Chemistry, physiology and pathology of free radicals. *Life Sci* 1999;65:1865–1874. [PubMed: 10576429]
- Bhat NR, Zhang P, Lee JC, Hogan EL. Extracellular signal-regulated kinase and p38 subgroups of mitogen-activated protein kinases regulate inducible nitric oxide synthase and tumor necrosis factor-alpha gene expression in endotoxin-stimulated primary glial cultures. *J Neurosci* 1998;18:1633–1641. [PubMed: 9464988]
- Bianca VD, Dusi S, Bianchini E, Dal Pra I, Rossi F. beta-amyloid activates the O-2 forming NADPH oxidase in microglia, monocytes, and neutrophils. A possible inflammatory mechanism of neuronal damage in Alzheimer's disease. *J Biol Chem* 1999;274:15493–15499. [PubMed: 10336441]
- Blander JM, Medzhitov R. Regulation of phagosome maturation by signals from toll-like receptors. *Science* 2004;304:1014–1018. [PubMed: 15143282]
- Blasi E, Barluzzi R, Bocchini V, Mazzolla R, Bistoni F. Immortalization of murine microglial cells by a v-raf/v-myc carrying retrovirus. *J Neuroimmunol* 1990;27:229–237. [PubMed: 2110186]
- Bornemann KD, Wiederhold KH, Pauli C, Ermini F, Stalder M, Schnell L, Sommer B, Jucker M, Staufenbiel M. Abeta-induced inflammatory processes in microglia cells of APP23 transgenic mice. *Am J Pathol* 2001;158:63–73. [PubMed: 11141480]
- Bornstein P. Diversity of function is inherent in matricellular proteins: an appraisal of thrombospondin 1. *J Cell Biol* 1995;130:503–506. [PubMed: 7542656]
- Brockman JA, Scherer DC, McKinsey TA, Hall SM, Qi X, Lee WY, Ballard DW. Coupling of a signal response domain in I kappa B alpha to multiple pathways for NF-kappa B activation. *Mol Cell Biol* 1995;15:2809–2818. [PubMed: 7739562]
- Brown K, Gerstberger S, Carlson L, Franzoso G, Siebenlist U. Control of I kappa B-alpha proteolysis by site-specific, signal-induced phosphorylation. *Science* 1995;267:1485–1488. [PubMed: 7878466]
- Burdick D, Soreghan B, Kwon M, Kosmoski J, Knauer M, Henschen A, Yates J, Cotman C, Glabe C. Assembly and aggregation properties of synthetic Alzheimer's A4/beta amyloid peptide analogs. *J Biol Chem* 1992;267:546–554. [PubMed: 1730616]
- Chen ZJ, Parent L, Maniatis T. Site-specific phosphorylation of IkappaBalpha by a novel ubiquitination-dependent protein kinase activity. *Cell* 1996;84:853–862. [PubMed: 8601309]
- Combs CK, Karlo JC, Kao SC, Landreth GE. beta-Amyloid stimulation of microglia and monocytes results in TNFalpha-dependent expression of inducible nitric oxide synthase and neuronal apoptosis. *J Neurosci* 2001;21:1179–1188. [PubMed: 11160388]
- Combs CK, Johnson DE, Cannady SB, Lehman TM, Landreth GE. Identification of microglial signal transduction pathways mediating a neurotoxic response to amyloidogenic fragments of beta-amyloid and prion proteins. *J Neurosci* 1999;19:928–939. [PubMed: 9920656]
- Cox D, Chang P, Zhang Q, Reddy PG, Bokoch GM, Greenberg S. Requirements for both Rac1 and Cdc42 in membrane ruffling and phagocytosis in leukocytes. *J Exp Med* 1997;186:1487–1494. [PubMed: 9348306]
- Dang PM, Stensballe A, Boussetta T, Raad H, Dewas C, Kroviarski Y, Hayem G, Jensen ON, Gougerot-Pocidal MA, El-Benna J. A specific p47phox -serine phosphorylated by convergent MAPKs mediates neutrophil NADPH oxidase priming at inflammatory sites. *J Clin Invest* 2006;116:2033–2043. [PubMed: 16778989]
- Davalos D, Grutzendler J, Yang G, Kim JV, Zuo Y, Jung S, Littman DR, Dustin ML, Gan WB. ATP mediates rapid microglial response to local brain injury in vivo. *Nat Neurosci* 2005;8:752–758. [PubMed: 15895084]
- DeLeo FR, Quinn MT. Assembly of the phagocyte NADPH oxidase: molecular interaction of oxidase proteins. *J Leukoc Biol* 1996;60:677–691. [PubMed: 8975869]
- El Bekay R, Alba G, Reyes ME, Chacon P, Vega A, Martin-Nieto J, Jimenez J, Ramos E, Olivan J, Pintado E, Sobrino F. Rac2 GTPase activation by angiotensin II is modulated by Ca2+/calcineurin


and mitogen-activated protein kinases in human neutrophils. *J Mol Endocrinol* 2007;39:351–363. [PubMed: 17975262]

- El Benna J, Faust RP, Johnson JL, Babior BM. Phosphorylation of the respiratory burst oxidase subunit p47phox as determined by two-dimensional phosphopeptide mapping. Phosphorylation by protein kinase C, protein kinase A, and a mitogen-activated protein kinase. *J Biol Chem* 1996;271:6374–6378. [PubMed: 8626435]
- English BK, Ihle JN, Myracle A, Yi T. Hck tyrosine kinase activity modulates tumor necrosis factor production by murine macrophages. *J Exp Med* 1993;178:1017–1022. [PubMed: 8350043]
- Fassbender K, Walter S, Kuhl S, Landmann R, Ishii K, Bertsch T, Stalder AK, Muehlhauser F, Liu Y, Ulmer AJ, Rivest S, Lentschat A, Gulbins E, Jucker M, Staufenbiel M, Brechtel K, Walter J, Multhaup G, Penke B, Adachi Y, Hartmann T, Beyreuther K. The LPS receptor (CD14) links innate immunity with Alzheimer's disease. *Faseb J* 2004;18:203–205. [PubMed: 14597556]
- Ferrer I. Stress kinases involved in tau phosphorylation in Alzheimer's disease, tauopathies and APP transgenic mice. *Neurotox Res* 2004;6:469–475. [PubMed: 15658002]
- Finco TS, Beg AA, Baldwin AS Jr. Inducible phosphorylation of I kappa B alpha is not sufficient for its dissociation from NF-kappa B and is inhibited by protease inhibitors. *Proc Natl Acad Sci U S A* 1994;91:11884–11888. [PubMed: 7991551]
- Flanary BE, Sammons NW, Nguyen C, Walker D, Streit WJ. Evidence that aging and amyloid promote microglial cell senescence. *Rejuvenation Res* 2007;10:61–74. [PubMed: 17378753]
- Floden AM, Combs CK. Beta-amyloid stimulates murine postnatal and adult microglia cultures in a unique manner. *J Neurosci* 2006;26:4644–4648. [PubMed: 16641245]
- Gangloff SC, Hijiya N, Haziot A, Goyert SM. Lipopolysaccharide structure influences the macrophage response via CD14-independent and CD14-dependent pathways. *Clin Infect Dis* 1999;28:491–496. [PubMed: 10194066]
- Geng Y, Gulbins E, Altman A, Lotz M. Monocyte deactivation by interleukin 10 via inhibition of tyrosine kinase activity and the Ras signaling pathway. *Proc Natl Acad Sci U S A* 1994;91:8602–8606. [PubMed: 8078929]
- Giovannini MG, Scali C, Prosperi C, Bellucci A, Vannucchi MG, Rosi S, Pepeu G, Casamenti F. Beta-amyloid-induced inflammation and cholinergic hypofunction in the rat brain in vivo: involvement of the p38MAPK pathway. *Neurobiol Dis* 2002;11:257–274. [PubMed: 12505419]
- Gregory CD, Devitt A. CD14 and apoptosis. *Apoptosis* 1999;4:11–20. [PubMed: 14634291]
- Heine H, Kirschning CJ, Lien E, Monks BG, Rothe M, Golenbock DT. Cutting edge: cells that carry A null allele for toll-like receptor 2 are capable of responding to endotoxin. *J Immunol* 1999;162:6971–6975. [PubMed: 10358136]
- Heinzelmann M, Polk HC Jr, Miller FN. Modulation of lipopolysaccharide-induced monocyte activation by heparin-binding protein and fucoidan. *Infect Immun* 1998;66:5842–5847. [PubMed: 9826363]
- Hensley K, Floyd RA, Zheng NY, Nael R, Robinson KA, Nguyen X, Pye QN, Stewart CA, Geddes J, Markesbery WR, Patel E, Johnson GV, Bing G. p38 kinase is activated in the Alzheimer's disease brain. *J Neurochem* 1999;72:2053–2058. [PubMed: 10217284]
- Ishibashi Y, Claus S, Relman DA. Bordetella pertussis filamentous hemagglutinin interacts with a leukocyte signal transduction complex and stimulates bacterial adherence to monocyte CR3 (CD11b/CD18). *J Exp Med* 1994;180:1225–1233. [PubMed: 7931059]
- Itagaki S, McGeer PL, Akiyama H, Zhu S, Selkoe D. Relationship of microglia and astrocytes to amyloid deposits of Alzheimer disease. *J Neuroimmunol* 1989;24:173–182. [PubMed: 2808689]
- Jana M, Palencia CA, Pahan K. Fibrillar amyloid-beta peptides activate microglia via TLR2: implications for Alzheimer's disease. *J Immunol* 2008;181:7254–7262. [PubMed: 18981147]
- Jin Y, Yan EZ, Fan Y, Zong ZH, Qi ZM, Li Z. Sodium ferulate prevents amyloid-beta-induced neurotoxicity through suppression of p38 MAPK and upregulation of ERK-1/2 and Akt/protein kinase B in rat hippocampus. *Acta Pharmacol Sin* 2005;26:943–951. [PubMed: 16038626]
- Kielian T. Toll-like receptors in central nervous system glial inflammation and homeostasis. *J Neurosci Res* 2006;83:711–730. [PubMed: 16541438]
- Kim SH, Smith CJ, Van Eldik LJ. Importance of MAPK pathways for microglial pro-inflammatory cytokine IL-1 beta production. *Neurobiol Aging* 2004;25:431–439. [PubMed: 15013563]

- Kim WS, Ordija CM, Freeman MW. Activation of signaling pathways by putative scavenger receptor class A (SR-A) ligands requires CD14 but not SR-A. *Biochem Biophys Res Commun* 2003;310:542–549. [PubMed: 14521944]
- Koenigsnecht J, Landreth G. Microglial phagocytosis of fibrillar beta-amyloid through a beta1 integrin-dependent mechanism. *J Neurosci* 2004;24:9838–9846. [PubMed: 15525768]
- Kong L, Ge BX. MyD88-independent activation of a novel actin-Cdc42/Rac pathway is required for Toll-like receptor-stimulated phagocytosis. *Cell Res* 2008;18:745–755. [PubMed: 18542102]
- Kopec KK, Carroll RT. Alzheimer's beta-amyloid peptide 1-42 induces a phagocytic response in murine microglia. *J Neurochem* 1998;71:2123–2131. [PubMed: 9798938]
- Laroux FS, Romero X, Wetzler L, Engel P, Terhorst C. Cutting edge: MyD88 controls phagocyte NADPH oxidase function and killing of gram-negative bacteria. *J Immunol* 2005;175:5596–5600. [PubMed: 16237045]
- Lee DJ, Cox D, Li J, Greenberg S. Rac1 and Cdc42 are required for phagocytosis, but not NF-kappaB-dependent gene expression, in macrophages challenged with *Pseudomonas aeruginosa*. *J Biol Chem* 2000;275:141–146. [PubMed: 10617597]
- Letiembre M, Liu Y, Walter S, Hao W, Pfander T, Wrede A, Schulz-Schaeffer W, Fassbender K. Screening of innate immune receptors in neurodegenerative diseases: A similar pattern. *Neurobiol Aging*. 2007
- Liu Y, Walter S, Stagi M, Cherny D, Letiembre M, Schulz-Schaeffer W, Heine H, Penke B, Neumann H, Fassbender K. LPS receptor (CD14): a receptor for phagocytosis of Alzheimer's amyloid peptide. *Brain* 2005;128:1778–1789. [PubMed: 15857927]
- Lorenzo A, Yankner BA. Beta-amyloid neurotoxicity requires fibril formation and is inhibited by congo red. *Proc Natl Acad Sci U S A* 1994;91:12243–12247. [PubMed: 7991613]
- Mandrekar S, Jiang Q, Lee CY, Koenigsnecht-Talboo J, Holtzman DM, Landreth GE. Microglia mediate the clearance of soluble A $\beta$  through fluid phase macropinocytosis. *J Neurosci* 2009;29:4252–4262. [PubMed: 19339619]
- McDonald DR, Brunden KR, Landreth GE. Amyloid fibrils activate tyrosine kinase-dependent signaling and superoxide production in microglia. *J Neurosci* 1997;17:2284–2294. [PubMed: 9065490]
- McDonald DR, Bamberger ME, Combs CK, Landreth GE. beta-Amyloid fibrils activate parallel mitogen-activated protein kinase pathways in microglia and THP1 monocytes. *J Neurosci* 1998;18:4451–4460. [PubMed: 9614222]
- Meda L, Bonaiuto C, Baron P, Otvos L Jr, Rossi F, Cassatella MA. Priming of monocyte respiratory burst by beta-amyloid fragment (25-35). *Neurosci Lett* 1996;219:91–94. [PubMed: 8971787]
- Medzhitov R. Toll-like receptors and innate immunity. *Nat Rev Immunol* 2001;1:135–145. [PubMed: 11905821]
- Minoretti P, Gazzaruso C, Vito CD, Emanuele E, Bianchi M, Coen E, Reino M, Geroldi D. Effect of the functional toll-like receptor 4 Asp299Gly polymorphism on susceptibility to late-onset Alzheimer's disease. *Neurosci Lett* 2006;391:147–149. [PubMed: 16157451]
- Neal MD, Leaphart C, Levy R, Prince J, Billiar TR, Watkins S, Li J, Cetin S, Ford H, Schreiber A, Hackam DJ. Enterocyte TLR4 mediates phagocytosis and translocation of bacteria across the intestinal barrier. *J Immunol* 2006;176:3070–3079. [PubMed: 16493066]
- Nimmerjahn A, Kirchhoff F, Helmchen F. Resting microglial cells are highly dynamic surveillants of brain parenchyma in vivo. *Science* 2005;308:1314–1318. [PubMed: 15831717]
- Osada Y, Sunatani T, Kim IS, Nakanishi Y, Shiratsuchi A. Signaling pathway involving GULP, MAPK, and Rac1 for SR-BI-induced phagocytosis of apoptotic cells. *J Biochem*. 2009
- Park JY, Kim HY, Jou I, Park SM. GM1 induces p38 and microtubule dependent ramification of rat primary microglia in vitro. *Brain Res* 2008;1244:13–23. [PubMed: 18930716]
- Perlmutter LS, Barron E, Chui HC. Morphologic association between microglia and senile plaque amyloid in Alzheimer's disease. *Neurosci Lett* 1990;119:32–36. [PubMed: 2097581]
- Pfeiffer A, Bottcher A, Orso E, Kapinsky M, Nagy P, Bodnar A, Spreitzer I, Liebisch G, Drobnik W, Gempel K, Horn M, Holmer S, Hartung T, Multhoff G, Schutz G, Schindler H, Ulmer AJ, Heine H, Stelter F, Schutt C, Rothe G, Szollosi J, Damjanovich S, Schmitz G. Lipopolysaccharide and ceramide docking to CD14 provokes ligand-specific receptor clustering in rafts. *Eur J Immunol* 2001;31:3153–3164. [PubMed: 11745332]

- Pick E. Microassays for superoxide and hydrogen peroxide production and nitroblue tetrazolium reduction using an enzyme immunoassay microplate reader. *Methods Enzymol* 1986;132:407–421. [PubMed: 3029539]
- Pyo H, Jou I, Jung S, Hong S, Joe EH. Mitogen-activated protein kinases activated by lipopolysaccharide and beta-amyloid in cultured rat microglia. *Neuroreport* 1998;9:871–874. [PubMed: 9579682]
- Qin L, Li G, Qian X, Liu Y, Wu X, Liu B, Hong JS, Block ML. Interactive role of the toll-like receptor 4 and reactive oxygen species in LPS-induced microglia activation. *Glia* 2005;52:78–84. [PubMed: 15920727]
- Richard KL, Filali M, Prefontaine P, Rivest S. Toll-like receptor 2 acts as a natural innate immune receptor to clear amyloid beta 1-42 and delay the cognitive decline in a mouse model of Alzheimer's disease. *J Neurosci* 2008;28:5784–5793. [PubMed: 18509040]
- Ridley AJ, Paterson HF, Johnston CL, Diekmann D, Hall A. The small GTP-binding protein rac regulates growth factor-induced membrane ruffling. *Cell* 1992;70:401–410. [PubMed: 1643658]
- Sanchez-Mejorada G, Rosales C. Signal transduction by immunoglobulin Fc receptors. *J Leukoc Biol* 1998;63:521–533. [PubMed: 9581795]
- Sanlioglu S, Williams CM, Samavati L, Butler NS, Wang G, McCray PB Jr, Ritchie TC, Hunninghake GW, Zandi E, Engelhardt JF. Lipopolysaccharide induces Rac1-dependent reactive oxygen species formation and coordinates tumor necrosis factor-alpha secretion through IKK regulation of NF-kappa B. *J Biol Chem* 2001;276:30188–30198. [PubMed: 11402028]
- Saura J, Tusell JM, Serratos J. High-yield isolation of murine microglia by mild trypsinization. *Glia* 2003;44:183–189. [PubMed: 14603460]
- Savage MJ, Lin YG, Ciallella JR, Flood DG, Scott RW. Activation of c-Jun N-terminal kinase and p38 in an Alzheimer's disease model is associated with amyloid deposition. *J Neurosci* 2002;22:3376–3385. [PubMed: 11978814]
- Scholtzova H, Kascak RJ, Bates KA, Boutajangout A, Kerr DJ, Meeker HC, Mehta PD, Spinner DS, Wisniewski T. Induction of toll-like receptor 9 signaling as a method for ameliorating Alzheimer's disease-related pathology. *J Neurosci* 2009;29:1846–1854. [PubMed: 19211891]
- Selkoe DJ. Toward a comprehensive theory for Alzheimer's disease. Hypothesis: Alzheimer's disease is caused by the cerebral accumulation and cytotoxicity of amyloid beta-protein. *Ann N Y Acad Sci* 2000;924:17–25. [PubMed: 11193794]
- Sendide K, Reiner NE, Lee JS, Bourgoin S, Talal A, Hmama Z. Cross-talk between CD14 and complement receptor 3 promotes phagocytosis of mycobacteria: regulation by phosphatidylinositol 3-kinase and cytohesin-1. *J Immunol* 2005;174:4210–4219. [PubMed: 15778383]
- Sierra A, Gottfried-Blackmore AC, McEwen BS, Bulloch K. Microglia derived from aging mice exhibit an altered inflammatory profile. *Glia* 2007;55:412–424. [PubMed: 17203473]
- Smits HA, de Vos NM, Wat JW, van der Bruggen T, Verhoef J, Nottet HS. Intracellular pathways involved in TNF-alpha and superoxide anion release by Abeta(1-42)-stimulated primary human macrophages. *J Neuroimmunol* 2001;115:144–151. [PubMed: 11282164]
- Stefanova I, Corcoran ML, Horak EM, Wahl LM, Bolen JB, Horak ID. Lipopolysaccharide induces activation of CD14-associated protein tyrosine kinase p53/56lyn. *J Biol Chem* 1993;268:20725–20728. [PubMed: 7691802]
- Streit WJ, Mrak RE, Griffin WS. Microglia and neuroinflammation: a pathological perspective. *J Neuroinflammation* 2004;1:14. [PubMed: 15285801]
- Sumimoto H, Hata K, Mizuki K, Ito T, Kage Y, Sakaki Y, Fukumaki Y, Nakamura M, Takeshige K. Assembly and activation of the phagocyte NADPH oxidase. Specific interaction of the N-terminal Src homology 3 domain of p47phox with p22phox is required for activation of the NADPH oxidase. *J Biol Chem* 1996;271:22152–22158. [PubMed: 8703027]
- Sun A, Liu M, Nguyen XV, Bing G. P38 MAP kinase is activated at early stages in Alzheimer's disease brain. *Exp Neurol* 2003;183:394–405. [PubMed: 14552880]
- Taborda CP, Casadevall A. CR3 (CD11b/CD18) and CR4 (CD11c/CD18) are involved in complement-independent antibody-mediated phagocytosis of *Cryptococcus neoformans*. *Immunity* 2002;16:791–802. [PubMed: 12121661]
- Tahara K, Kim HD, Jin JJ, Maxwell JA, Li L, Fukuchi K. Role of toll-like receptor signalling in Abeta uptake and clearance. *Brain* 2006;129:3006–3019. [PubMed: 16984903]


- Tang G, Yue Z, Talloczy Z, Hagemann T, Cho W, Messing A, Sulzer DL, Goldman JE. Autophagy induced by Alexander disease-mutant GFAP accumulation is regulated by p38/MAPK and mTOR signaling pathways. *Hum Mol Genet* 2008;17:1540–1555. [PubMed: 18276609]
- Traenckner EB, Pahl HL, Henkel T, Schmidt KN, Wilk S, Baeuerle PA. Phosphorylation of human I kappa B-alpha on serines 32 and 36 controls I kappa B-alpha proteolysis and NF-kappa B activation in response to diverse stimuli. *EMBO J* 1995;14:2876–2883. [PubMed: 7796813]
- Triantafilou M, Brandenburg K, Kusumoto S, Fukase K, Mackie A, Seydel U, Triantafilou K. Combinational clustering of receptors following stimulation by bacterial products determines lipopolysaccharide responses. *Biochem J* 2004;381:527–536. [PubMed: 15040785]
- Triantafilou M, Gamper FG, Haston RM, Mouratis MA, Morath S, Hartung T, Triantafilou K. Membrane sorting of toll-like receptor (TLR)-2/6 and TLR2/1 heterodimers at the cell surface determines heterotypic associations with CD36 and intracellular targeting. *J Biol Chem* 2006;281:31002–31011. [PubMed: 16880211]
- Udan ML, Ajit D, Crouse NR, Nichols MR. Toll-like receptors 2 and 4 mediate Abeta(1-42) activation of the innate immune response in a human monocytic cell line. *J Neurochem* 2008;104:524–533. [PubMed: 17986235]
- Underhill DM, Ozinsky A, Hajjar AM, Stevens A, Wilson CB, Bassetti M, Aderem A. The Toll-like receptor 2 is recruited to macrophage phagosomes and discriminates between pathogens. *Nature* 1999;401:811–815. [PubMed: 10548109]
- Van Muiswinkel FL, Raupp SF, de Vos NM, Smits HA, Verhoef J, Eikelenboom P, Nottet HS. The amino-terminus of the amyloid-beta protein is critical for the cellular binding and consequent activation of the respiratory burst of human macrophages. *J Neuroimmunol* 1999;96:121–130. [PubMed: 10227431]
- van Oosten M, van Amersfoort ES, van Berkel TJ, Kuiper J. Scavenger receptor-like receptors for the binding of lipopolysaccharide and lipoteichoic acid to liver endothelial and Kupffer cells. *J Endotoxin Res* 2001;7:381–384. [PubMed: 11753207]
- Walter S, Letiembre M, Liu Y, Heine H, Penke B, Hao W, Bode B, Manietta N, Walter J, Schulz-Schuffer W, Fassbender K. Role of the toll-like receptor 4 in neuroinflammation in Alzheimer's disease. *Cell Physiol Biochem* 2007;20:947–956. [PubMed: 17982277]
- Wilkinson B, Koenigsnecht-Talboo J, Grommes C, Lee CY, Landreth G. Fibrillar beta-amyloid-stimulated intracellular signaling cascades require Vav for induction of respiratory burst and phagocytosis in monocytes and microglia. *J Biol Chem* 2006;281:20842–20850. [PubMed: 16728400]
- Wong WS, Simon DI, Rosoff PM, Rao NK, Chapman HA. Mechanisms of pertussis toxin-induced myelomonocytic cell adhesion: role of Mac-1(CD11b/CD18) and urokinase receptor (CD87). *Immunology* 1996;88:90–97. [PubMed: 8707356]
- Yamamori T, Inanami O, Nagahata H, Cui Y, Kuwabara M. Roles of p38 MAPK, PKC and PI3-K in the signaling pathways of NADPH oxidase activation and phagocytosis in bovine polymorphonuclear leukocytes. *FEBS Lett* 2000;467:253–258. [PubMed: 10675549]
- Yan SR, Al-Hertani W, Byers D, Bortolussi R. Lipopolysaccharide-binding protein- and CD14-dependent activation of mitogen-activated protein kinase p38 by lipopolysaccharide in human neutrophils is associated with priming of respiratory burst. *Infect Immun* 2002;70:4068–4074. [PubMed: 12117913]
- Zhu X, Rottkamp CA, Boux H, Takeda A, Perry G, Smith MA. Activation of p38 kinase links tau phosphorylation, oxidative stress, and cell cycle-related events in Alzheimer disease. *J Neuropathol Exp Neurol* 2000;59:880–888. [PubMed: 11079778]
- Zuluaga S, Gutierrez-Uzquiza A, Bragado P, Alvarez-Barrientos A, Benito M, Nebreda AR, Porras A. p38alpha MAPK can positively or negatively regulate Rac-1 activity depending on the presence of serum. *FEBS Lett* 2007;581:3819–3825. [PubMed: 17658519]



**Figure 1. CD14, TLR4, and TLR2 mediate the recognition and binding of fAβ**

THP-1 monocytes were incubated in the presence or absence of function blocking antibodies to CD14 (MY4, **A**), TLR4 (HTA125, **A**), TLR2 (T2.5, **A**), CD36 (FA6-152, **B**), CD47 (B6H12, **B**), or their isotype controls (IgG<sub>2b</sub>, IgG<sub>2a</sub>, IgG<sub>1</sub>). Cells were then added to fAβ bound to a glass slide and the number of adherent cells was determined. **C**, THP-1 monocytes were incubated in the presence or absence of 0.1 μg/ml polymixin B. Cells were then added to fAβ bound to a glass slide and the number of adherent cells was determined. The data shown are the mean ± SEM of three independent experiments. (\*p<0.05, \*\*p<0.01, \*\*\*p<0.001)






**Figure 2. Phagocytosis stimulated by fAβ requires CD14, TLR4, and TLR2**


**A**, BV2 microglia were pretreated with function blocking antibodies to CD14 (MY4), TLR4 (HTA125), or TLR2 (T2.5), prior to stimulation with fAβ. Fluorescent microspheres were then added for 30 minutes. Cells were stained with phalloidin for visualization of actin and DAPI. Magnification is 40×. **B**, Confocal images (100×) of BV2 microglia treated with fAβ containing microspheres in both the x-y plane (left) and z-scan cut (right) through the cells. **C**, Primary microglia from C57BL/6 (WT), CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, or TLR2<sup>-/-</sup> mice were stimulated with fAβ, followed by incubation with fluorescent microspheres. The fraction of cells containing microspheres was determined by counting three random fields (>100 cells) on an inverted

microscope. The data shown are the mean  $\pm$  SEM of three independent experiments performed in duplicate. (\*\*\*) $p < 0.001$ )




**Figure 3. CD14, TLR4, and TLR2 are necessary for fAβ-stimulated reactive oxygen species production**

Primary microglia from C57BL/6 (WT), CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, TLR2<sup>-/-</sup> mice were stimulated with fAβ (A), LPS (B, C), Pam3CSK4 (D), or PMA (B-D). Reduction of NBT results in the formation of a dark blue formazan deposit viewable by a light microscope. Three random fields (>100 cells) were counted and the fraction of cells with blue formazan deposits was determined. The data shown are the mean ± SEM of three independent experiments performed in duplicate. (\*p<0.05, \*\*p<0.01, \*\*\*p<0.001)




**Figure 4. Activation of the Src tyrosine kinases by fAβ requires CD14, TLR4, and TLR2**  
 Primary microglia from C57BL/6 (WT), CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, or TLR2<sup>-/-</sup> mice were stimulated with fAβ (A), LPS (B, C), Pam3CSK4 (D), or immune IgG (B-D). Cell lysates were analyzed by Western blot analysis using an anti-phospho-Src family antibody. Blots were stripped and reprobed with an anti-cSrc antibody as a protein loading control. A, Western blot band intensity of phosphorylated Src kinases was normalized to cSrc levels and expressed as relative density. Western blots are representative, and the densitometry data is the mean ± SEM from at least three independent experiments. (\*p<0.05)




**Figure 5. CD14, TLR4, and TLR2 are required for fA $\beta$ -induced phosphorylation of the Rac GEF Vav and subsequent translocation of Rac to the membrane**


**A.** Primary microglia from C57BL/6 (WT), CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, or TLR2<sup>-/-</sup> mice were stimulated with fA $\beta$ . Vav was immunoprecipitated from cell lysates with an anti-Vav antibody and analyzed by Western blot analysis using the anti-phosphoTyr antibody 4G10. Blots were stripped and reprobed with an anti-Vav antibody as a protein loading control. **B.** Primary microglia from C57BL/6 (WT), CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, or TLR2<sup>-/-</sup> mice were stimulated with fA $\beta$ . Membrane fractions were analyzed by Western blot analysis with an anti-Rac antibody and an anti-flotillin antibody as a protein loading control. Western blots are representative, and the densitometry data is the mean  $\pm$  SEM from three independent experiments. (\*\*p<0.01)



**Figure 6. Activation of the p38 MAP kinase by fAβ requires CD14, TLR4, and TLR2**  
 Primary microglia from C57BL/6 (WT), CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, or TLR2<sup>-/-</sup> mice were stimulated with fAβ (A), LPS (B, C), Pam3CSK4 (D), or immune IgG (B-D). Cell lysates were analyzed by Western blot analysis using an anti-phospho-p38 antibody. Blots were stripped and reprobed with an anti-p38 antibody as a protein loading control. Western blot band intensity of phosphorylated-p38 was normalized to p38 levels and expressed as relative density. Western blots are representative, and the densitometry data is the mean ± SEM from at least three independent experiments. (\*\*p < 0.01, \*\*\*p < 0.001)



**Figure 7. Fibrillar A $\beta$ -stimulated ROS production and phagocytosis requires the activity of p38**  
**A**, Primary microglia from C57BL/6 mice were pretreated with SB203580 prior to stimulation with fA $\beta$ . Reduction of NBT results in the formation of a dark blue formazan deposit viewable by a light microscope. Three random fields (>100 cells) were counted and the fraction of cells with blue formazan deposits was determined. The data shown are the mean  $\pm$  SEM of three independent experiments performed in duplicate. (\*\*\*) $p < 0.001$   
**B**, BV2 microglia were pretreated with SB203580 prior to stimulation with fA $\beta$ . Fluorescent microspheres were added, following which the fraction of cells containing microspheres was determined by counting three random fields (>100 cells) on an inverted microscope. The data shown are the mean  $\pm$  SEM of three independent experiments performed in duplicate and normalized to actin levels. (\*\*\*) $p < 0.001$



**Figure 8. CD14, TLR4, and TLR2 are required for fAβ-induced NFκB activation**  
Primary microglia from C57BL/6 (WT), CD14<sup>-/-</sup>, TLR4<sup>-/-</sup>, or TLR2<sup>-/-</sup> mice were stimulated with fAβ. Cell lysates were analyzed by Western blot analysis using an anti-phospho-IκBα antibody. Blots were stripped and reprobbed with an anti-actin antibody as a protein loading control. Western blot band intensity of phosphorylated-IκBα was normalized to actin levels and expressed as relative density. Western blots are representative, and the densitometry data represent the mean ± SEM from at least three independent experiments. (\*p<0.05)