



Published in final edited form as:

*AIDS*. 2020 October 01; 34(12): 1789–1794. doi:10.1097/QAD.0000000000002632.

## Clinical Characteristics, Comorbidities and Outcomes Among Persons with HIV Hospitalized with Coronavirus Disease 2019 in Atlanta, GA

Lauren F. COLLINS, MD<sup>1,2</sup>, Caitlin A. MORAN, MD<sup>1,2</sup>, Nora T. OLIVER, MD<sup>1,3</sup>, Abeer MOANNA, MD<sup>1,3</sup>, Cecile D. LAHIRI, MD<sup>1,2</sup>, Jonathan A. COLASANTI, MD<sup>1,2</sup>, Colleen F. KELLEY, MD<sup>1,2,4</sup>, Minh Ly NGUYEN, MD<sup>1,2</sup>, Vincent C. MARCONI, MD<sup>1,2,3</sup>, Wendy S. ARMSTRONG, MD<sup>1,2</sup>, Ighowwerha OFOTOKUN, MD<sup>1,2</sup>, Anandi N. SHETH, MD<sup>1,2</sup>

<sup>1</sup>Division of Infectious Diseases, Emory University School of Medicine, Atlanta, GA, USA;

<sup>2</sup>Grady Healthcare System, Infectious Diseases Program, Atlanta, GA, USA;

<sup>3</sup>Atlanta Veterans Affairs Medical Center, Decatur, GA, USA;

<sup>4</sup>The Hope Clinic of the Emory Vaccine Center, Decatur, GA, USA

### Abstract

**Background:** There are limited data describing the presenting characteristics and outcomes among U.S. persons with HIV (HIV) requiring hospitalization for coronavirus disease 2019 (COVID-19).

**Methods:** We performed a case series of all PWH sequentially admitted with COVID-19 from 3/8/2020 to 4/23/2020 at three hospitals in Atlanta, Georgia. Sociodemographic, clinical and HIV-associated characteristics were collected.

**Results:** Of 530 confirmed COVID-19 cases hospitalized during this period, 20 occurred among PWH (3.8%). The median age was 57 (Q1-Q3, 48–62) years, 65% were men, and 85% were non-Hispanic Black. Presenting median symptom duration was 5 (Q1-Q3, 3–7) days; cough (90%), fever (65%), malaise (60%) and dyspnea (60%) were most common. On admission, 40% of patients required oxygenation support and 65% had an abnormal chest radiograph. Median length of hospitalization was 5 (Q1-Q3, 4–12) days, 30% required intensive care, 15% required intubation, and 15% died. Median CD4 count prior to admission was 425 (Q1-Q3, 262–815) cells/mm<sup>3</sup> and 90% of patients had HIV-1 RNA <200 copies/ml. Half of the patients had 5 comorbidities; hypertension (70%), dyslipidemia (60%) and diabetes (45%) were most prevalent. All three patients who died had CD4 count >200, HIV suppression and each had a total of 5 comorbidities.

**Conclusions:** This multisite series in the Southern U.S. provides characteristics and early outcomes of hospitalized PWH with COVID-19. Nearly all patients had controlled HIV and a high comorbidity burden. Additional study of COVID-19 among PWH is needed to determine the role of age, comorbidities, and HIV control in mediating COVID-19 presentation and its sequelae.

## Keywords

Human immunodeficiency virus; persons living with HIV; severe acute respiratory syndrome coronavirus 2; coronavirus disease 2019; comorbidity burden

---

## Introduction

The severe acute respiratory syndrome-coronavirus-2 (SARS-CoV-2) pandemic continues to spread unabated globally, with >1 million confirmed cases reported in the U.S. Coronavirus disease-2019 (COVID-19), a novel pneumonia caused by SARS-CoV-2, particularly devastates those of older age and with multimorbidity. The Centers for Disease Control and Prevention categorizes people with “poorly controlled HIV or AIDS” at higher risk than the general population for severe illness from COVID-19<sup>1</sup>, however, supportive data are lacking. The Southeast persists as the epicenter of the U.S. HIV epidemic with the majority of new diagnoses, lower rates of virologic suppression and higher rates of deaths from late-stage AIDS than other parts of the country<sup>2</sup>. To understand how COVID-19 may affect persons with HIV (PWH) in the Southern U.S., a prematurely aging population with a high comorbidity burden<sup>3,4</sup>, we analyzed cases among hospitalized PWH in Atlanta, Georgia.

## Methods

PWH admitted with COVID-19 at one of three hospitals in Atlanta, Georgia between 3/8/2020 and 4/23/2020 were recorded. The included hospitals were: (1) Grady Memorial Hospital, a 961-bed public safety-net hospital, (2) Emory University Hospital, a 587-bed tertiary academic medical center, and (3) the Atlanta Veterans Affairs Medical Center, a 466-bed hospital serving veterans in the metropolitan and surrounding areas. COVID-19 was confirmed by detection of SARS-CoV-2 by reverse transcription-polymerase chain reaction (RT-PCR) from a respiratory specimen.

Sociodemographic, clinical and HIV-associated characteristics were manually extracted from the electronic medical record. Non-AIDS comorbidities were included if documented by a healthcare provider in the outpatient setting or during hospitalization. Viral suppression was defined as HIV-1 RNA <200 copies/ml. The research was approved by the Emory University Institutional Review Board.

## Results

Of 530 confirmed COVID-19 cases admitted during this period, 20 occurred among PWH (3.8%). The median age was 57 (Q1-Q3, 48–62) years, 65% were men, and 85% were non-Hispanic Black; 60% were admitted to the public safety-net hospital. While 50% denied recent travel or a known exposure, seven (35%) reported known sick contacts. Three of these were nursing home residents, two had ill household contacts, one was recently incarcerated, and one interacted with clients from Italy at the workplace in Atlanta (Table). Median symptom duration on admission was 5 (Q1-Q3, 3–7) days; cough (90%), fever (65%), malaise (60%) and dyspnea (60%) were most common. Two patients experienced both anosmia and ageusia.

Overall, presenting vital sign abnormalities were subtle (Table). Forty percent of patients required oxygenation support in the emergency department including two patients who were intubated on arrival. Absolute lymphocyte count was depressed, and serum creatinine, lactate dehydrogenase, C-reactive protein and D-dimer were all elevated (Table). Chest radiograph was abnormal in 65% of patients, with 40% showing diffuse multifocal or bibasilar opacities and 15% demonstrating a focal infiltrate. Over half (55%) of patients received a therapeutic for SARS-CoV-2 including open-label hydroxychloroquine +/- azithromycin or clinical trial enrollment. Median length of hospitalization was 5 (Q1-Q3, 4–12) days, 30% required intensive care, 15% required intubation, and 15% died. One patient remained admitted to date requiring high-flow oxygen support via nasal cannula on hospital day six.

All patients had SARS-CoV-2 detected by RT-PCR on initial specimen testing (19/20 specimens were collected by nasopharyngeal swab and 1/20 by mini-bronchoalveolar lavage). Three patients had repeat testing performed for discharge planning. Each individual had two nasopharyngeal swabs obtained several days after their initial test, including one individual who had a negative test result on both repeat tests (obtained on hospital days 18 and 20) and two individuals who had positive test results on both repeat tests (obtained on hospital days 7 and 10, and on hospital days 12 and 14, respectively).

Fifty-five percent of patients had HIV diagnosed 10 years ago. The most frequently reported HIV acquisition risk factor was heterosexual behavior (55%), followed by men who have sex with men activity (25%) and injection drug use (10%). Median absolute CD4 count and percentage prior to admission were 425 (Q1-Q3, 262–815) cells/mm<sup>3</sup> and 29% (Q1-Q3, 21–36), respectively, and 90% of patients had viral suppression (Table). In the two years prior to hospitalization, 82% of patients were continuously virologically suppressed over a median of 4.5 (Q1-Q3, 3.3–5.0) HIV care visits per patient. All 20 patients were prescribed antiretroviral therapy prior to hospitalization, and 95% reported therapy adherence on admission. Eighty percent were prescribed an integrase strand transfer inhibitor-containing regimen, most commonly bictegravir/emtricitabine/tenofovir alafenamide (35%) or dolutegravir/abacavir/lamivudine (25%).

The burden of non-AIDS comorbidities was high in this cohort, with 50% of patients having 5 comorbidities, and only 15% with no comorbidities (Table). The most prevalent comorbidities were hypertension (70%), dyslipidemia (60%), diabetes (45%) and psychiatric illness (40%). Nine patients were prescribed an angiotensin-converting enzyme inhibitor or angiotensin II receptor blocker on their medication list prior to admission. Forty percent of patients reported current/former cigarette smoking, 50% currently used alcohol, and 20% reported a history of crack/cocaine use. One patient had chronic active hepatitis B virus and five patients had a history of hepatitis C virus, including two who spontaneously cleared and three who achieved sustained virologic response after treatment with direct-acting antivirals.

A summary of demographic, comorbidity, HIV-specific and treatment characteristics for the three patients whom died is provided in the Table 2.

## Discussion

This multisite series of PWH hospitalized with COVID-19 in Atlanta is notable for several reasons. The median age of 57 years is 4–6 years younger than reports of non-HIV hospitalized COVID-19 U.S. cohorts<sup>5,6</sup>. Half of PWH in this study had 5 comorbidities, an alarming burden that exceeds previous reports of hospitalized COVID-19 cases including adults in New York and California<sup>5,6</sup> and specifically among PWH in Spain<sup>7</sup>. Among hospitalized PWH in this series, nearly all had CD4 counts  $\geq 200$  and viral suppression (inclusive of all three patients who died, Table 2), despite the three hospitals' catchment areas treating significant proportions of demographically-similar PWH with uncontrolled HIV and AIDS<sup>8</sup>. It is possible that the premature onset of multimorbidity among PWH considerably impacts SARS-CoV-2 infection and illness severity, distinct from HIV infection or its associated immunosuppression.

The immunologic intersection between SARS-CoV-2 and chronic HIV infection remains to be examined. Clearly, in a subset of people with COVID-19, a hyper-immune response to SARS-CoV-2 leads to severe lung damage and poor outcomes exemplified by elevated systemic levels of pro-inflammatory cytokines, coagulation dysfunction, and dysregulated cellular responses<sup>9</sup>. Curiously, we and others<sup>5</sup> have not yet observed a high prevalence of severe COVID-19 disease among PWH in general, and particularly not among PWH with advanced immunosuppression, i.e., CD4  $< 200$ . Further research will be necessary to understand whether the immune response to SARS-CoV-2 differs among PWH and how multimorbidity, which is common in HIV<sup>3,4</sup>, influences clinical outcomes of COVID-19 among PWH specifically.

PWH in the Southern U.S. represent a group particularly vulnerable to COVID-19 severity and sequelae, given the confluence of income inequality, housing instability and food insecurity – disproportionate among minorities because of structural racism and discrimination – limiting the ability to adopt optimal infection prevention measures and augmenting comorbidity risk, which in turn confers worse outcomes<sup>2</sup>. The predominance of non-Hispanic Black men in this cohort reflects the demographic of HIV/AIDS in Atlanta; however, more data are needed to determine the role of racial/ethnic disparities in COVID-19 presentation and outcomes, especially as this can be exacerbated in high HIV-burden areas<sup>10</sup>.

The indirect effects of the SARS-CoV-2 pandemic on PWH and those at risk for HIV are not yet fully understood. Closure of clinics, reduced access to medications and fewer support services may lead to decreased rates of retention in care and lower rates of virologic suppression, particularly among the most marginalized<sup>11</sup>. In addition, many clinics have scaled back testing for HIV and other sexually transmitted infections, and others have suspended visits for pre-exposure prophylaxis (PrEP). The overall impact on the HIV care continuum and the *Ending the HIV Epidemic* initiative must be closely monitored.

It is anticipated that additional resources are needed to support the unique needs of PWH, including to increase access to SARS-CoV-2 testing and clinical trials, and more broadly to build socioeconomic and healthcare infrastructure that will enhance HIV-related,

comorbidity, mental and sexual health and well-being<sup>12</sup>. This report supports dedicated larger study of COVID-19 among PWH to determine the role of age, comorbidities, immunocompetency, HIV control and antiretroviral agents in mediating COVID-19 presentation and its sequelae.

## Acknowledgements:

We sincerely thank the patients, clinicians and staff at Grady Memorial Hospital, Emory University Hospital and the Atlanta Veteran's Affairs Medical Center. We also thank the Emory-Grady HIV Clinical Cohort Registry for contribution of data.

**Financial support:** This work was supported by the Emory Center for AIDS Research (award number P30-AI-050409) as well as the Emory Specialized Center of Research Excellence (SCORE) on Sex Differences (award number U54AG062334; to I. O.). L. F. C. is also supported by the National Center for Advancing Translational Sciences (NCATS) of the National Institutes of Health (NIH) (award numbers UL1TR002378 and TL1TR002382) and C.A.M. by the NCATS of the NIH (award number KL2TR002381). C.D.L. is also supported by the National Institute of Allergy and Infectious Diseases (NIAID) of the NIH (award number K23-AI124913).

## References

- Centers for Disease Control and Prevention. Coronavirus Disease 2019 (COVID-19): People Who Are At Higher Risk. <https://www.cdc.gov/coronavirus/2019-ncov/need-extra-precautions/people-at-higher-risk.html>. Accessed April 26, 2020.
- Colasanti JA & Armstrong WS Challenges of reaching 90-90-90 in the Southern United States. *Curr Opin HIV AIDS* 14, 471–480 (2019). [PubMed: 31425179]
- Lerner AM, Eisinger RW & Fauci AS Comorbidities in Persons With HIV: The Lingering Challenge. *JAMA* (2019) doi:10.1001/jama.2019.19775.
- Collins LF et al. The Prevalence and Burden of Non-AIDS Comorbidities among Women living with or at-risk for HIV Infection in the United States. *Clin. Infect. Dis* (2020) doi:10.1093/cid/ciaa204.
- Richardson S et al. Presenting Characteristics, Comorbidities, and Outcomes Among 5700 Patients Hospitalized With COVID-19 in the New York City Area. *JAMA* (2020) doi:10.1001/jama.2020.6775.
- Myers LC, Parodi SM, Escobar GJ & Liu VX Characteristics of Hospitalized Adults With COVID-19 in an Integrated Health Care System in California. *JAMA* (2020) doi:10.1001/jama.2020.7202.
- Blanco JL et al. COVID-19 in patients with HIV: clinical case series. *Lancet HIV* (2020) doi:10.1016/S2352-3018(20)30111-9.
- Centers for Disease Control and Prevention. Estimated HIV incidence and prevalence in the United States, 2010–2016. HIV Surveillance Supplemental Report 2019;24(No. 1). <http://www.cdc.gov/hiv/library/reports/hiv-surveillance.html>. Published February 2019. Accessed [3/21/2020].
- Chen G et al. Clinical and immunological features of severe and moderate coronavirus disease 2019. *J. Clin. Invest* (2020) doi:10.1172/JCI137244.
- The Centers for Disease Prevention and Control. Characteristics and Clinical Outcomes of Adult Patients Hospitalized with COVID-19 — Georgia, March 2020. *Morbidity and Mortality Weekly Report* 69, (2020).
- Colasanti J et al. Continuous Retention and Viral Suppression Provide Further Insights Into the HIV Care Continuum Compared to the Cross-sectional HIV Care Cascade. *Clin. Infect. Dis* 62, 648–654 (2016). [PubMed: 26567263]
- Shoptaw S, Goodman-Meza D & Landovitz RJ Collective Call to Action for HIV/AIDS Community-Based Collaborative Science in the Era of COVID-19. *AIDS Behav* (2020) doi:10.1007/s10461-020-02860-y.

**Table.**

Clinical Characteristics, Comorbidities and Outcomes Among Persons Living with HIV Hospitalized with Coronavirus Disease 2019 in Atlanta, Georgia

<i>Characteristic, median (Q1-Q3), mean (sd) or n (%)</i>	<b>Data (n=20<sup>a</sup>)</b>
<b>Demographics</b>	
Age, years	57 (48–62)
Sex	
Male	13/20 (65)
Female	6/20 (30)
Male-to-female transgender	1/20 (5)
Race/ethnicity	
Non-Hispanic Black	17/20 (85)
Non-Hispanic White	1/20 (5)
Non-Hispanic/Multiracial	1/20 (5)
Hispanic/Latino	1/20 (5)
Body mass index, kg/m <sup>2</sup>	28 (24–31)
Medical Facility	
Public Safety-net Hospital	12/20 (60)
Tertiary Academic Medical Center	4/20 (20)
Veterans Affairs Medical Center	4/20 (20)
<b>Exposure and Symptom Inventory on Admission</b>	
Potential Exposure	
No recent travel/known sick contact	10/20 (50)
Known sick contact(s) in the home	2/20 (10)
Recent domestic travel	1/20 (5)
Group living situation <sup>b</sup>	5/20 (25)
Ongoing work outside the home <sup>c</sup>	2/20 (10)
Duration of symptoms, days	5 (3–7)
Symptoms reported	
Fever	13/20 (65)
Chills	10/20 (50)
Malaise	12/20 (60)
Cough	18/20 (90)
Shortness of breath	12/20 (60)
Chest tightness or pleuritic chest pain	4/20 (20)
Diarrhea	6/20 (30)
Nausea or vomiting	6/20 (30)
Myalgias	8/20 (40)
Headache	4/20 (20)
Sore throat	3/20 (15)
Anosmia	2/20 (10)

<i>Characteristic, median (Q1-Q3), mean (sd) or n (%)</i>	<b>Data (n=20<sup>a</sup>)</b>
Ageusia	2/20 (10)
<b>Vital Signs on Admission</b>	
Temperature, °C	37.9 (37.2–38.7)
Respiratory rate, breaths per minutes	20 (18–21)
Oxygen saturation on ambient air, %	96 (84–97)
Oxygen support required	
None	12/20 (60)
Nasal cannula	6/20 (30)
Invasive ventilation	2/20 (10)
<b>Laboratory Findings on Admission</b>	
White blood cells, K/mcL	6 (5.3–7.1)
Absolute lymphocyte count, K/mcL	0.98 (0.7–1.4)
Creatinine, mg/dL	1.4 (1.1–1.6)
Lactate dehydrogenase, U/L <sup>d</sup>	208 (173–353)
C-reactive protein, mg/L <sup>d</sup>	73 (31–163)
Ferritin, ng/mL <sup>d</sup>	357 (172–530)
D-dimer, ng/mL <sup>d</sup>	502 (323–1444)
<b>Chest Imaging</b>	
Chest radiograph findings on admission	
No acute process	7/20 (35)
Bibasilar opacities	5/20 (25)
Diffuse airspace opacities	3/20 (15)
Focal infiltrate	5/20 (15)
Computed tomography of the chest	
Not performed during hospitalization	14/20 (70)
Findings of bilateral diffuse GGO	5/20 (25)
Findings of localized GGO and focal infiltrate	1/20 (5)
<b>Treatment</b>	
Therapy for SARS-CoV-2	
Supportive care only	9/20 (45)
Hydroxychloroquine	8/20 (40)
Hydroxychloroquine/azithromycin	2/20 (10)
Enrolled in remdesivir versus placebo trial	1/20 (5)
Number of days therapy given for SARS-CoV-2	5 (5–5)
Antibiotics for secondary pneumonia	
None	7/20 (35)
Community-acquired coverage	9/20 (45)
Healthcare-associated coverage	4/20 (20)
Number of total antibiotic days for secondary pneumonia	5 (3–7)

<i>Characteristic, median (Q1-Q3), mean (sd) or n (%)</i>	<b>Data (n=20<sup>a</sup>)</b>
<b>Clinical course</b>	
Length of hospitalization, days <sup>e</sup>	5 (4–12)
Required intensive care	6/20 (30)
Number of intensive care days	7 (3–13)
Highest level of respiratory support needed	
None	9/20 (45)
Nasal cannula	5/20 (25)
Non-invasive ventilation	3/20 (15)
Invasive ventilation	3/20 (15)
Pathogen-confirmed secondary infection <sup>f</sup>	5/20 (25)
Final disposition	
Still hospitalized	1/20 (5)
Home	13/20 (65)
Nursing facility as permanent residence	1/20 (5)
Hotel for those with confirmed COVID-19	2/20 (10)
Deceased	3/20 (15)
<b>HIV-specific Indices</b>	
CD4 count prior to admission, cells/mm <sup>3</sup>	425 (262–815)
CD4 percentage prior to admission, %	29 (21–36)
Most recent HIV viral load	
<200 copies/ml	18/20 (90)
200–1000 copies/ml	1/20 (5)
>1000 copies/ml	1/20 (5)
Number of HIV clinic visits attended in past two years	4.5 (3.25–5)
Proportion of visits with HIV virologic suppression in past two years, %	82.2 (27.2)
Prescribed antiretroviral therapy	20/20 (100)
Reported adherence to antiretroviral therapy	19/20 (95)
Antiretroviral therapy class prescribed <sup>g</sup>	
Non-nucleoside reverse transcriptase inhibitor	2/20 (10)
Protease inhibitor, boosted	4/20 (20)
Integrase strand transfer inhibitor	16/20 (80)
<b>Non-AIDS Comorbidities</b>	
Hypertension	14/20 (70)
Dyslipidemia	12/20 (60)
Diabetes mellitus, type 2	9/20 (45)
Cardiovascular disease	6/20 (30)
Chronic lung disease <sup>h</sup>	6/20 (30)
Obesity	6/20 (30)



Characteristic, median (Q1-Q3), mean (sd) or n (%)	Data (n=20 <sup>a</sup> )
Chronic kidney disease	5/20 (25)
Non-AIDS Cancer <sup>i</sup>	3/20 (15)
Psychiatric illness (depression, anxiety)	8/20 (40)
<b>Non-AIDS Comorbidity Burden</b>	
Number of prevalent comorbidities per patient <sup>j</sup>	
0	3/20 (15)
1-2	3/20 (15)
3-4	4/20 (20)
5	10/20 (50)
<b>Substance use</b>	
Cigarette smoking	
Current	3/20 (15)
Former	5/20 (25)
Never	12/20 (60)
Alcohol consumption	
Current	10/20 (50)
Former	2/20 (10)
Never	8/20 (40)
History of illicit drug use <sup>g</sup>	
Marijuana	3/20 (15)
Crack/cocaine	4/20 (20)
Methamphetamines	2/20 (10)
Opioid use disorder on methadone	1/20 (5)

**Abbreviations:** GGO = ground glass opacities; PCR = polymerase chain reaction; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2

<sup>a</sup>Unless otherwise noted.

<sup>b</sup>Nursing home (n=3), incarceration (n=1), substance abuse recovery home (n=1).

<sup>c</sup>Includes a patient who interacted with travelers from Italy at work, of whom there were 3 confirmed cases of coronavirus disease 2019.

<sup>d</sup>Data available for the following number of individuals: lactate dehydrogenase (n=18), C-reactive protein (n=16), ferritin (n=13), D-dimer (n=16), international normalized ratio (n=11).

<sup>e</sup>Includes one individual who remains hospitalized at day 6.

<sup>f</sup>Includes one individual with *Streptococcus pneumoniae* bacteremia; one with *Streptococcus pyogenes* pneumonia; one with methicillin-resistant *Staphylococcus aureus* bacteremia as well as polymicrobial ventilator-associated pneumonia (*Acinetobacter baumannii*, *Klebsiella aerogenes*); and two individuals with *Clostridioides difficile* colitis.

<sup>g</sup>Not mutually exclusive.

<sup>h</sup>Includes chronic obstructive pulmonary disease (n=3), asthma (n=1), obstructive sleep apnea (n=1), pulmonary fibrosis (n=1).

<sup>i</sup>Includes one patient with renal cell carcinoma status post nephrectomy and subsequent deceased-donor kidney transplantation currently on immunosuppression; one patient with prostate cancer who has repeatedly declined treatment; one patient with Hodgkin's lymphoma treated with chemotherapy (n=1).

<sup>j</sup>Out of the 9 listed above.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

**Table 2.**

Characteristics of Persons with HIV Who Died of Coronavirus Disease 2019 in Atlanta, Georgia

<i>Characteristic</i>	<b>Patient 1</b>	<b>Patient 2</b>	<b>Patient 3</b>
Age, years	54	61	66
Sex	Male	Male	Male
Race/ethnicity	NHB	NHB	NHB
Known SARS-CoV-2 exposure	Nursing home	No	No
Intubation required	Yes	Yes	Do not resuscitate or intubate order in place
C-reactive protein on admission, mg/L	158.5	448.3	67.8
D-dimer on admission, ng/mL	1574	3251	501
Length of hospitalization	14 days	18 days	5 days
Treatment for SARS-CoV-2	Supportive care	HCQ Azithromycin	HCQ
Number of prevalent non-AIDS comorbidities	5	5	5
List of prevalent non-AIDS comorbidities	Hypertension Dyslipidemia Diabetes CVD CKD	Hypertension Dyslipidemia Diabetes CVD Prostate cancer	Hypertension Dyslipidemia Diabetes COPD CKD
CD4 count (cells/mm <sup>3</sup> )/percentage prior to admission	238/19%	391/40%	974/31%
HIV-1 RNA prior to or during hospitalization, copies/ml	Not detected	Not detected	22 copies/ml (1.34 log <sub>10</sub> )
Antiretroviral therapy prescribed prior to admission	3TC, ETR DRV/r	TAF/FTC DRV/c	ABC/3TC DTG

**Abbreviations:** 3TC = lamivudine; ABC = abacavir; CKD = chronic kidney disease; COPD = chronic obstructive pulmonary disease; CVD = cardiovascular disease; DRV/r or DRV/c = darunavir boosted with ritonavir (r) or cobicistat (c); DTG = dolutegravir; ETR = etravirine; FTC = emtricitabine; HCQ = hydroxychloroquine; NHB = Non-Hispanic Black; SARS-CoV-2 = severe acute respiratory syndrome coronavirus 2; TAF = tenofovir alafenamide fumarate