

Published in final edited form as:

Toxins (Basel). 2009 December 1; 1(2): 123–133. doi:10.3390/toxins1020123.

Actin Crosslinking Toxins of Gram-Negative Bacteria

Karla J. F. Satchell

Northwestern University, Feinberg School of Medicine, 320 E. Superior Avenue, Chicago, IL 60611, USA; k-satchell@northwestern.edu; Tel.: +1-312-503-2162; Fax: +1-312-503-1339

Abstract

Actin crosslinking toxins produced by Gram-negative bacteria represent a small but unique class of bacterial protein toxins. For each of these toxins, a discrete actin crosslinking domain (ACD) that is a distant member of the ATP-dependent glutamine synthetase family of protein ligases is translocated to the eukaryotic cell cytosol. This domain then incorporates a glutamate-lysine crosslink between actin monomers, resulting in destruction of the actin cytoskeleton. Recent studies argue that the function of these toxins during infection is not destruction of epithelial layers, but rather may specifically target phagocytic cells to promote survival of bacteria after the onset of innate immune defenses. This review will summarize key experiments performed over the past 10 years to reveal the function of these toxins.

Keywords

bacterial protein toxin; *Vibrio*; MARTX; VgrG; actin; ACD

1. Bacterial Protein Toxins with an Actin Crosslinking Domain

1.1. Overview of actin targeting toxins

Many toxins have been characterized that disrupt the eukaryotic actin cytoskeleton. These include fungal and marine small molecule toxins that bind to actin to prevent actin polymerization [1,2]. Other bacterial protein toxins and effectors can alter the normal regulation of cytoskeletal assembly, by directly or indirectly modifying the activity of small Rho family GTPases [3]. Finally, toxins have been identified that covalently modify actin itself, for example by the addition of an ADP-ribosyl group that then prevents its incorporation into F-actin [4,5].

Studies over the past ten years have been directed toward characterizing a distinct mechanism to disrupt actin filament assembly by covalently crosslinking actin monomers. The crosslinks can be introduced repetitively to form long chains of dysfunctional actin that cannot be incorporated into F-actin filaments, resulting in irreversible destruction of the host cell cytoskeleton [6].

1.2. Discovery of the actin crosslinking domain of MARTX_{Vc}

An actin crosslinking toxin was first discovered in current pandemic strains of the human diarrheal pathogen *Vibrio cholerae* [6]. When added to mammalian cells in culture, *V. cholerae* was shown to secrete a factor that induced rapid cell rounding, ultimately resulting

in loss of all polymerized actin [6,7]. When actin in *V. cholerae*-treated cells was monitored by western blotting as a potential target for covalent modification, it was noted that all the monomeric actin in the cell disappeared within 180 min of addition of bacteria and actin was detected as higher weight molecular forms. The change in the electrophoretic mobility of actin corresponded to the formation of dimers, trimers, and higher order oligomers, revealing that actin in *V. cholerae*-treated cells had been crosslinked into chains [6].

This actin crosslinking activity depended upon the gene *rtxA*, the largest gene of the *V. cholerae* genome [6,7] and actin crosslinking activity was found to be widely present among both clinical and environmental isolates of *V. cholerae* [8-11]. The protein encoded by *rtxA*, now referred to as the Multifunctional, Autoprocessing RTX toxin of *V. cholerae* (MARTX_{V_C}), is the founding member of a new family of toxins [12]. The MARTX toxins are a grouping of toxins that are all large in size (>350 kDa) and are secreted from the bacteria by dedicated Type I secretion systems. Conserved features of all MARTX toxins include N- and C-terminal glycine rich repeats and an autoprocessing cysteine protease. These conserved structural elements are proposed to function in toxin translocation to deliver one to five effector domains found in the central region of the toxins to the eukaryotic cell cytosol [12].

To identify the specific effector of MARTX_{V_C} responsible for actin crosslinking, subfragments of *V. cholerae* *rtxA* gene were ectopically expressed in epithelial cells as fusions to GFP and GFP-positive cells were visualized for rounding. Cells transfected with DNA corresponding to amino acids 1963-2375 of the *rtxA* gene (according to the original annotation by Lin *et al.* [7]) resulted in cell rounding and cells were confirmed to contain crosslinked actin by western blotting. This region was thus designated the actin crosslinking domain, or ACD [13], and it is now recognized as an effector delivered by autoprocessing of the large MARTX_{V_C} toxin [14].

1.3. Other proteins with actin crosslinking domains

Subsequent to the discovery of the ACD within MARTX_{V_C}, two other MARTX toxins have been identified by protein sequence analyses that also carry ACD effectors. Genomic sequencing of *Aeromonas hydrophila* strain ATCC 7966T revealed this pathogen associated with disease in invertebrates and an emerging pathogen in humans has a gene for a MARTX toxin that includes an ACD effector [12,15]. By contrast, genomic sequencing of human pathogenic Biotype 1 *V. vulnificus* strains revealed they also encode a MARTX toxin [16,17], but this toxin was not associated with actin crosslinking and did not have an ACD [13,18]. However, an environmental *V. vulnificus* Biotype 2 strain isolated from a diseased eel was found to carry a gene for a second MARTX toxin on the pR99 virulence plasmid (MARTX_{R99}) and this toxin included an ACD [19] (Figure 1). At this time, neither of these ACDs have been confirmed biochemically to be functional for actin crosslinking.

A second protein encoded by *V. cholerae* was also found to have an ACD [13]. This protein, VgrG-1, is a bacterial effector that is secreted from *V. cholerae* dependent upon a Type 6 secretion (T6S) system [20]. The first domain of the protein belongs to the VgrG family, a group of proteins required to form a structural element of the T6S apparatuses [21]. VgrG-1 is unusual among the Vgr proteins because it is an “evolved VgrG” with an attached second domain that is delivered to the cell cytosol and carries effector function, in this case, an ACD that has been demonstrated experimentally to have actin crosslinking activity [22] (Figure 1).

The final putative actin crosslinking toxin is hypothetical protein AND4_00045 identified in the draft genome of environmental *Vibrio* sp. AND4 isolated directly from the Andaman Sea in Thailand. Unlike the MARTX toxins or VgrG-1, this protein is a stand-alone actin

crosslinking protein with no other associated domain for translocation [23]. Thus, if it is an effector used for pathogenesis of aquatic animals, it must be translocated directly to the eukaryotic cell cytosol, possibly by a Type III or T6S system also identified in the draft genome.

2. Characterization of the Actin Crosslinking Activity

2.1. ACD directly crosslinks actin

Early models of the mechanism of actin crosslinking recognized that crosslinking could occur either directly, in which the ACD is the catalytic enzyme, or indirectly, by activation of an unknown host cell protein with this activity [6]. To demonstrate direct action, the ACD portion of MARTX_{VC} was purified as a fusion with the N-terminus of Anthrax Toxin Lethal Factor (LF_N). LF_N has been shown previously to mediate cytosolic delivery of heterologous fusion proteins through pores formed by Anthrax Toxin Protective Antigen (PA) [25,26]. This approach facilitated pretesting the LF_NACD for biochemical activity in the cytosol of mammalian cells after delivery by PA, prior to initiating experiments to establish *in vitro* crosslinking [27].

Subsequent experiments showed that LF_NACD does crosslink actin both *in vivo* after delivery by PA, as well as *in vitro* when added to eukaryotic cell lysates. Further refinement of the crosslinking reaction revealed the *in vitro* reaction is catalytic and progresses to completion in less than 10 min. This crosslinking occurs under reaction conditions composed of only purified actin, ATP, and Mg²⁺, indicating that no other proteins or cellular co-factors are required for crosslinking of actin by the ACD; and thus, ACD itself is the catalytic enzyme [27].

2.2. Requirement of G-actin, ATP, and Mg²⁺ for actin crosslinking

Polymerization of monomeric G-actin into F-actin filaments is dependent upon hydrolysis of ATP [28] and occurs in the presence of 2 mM Mg²⁺. Hence, the requirement of Mg²⁺ and ATP during the *in vitro* crosslinking reaction was initially thought to indicate that actin must be polymerized to be crosslinked. However, it was shown that crosslinking *in vitro* progressed efficiently when actin was locked as monomeric G-actin by association with latrunculin or kabiramide C or by modification with tetramethyl red [29]. Similarly, actin crosslinking progressed *in vivo* after addition of MARTX_{VC}-producing *V. cholerae* to cells that were pretreated with cytochalasin or latrunculin to inhibit actin polymerization [27,30]. By contrast, stabilization of F-actin *in vitro* by the addition of phalloidin or *in vivo* using dolastatin 11 inhibited the efficiency of actin crosslinking [27,29]. These data indicated that G-actin, not F-actin, is the substrate for ACD in the crosslinking reaction. Furthermore, these data support that the requirement for ATP and Mg²⁺ during ACD-mediated actin crosslinking is not to drive actin polymerization.

Interestingly, by contrast to studies with MARTX_{VC}, crosslinking of actin in macrophages due to VgrG-1 was completely blocked by cytochalasin. However, this inhibition was not due to the ACD of VgrG-1 requiring polymerized actin instead of monomeric actin; rather it revealed a requirement for the bacteria to be phagocytosed before the VgrG-1 effector could be delivered to the cytosol by T6S [21]. Next, to account for the requirement of ATP and Mg²⁺ in the crosslinking reaction, it was considered if the ACD itself is an ATPase. During an *in vitro* crosslinking reaction, inorganic phosphate (P_i) is released in a dose dependent manner such that one molecule of ATP is used for each crosslink introduced. This ATP hydrolysis is not due to polymerization of actin, but is an essential aspect of the crosslinking reaction itself [29] (see Figure 2).

2.3. Most G-actin binding proteins do not inhibit the crosslinking reaction

It is well-recognized that in the living cell, G-actin does not exist in a free form and is typically bound to one of the several actin binding proteins (ABPs) [31]. Remarkably, prior association of actin with the common ABPs profilin, thymosin- β 4, and gelsolin did not have any noticeable effect on actin crosslinking by ACD. Among the ABPs tested, only DNaseI completely blocked the formation of actin oligomers, and cofilin partially inhibited. These data show that *in vivo*, crosslinking of actin by MARTX_{Vc} likely progresses without interference of most ABPs [29].

Furthermore, actin crosslinking can occur when actin is previously crosslinked to itself. As G-actin is the substrate for crosslinking, higher order oligomers could arise by continued addition of G-actin monomers to longer chains or by the joining of oligomers. It was found that actin dimers disappear during crosslinking at ~4 fold slower rate than the monomers. This result indicated that although ACD-crosslinked oligomers can be further joined to form higher order species, the efficiency of the reaction diminishes with increased size of oligomers [29]. Thus, the formation of long crosslinked actin chains may occur predominantly by addition of monomers to linked oligomers.

3. The Chemical Nature of the Crosslink

The chemical crosslink introduced into actin has been solved and shown to be an isopeptide linkage between glutamate 270 on one actin monomer with lysine 50 on a second actin monomer (Figure 2).

No other crosslinks are introduced and mutation of these residues within actin prevented crosslinking entirely, both *in vitro* and *in vivo* [32]. The crosslinked residues occur in flexible loops of actin in regions isolated away from binding sites for latrunculin, kabiramide C, and ABP binding sites. Lysine 50 is a residue on the DNaseI binding loop accounting for the lack of crosslinking when actin was pre-associated with DNaseI [29,32].

As seen in the crystal structure of the dimerized form of crosslinked actin (PDBID: 3CJC), the two actin molecules are tilted with respect to the normal alignment of polymerized actin. When added to a crosslinking reaction, the altered structure cannot facilitate addition of new monomers such that dimerized actin competes with monomeric actin in F-actin growth and halts further polymerization [32]. Thus, actin crosslinking ultimately rounds cells both by actively blocking growth of actin filaments and by depleting the G-actin pool.

4. The Catalytic Mechanism Is Similar to Glutamate Synthetases

Given the novel nature of the crosslink in actin, it is not surprising that little information regarding the mechanism of catalysis could be derived using bioinformatic analyses.

Protein alignments using the primary peptide sequence of the ACDs identified only that ACD proteins are highly conserved, but showed no homology to other known or hypothetical proteins, suggesting actin crosslinking by an ACD may be restricted to the *Vibrionaceae* bacteria (Figure 3).

Without sequence-based information to inform research strategies, the catalytic mechanism was probed by a random genetic approach to identify essential amino acids within the ACD [33]. This analysis revealed that glutamic acid 1990 of MARTX_{Vc} is absolutely essential for actin crosslinking while mutation of six other residues resulted in decreased actin crosslinking efficiency. No other important residues were identified. A structural alignment algorithm revealed these residues could be mapped against the conserved catalytic site of proteins in the glutamine synthetase family of ATP-dependent protein ligases. Modeling

ACD using the crystal structure of γ -glutamylcysteine synthetase (GshA) from *Escherichia coli* [34] revealed that E1990 likely participates in the transfer of the γ -phosphate from ATP to actin glutamate 270 side chain to activate the ligation reaction with lysine 50, resulting in release of P_1 (Figure 2). Thus, actin crosslinking to link glutamate to lysine occurs by a ligation reaction similar to that required to attach ammonia to glutamate to form glutamine or cysteine to glutamate to form γ -glutamylcysteine [33]. Given the strong conservation of important residues among the five identified ACDs (Figure 3), it is reasonable to predict that all ACDs will crosslink actin by an identical mechanism.

5. Phagocytic Cells as the Primary Target of ACD

The predominance of work studying the action of actin crosslinking has been conducted using epithelial cells in culture. However, the role of these toxins in disease progression is unclear because, particularly for cholera disease, there is limited epithelial damage. Recent studies have been directed toward understanding whether actin crosslinking toxins more specifically target innate immune cells. MARTX_{VC}, in conjunction with other secreted factors, has been demonstrated to be important to prevent clearance of the bacteria from the small intestine [35,36]. This effect of toxin expression on the ability of the bacteria to persist suggests MARTX_{VC} effectively blocks clearance by slowing or completely deactivating phagocytosis of the bacteria. Consistent with this model, the virulence plasmid that encodes MARTX_{R99} promotes survival of *V. vulnificus* Biotype 2 strains in eel whole blood suggesting it may inactivate phagocytes [19]. Furthermore, the actin crosslinking Type VI secretion effector VgrG-1 from *V. cholerae* crosslinks actin within phagocytic macrophages and amoebae, but not epithelial cells [20,21]. Finally, *A. hydrophila* has been demonstrated to evade phagocytosis, although this may be as much or more related to its T6S system and associated actin ADP-ribosylating effector [5,37]. Thus, it seems possible that the function of the actin crosslinking toxins could be to directly block bacterial engulfment by phagocytic cells to evade clearance from the site of infection. This has been experimentally demonstrated for the function of VgrG-1 proteins and infection data suggest other actin crosslinking toxins may function in similar capacity to facilitate colonization and disease [38].

Acknowledgments

K.J.F.S is funded by an Investigator in Pathogenesis of Infectious Diseases Award from the Burroughs Wellcome Fund and by the National Institutes of Health Awards AI051490 and AI072461.

References

1. Espina B, Rubiolo JA. Marine toxins and the cytoskeleton: pectenotoxins, unusual macrolides that disrupt actin. *FEBS J* 2008;275:6082–6088. [PubMed: 19016860]
2. Natori S. Cytochalasins-actin filament modifiers as a group of mycotoxins. *Dev Toxicol Environ Sci* 1986;12:291–299. [PubMed: 3549232]
3. Aktories K, Barbieri JT. Bacterial cytotoxins: Targeting eukaryotic switches. *Nature Rev* 2005;3:397–410.
4. Richard JF, Petit L, Gibert M, Marvaud JC, Bouchaud C, Popoff MR. Bacterial toxins modifying the actin cytoskeleton. *Int Microbiol* 1999;2:185–194. [PubMed: 10943412]
5. Suarez G, Sierra JC, Erova TE, Sha J, Horneman AJ, Chopra AK. A type VI secretion system effector protein VgrG1 from *Aeromonas hydrophila* that induces host cell toxicity by ADP-ribosylation of actin. *J Bacteriol*. 2009 published online 10/30/2009.
6. Fullner KJ, Mekalanos JJ. *In vivo* covalent crosslinking of actin by the RTX toxin of *Vibrio cholerae*. *EMBO J* 2000;19:5315–5323. [PubMed: 11032799]

7. Lin W, Fullner KJ, Clayton R, Sexton JA, Rogers MB, Calia KE, Calderwood SB, Fraser C, Mekalanos JJ. Identification of a *Vibrio cholerae* RTX toxin gene cluster that is tightly linked to the cholera toxin prophage. *Proc Natl Acad Sci USA* 1999;96:1071–1076. [PubMed: 9927695]
8. Chow KH, Ng TK, Yuen KY, Yam WC. Detection of RTX Toxin Gene in *Vibrio cholerae* by PCR. *J Clin Microbiol* 2001;39:2594–2597. [PubMed: 11427575]
9. Cordero CL, Sozhamannan S, Satchell KJ. RTX toxin actin cross-linking activity in clinical and environmental isolates of *Vibrio cholerae*. *J Clin Microbiol* 2007;45:2289–2292. [PubMed: 17522276]
10. Dalsgaard A, Serichantalergs O, Forslund A, Lin W, Mekalanos J, Mintz E, Shimada T, Wells JG. Clinical and environmental isolates of *Vibrio cholerae* serogroup O141 carry the CTX phage and the genes encoding the toxin-coregulated pili. *J Clin Microbiol* 2001;39:4086–4092. [PubMed: 11682534]
11. Rahman MH, Biswas K, Hossain MA, Sack RB, Mekalanos JJ, Faruque SM. Distribution of genes for virulence and ecological fitness among diverse *Vibrio cholerae* population in a cholera endemic area: tracking the evolution of pathogenic strains. *DNA Cell Biol* 2008;27:347–355. [PubMed: 18462070]
12. Satchell KJ. MARTX: Multifunctional-Autoprocessing RTX Toxins. *Infect Immun* 2007;75:5079–5084. [PubMed: 17646359]
13. Sheahan KL, Cordero CL, Satchell KJ. Identification of a domain within the multifunctional *Vibrio cholerae* RTX toxin that covalently cross-links actin. *Proc Natl Acad Sci USA* 2004;101:9798–9803. [PubMed: 15199181]
14. Prochazkova K, Shuvalova LA, Minasov G, Voburka Z, Anderson WF, Satchell KJ. Structural and molecular mechanism for autoprocessing of MARTX Toxin of *Vibrio cholerae* at multiple sites. *J Biol Chem* 2009;284:26557–26568. [PubMed: 19620709]
15. Seshadri R, Joseph SW, Chopra AK, Sha J, Shaw J, Graf J, Haft D, Wu M, Ren Q, Rosovitz MJ, Madupu R, Tallon L, Kim M, Jin S, Vuong H, Stine OC, Ali A, Horneman AJ, Heidelberg JF. Genome sequence of *Aeromonas hydrophila* ATCC 7966T: jack of all trades. *J Bacteriol* 2006;188:8272–8282. [PubMed: 16980456]
16. Chen CY, Wu KM, Chang YC, Chang CH, Tsai HC, Liao TL, Liu YM, Chen HJ, Shen AB, Li JC, Su TL, Shao CP, Lee CT, Hor LI, Tsai SF. Comparative genome analysis of *Vibrio vulnificus*, a marine pathogen. *Genome Res* 2003;13:2577–2587. [PubMed: 14656965]
17. Kim YR, Lee SE, Kim CM, Kim SY, Shin EK, Shin DH, Chung SS, Choy HE, Progulsk-Fox A, Hillman JD, Handfield M, Rhee JH. Characterization and pathogenic significance of *Vibrio vulnificus* antigens preferentially expressed in septicemic patients. *Infect Immun* 2003;71:5461–5471. [PubMed: 14500463]
18. Kim YR, Lee SE, Kook H, Yeom JA, Na HS, Kim SY, Chung SS, Choy HE, Rhee JH. *Vibrio vulnificus* RTX toxin kills host cells only after contact of the bacteria with host cells. *Cell Microbiol* 2008;10:848–862. [PubMed: 18005241]
19. Lee CT, Amaro C, Wu KM, Valiente E, Chang YF, Tsai SF, Chang CH, Hor LI. A common virulence plasmid in biotype 2 *Vibrio vulnificus* and its dissemination aided by a conjugal plasmid. *J Bacteriol* 2008;190:1638–1648. [PubMed: 18156267]
20. Pukatzki S, Ma AT, Sturtevant D, Krastins B, Sarracino D, Nelson WC, Heidelberg JF, Mekalanos JJ. Identification of a conserved bacterial protein secretion system in *Vibrio cholerae* using the *Dictyostelium* host model system. *Proc Natl Acad Sci USA* 2006;103:1528–1533. [PubMed: 16432199]
21. Ma AT, McAuley S, Pukatzki S, Mekalanos JJ. Translocation of a *Vibrio cholerae* type VI secretion effector requires bacterial endocytosis by host cells. *Cell Host Microbe* 2009;5:234–243. [PubMed: 19286133]
22. Pukatzki S, Ma AT, Revel AT, Sturtevant D, Mekalanos JJ. Type VI secretion system translocates a phage tail spike-like protein into target cells where it cross-links actin. *Proc Natl Acad Sci USA* 2007;104:15508–15513. [PubMed: 17873062]
23. Hagstrom A, Ferreira S, Johnson J, Kravitz S, Beeson K, Sutton G, Rogers Y-H, Friedman R, Frazier M, Venter JC. *Vibrio* sp. AND4 1103602000423, whole genome shotgun sequence, direct submission. NCBI reference sequence NZ_ABGR01000041.1. 2007

24. Sheahan KL, Satchell KJ. Inactivation of small Rho GTPases by the multifunctional RTX toxin from *Vibrio cholerae*. *Cell Microbiol* 2007;9:1324–1335. [PubMed: 17474905]
25. Milne JC, Blanke SR, Hanna PC, Collier RJ. Protective antigen-binding domain of anthrax lethal factor mediates translocation of a heterologous protein fused to its amino- or carboxy-terminus. *Mol Microbiol* 1995;15:661–666. [PubMed: 7783638]
26. Spyres LM, Qa'Dan M, Meader A, Tomasek JJ, Howard EW, Ballard JD. Cytosolic delivery and characterization of the TcdB glucosylating domain by using a heterologous protein fusion. *Infect Immun* 2001;69:599–601. [PubMed: 11119561]
27. Cordero CL, Kudryashov DS, Reisler E, Satchell KJ. The actin cross-linking domain of the *Vibrio cholerae* RTX toxin directly catalyzes the covalent cross-linking of actin. *J Biol Chem* 2006;281:32366–32374. [PubMed: 16954226]
28. Korn ED, Carlier MF, Pantaloni D. Actin polymerization and ATP hydrolysis. *Science* 1987;238:638–644. [PubMed: 3672117]
29. Kudryashov DS, Cordero CL, Reisler E, Satchell KJ. Characterization of the enzymatic activity of the actin cross-linking domain from the *Vibrio cholerae* MARTX_{VC} toxin. *J Biol Chem* 2008;283:445–452. [PubMed: 17951576]
30. Fullner KJ, Mekalanos JJ. Genetic characterization of a new type IV pilus gene cluster found in both classical and El Tor biotypes of *Vibrio cholerae*. *Infect Immun* 1999;67:1393–1404. [PubMed: 10024587]
31. Pollard TD, Borisy GG. Cellular motility driven by assembly and disassembly of actin filaments. *Cell* 2003;112:453–465. [PubMed: 12600310]
32. Kudryashov DS, Durer ZA, Ytterberg AJ, Sawaya MR, Pashkov I, Prochazkova K, Yeates TO, Loo RR, Loo JA, Satchell KJ, Reisler E. Connecting actin monomers by iso-peptide bond is a toxicity mechanism of the *Vibrio cholerae* MARTX toxin. *Proc Natl Acad Sci USA* 2008;105:18537–18542. [PubMed: 19015515]
33. Geissler B, Bonebrake A, Sheahan KL, Walker ME, Satchell KJ. Genetic determination of essential residues of the *Vibrio cholerae* actin cross-linking domain reveals functional similarity with glutamine synthetases. *Mol Microbiol* 2009;73:858–868. [PubMed: 19656298]
34. Hibi T, Nii H, Nakatsu T, Kimura A, Kato H, Hiratake J, Oda J. Crystal structure of gamma-glutamylcysteine synthetase: insights into the mechanism of catalysis by a key enzyme for glutathione homeostasis. *Proc Natl Acad Sci USA* 2004;101:15052–15057. [PubMed: 15477603]
35. Olivier V, Queen J, Satchell KJ. Successful small intestine colonization of adult mice by *Vibrio cholerae* requires ketamine anesthesia and accessory toxins. *PLoS One* 2009;4:e7352. [PubMed: 19812690]
36. Olivier V, Salzman NH, Satchell KJ. Prolonged colonization of mice by *Vibrio cholerae* El Tor O1 depends on accessory toxins. *Infect Immun* 2007;75:5043–5051. [PubMed: 17698571]
37. Suarez G, Sierra JC, Sha J, Wang S, Erova TE, Fadl AA, Foltz SM, Horneman AJ, Chopra AK. Molecular characterization of a functional type VI secretion system from a clinical isolate of *Aeromonas hydrophila*. *Microbial Pathog* 2008;44:344–361.
38. Satchell KJ. Bacterial martyrdom: Phagocytes disabled by type VI secretion after engulfing bacteria. *Cell Host Microbe* 2009;5:213–214. [PubMed: 19286128]




Figure 1.

Actin crosslinking toxins. Scaled diagrams of the five known and putative actin crosslinking toxins. MARTX toxins of *V. cholerae* (MARTX_{Vc}), *A. hydrophila* (MARTX_{Ah}), and *V. vulnificus* Biotype 2 virulence plasmid pR99 (MARTX_{R99}). Conserved MARTX repeat regions (grey) are predicted to converge at the eukaryotic plasma membrane to form a pore through which the centrally-located effectors are translocated. After translocation, the cysteine protease domain (CPD) initiates autoprocessing to release effectors, including the actin crosslinking domain (ACD-red), Rho-inactivation domain (RID-purple) [24], α/β hydrolase effector (α/β -green), Mcf conserved effector of unknown function (Mcf-orange), and *Pasteurella multocida* toxin conserved effector of unknown function (PMT C1/C2-brown)[12]. Note that MARTX_{R99} has a duplication of the Mcf effector. *V. cholerae* VgrG-1 is an effector of the *V. cholerae* VAS T6S system. The Vgr conserved domain (Vgr-pink) forms part of the translocation apparatus for transfer of the ACD across the phagosomal membrane after engulfment by phagocytes. *Vibrio* sp. AND4 hypothetical protein AND4_00045 is a putative stand-alone actin crosslinking toxin that is potentially an effector of an unknown transport apparatus.




Figure 2.

Crosslink introduced between actin side chains. Schematic shows E270 and K50 side chains originating from two different actin monomers with dashed box indicating H₂O molecule lost during the ligation reaction. Lower equation depicts the predicted phosphotransfer based on the mechanism for formation of γ -glutamylcysteine from glutamic acid and cysteine.




Figure 3.

Alignment of ACD amino acid sequences from actin crosslinking toxins. Alignment was generated with CLUSTALW using MacVector 9.5.2 software based on sequences available at NCBI PubMed with accession numbers: *V. cholerae* VgrG1 ACD (NP_231059.1); *Vibrio sp.* AND4_00045 ACD (ZP_02197558.1); *V. cholerae* (Vc) MARTX ACD (AAD21057.1); *A. hydrophila* (Ah) MARTX ACD (YP_855898.1); and *V. vulnificus* Biotype 2 virulence plasmid pR99 MARTX ACD (YP_001393065.1). Red boxes indicate residues identified by genetic analysis as important for crosslinking with the star indicating the glutamic acid required for phosphotransfer. Black arrows represent predicted beta-strands that might form a glutamine synthetase ligase-like active site (see Figure 4).




Figure 4. Structural model of *E. coli* GshA. *E. coli* GshA (PDB 1VA6; [34]) active site is composed of a beta-sheet (grey) that contributes amino acid side chains that bind three Mg^{2+} ions (blue) and ATP (ADP in crystal structure, red). The active site shares conserved residues with ACD amino acids demonstrated to be important for crosslinking activity (magenta). The essential ACD glutamate residue (green) is likely required for phosphotransfer during the ligation reaction as shown in Figure 2. Model prepared using MacPyMol (Delano Scientific) based on information from Geissler *et al.* [33].