



Published in final edited form as:

J Gen Virol. 2010 May ; 91(Pt 5): 1311–1314. doi:10.1099/vir.0.019307-0.

The HR motif in the RNA-dependent RNA polymerase L protein of Chandipura virus is required for unconventional mRNA capping activity

Tomoaki Ogino and Amiya K. Banerjee

Department of Molecular Genetics, Section of Virology, Lerner Research Institute, Cleveland Clinic, Cleveland, Ohio 44195, USA

Summary

Chandipura virus (CHPV) is an emerging human pathogen associated with acute encephalitis and is closely related to vesicular stomatitis virus (VSV), a prototype of rhabdoviruses. Here, we demonstrate that the RNA polymerase L protein of CHPV exhibits a VSV-like RNA:GDP polyribonucleotidyltransferase (PRNTase) activity, which transfers the 5'-monophosphorylated viral mRNA-start sequence to GDP to produce a capped RNA, and the conserved HR motif in the CHPV L protein is essential for the PRNTase activity. Interestingly, the CHPV L protein was found to form two distinct SDS-resistant complexes with the CHPV mRNA- and leader RNA-start sequences; mutations in the HR motif significantly reduced the formation of the former complex (a putative covalent enzyme-pRNA intermediate in the PRNTase reaction), but not the latter complex. These results suggest that the rhabdoviral L proteins universally use the active site HR motif for the PRNTase reaction at the step of the enzyme-pRNA intermediate formation.

Text

Unconventional mRNA capping of vesicular stomatitis virus (VSV, Indiana), a prototype nonsegmented negative-strand (NNS) RNA virus belonging to the genus *Vesiculovirus* of the family *Rhabdoviridae* in the order *Mononegavirales*, is catalyzed by the multifunctional RNA-dependent RNA polymerase (RdRp) L protein (Ogino & Banerjee, 2007; 2008). At the first step, the guanosine 5'-triphosphatase (GTPase) activity of the VSV L protein produces GDP (an RNA acceptor) by hydrolysis of GTP (Ogino & Banerjee, 2007; 2008). Subsequently, the RNA:GDP polyribonucleotidyltransferase (PRNTase) domain in the VSV L protein transfers the 5'-monophosphorylated (p-) RNA moiety of the 5'-triphosphorylated (ppp-) RNA with the conserved VSV mRNA-start sequence (AACAG) to GDP to generate the Gpp-pA capped RNA through a covalent enzyme-pRNA intermediate (Ogino & Banerjee, 2007; 2008; Ogino *et al.*, 2010). The latter intermediate is composed of pRNA covalently linked to the active site histidine residue at position 1,227 (H1227) within the conserved HR (H1227-R1228) motif via a phosphoamide bond (Ogino *et al.*, 2010). By mutational analyses, we have recently demonstrated that the HR motif and a basic amino

Correspondence: Amiya K. Banerjee, Tel.: +1 (216) 444-0625, Fax: +1 (216) 444-2998, banerja@ccf.org.

This is an author manuscript that has been accepted for publication in *Journal of General Virology*, copyright Society for General Microbiology, but has not been copy-edited, formatted or proofed. Cite this article as appearing in *Journal of General Virology*. This version of the manuscript may not be duplicated or reproduced, other than for personal use or within the rule of 'Fair Use of Copyrighted Materials' (section 17, Title 17, US Code), without permission from the copyright owner, Society for General Microbiology. The Society for General Microbiology disclaims any responsibility or liability for errors or omissions in this version of the manuscript or in any version derived from it by any other parties. The final copy-edited, published article, which is the version of record, can be found at <http://vir.sgmjournals.org>, and is freely available without a subscription.

acid residue (R1221) in the vicinity of the HR motif are necessary for the PRNTase activity of the VSV L protein at the step of the enzyme-pRNA intermediate formation (Ogino *et al.*, 2010). The HR motif is strikingly conserved in the L proteins of more than 80 NNS RNA viruses (e.g., rabies, measles, mumps, Ebola, and Borna disease viruses) belonging to different families, while the R1221 residue is conserved only in the L proteins of vesiculoviruses (e.g., VSV, Chandipura virus [CHPV]), lyssaviruses (e.g., rabies virus), and ephemerovirus (bovine ephemeral fever virus) belonging to the *Rhabdoviridae* family (Ogino *et al.*, 2010).

To examine the universality of the RNA capping mechanism by the rhabdoviral L proteins, we generated the recombinant L protein of CHPV, an emerging vesiculovirus that causes acute viral encephalitis in children with high mortality in India (Chadha *et al.*, 2005; Rao *et al.*, 2004). The recombinant CHPV L protein was expressed as a carboxyl-terminal octahistidine-tagged form (2,092 plus 10 [SRHHHHHHHH] amino acids) in Sf21 insect cells infected with a recombinant baculovirus carrying the cDNA encoding the CHPV L protein (Marriott, 2005), and purified by chromatography using nickel-nitrilotriacetic acid agarose (Qiagen), as described for the recombinant VSV L protein (Ogino & Banerjee, 2007; Ogino *et al.*, 2010). First, we measured the RNA capping activity of the CHPV L protein (Fig. 1) according to the method developed for the VSV L protein (Ogino & Banerjee, 2007; 2008; Ogino *et al.*, 2010). To analyze the substrate specificity of the CHPV L protein (PRNTase), we used 5'-tri- (ppp-) or di- (pp-) phosphorylated oligo-RNAs corresponding to the CHPV mRNA- and leader RNA-start sequences (AACAG and ACGAA, respectively), which are same as VSV (Marriott, 2005), and [α -³²P]GTP or [α -³²P]GDP. When pppAACAG was incubated with [α -³²P]GTP in the presence of the CHPV L protein, a distinct GpppA cap structure was produced (lane 2). Similar to the VSV L protein, ppAACAG and ppACGAA did not serve as the substrate for the CHPV L protein (lanes 3 and 4). Since the cellular capping enzyme composed of RNA 5'-triphosphatase and GTP:RNA guanylyltransferase can use ppp- or pp-RNA with any sequence as the substrate to produce Gp-ppRNA (Furuichi & Shatkin, 2000; Shuman, 2001), the capping activity associated with the CHPV L protein is different from the cellular activity, thus, is thought to be intrinsic to the viral protein. Furthermore, the CHPV L protein was found to produce the GpppA cap structure with [α -³²P]GDP (an inert substrate for the cellular capping enzyme) instead of [α -³²P]GTP (lane 6), indicating that the CHPV L protein catalyzes a similar unconventional RNA capping reaction like VSV by the PRNTase activity. Under standard conditions, the CHPV L protein (0.3 μ g) produced 0.74 ± 0.04 fmol (the mean \pm standard deviation of three independent determinations) of GpppA with [α -³²P]GDP. The specific activity of the CHPV L protein in RNA capping with GDP (2.5 fmol GpppA/ μ g protein) was approximately 800-fold lower than that of the VSV L protein (2.1 pmol GpppA/ μ g protein) (Ogino *et al.*, 2010), despite its 60% amino acid sequence identity to the VSV L protein (Marriott, 2005). It is noteworthy that the transcriptase activity of CHPV is significantly lower than that of VSV (Chang *et al.*, 1974).

To ascertain the significance of the R1211, H1217, and R1218 residues in the CHPV L protein [the counterparts of the R1221, H1227, and R1228 residues of the VSV L protein (Fig. 2a)] in the PRNTase activity, these residues were replaced with alanine (Fig. 2b, lanes 2–4). In addition, we prepared a CHPV L mutant with an alanine substitution for the non-conserved H1226 residue (the counterpart of the non-essential H1236 residue of the VSV L protein, Fig 2a) (Fig. 2b, lane 5). As shown in Fig. 2c (lanes 3–6), the R1211A, H1217A, and R1218A mutants, but not the H1226A mutant, were completely inactive in RNA capping with pppAACAG and [α -³²P]GDP, indicating that the CHPV L protein uses these active site amino acid residues (R1211, H1217, and R1218), which are identical to those in the VSV L protein, for the VSV-like PRNTase activity.

Our demonstration that the CHPV L protein exhibits the capping (PRNTase) activity prompted us to detect a covalent CHPV L protein-pRNA complex, an enzyme-pRNA intermediate in the PRNTase reaction. As described for the VSV L protein (Ogino & Banerjee, 2007; Ogino *et al.*, 2010), the enzyme-pRNA intermediate (L-pRNA complex) formation assay was performed using the purified recombinant CHPV L protein. Briefly, the CHPV L protein was incubated with ³²P-labeled pppRNA, and the resulting complexes were analyzed by SDS-7.5% PAGE followed by autoradiography. As shown in Fig. 3a (lane 2), the CHPV L protein reacted with pppAACAG (mRNA-start sequence), resulting in generation of an SDS-resistant L-RNA complex, but not with ppAACAG (lane 3). Interestingly, the CHPV L protein formed an L-RNA complex with pppACGAA (leader RNA-start sequence, lane 4) to a similar extent as that formed with the mRNA-start sequence (lane 2). This result is in contrast to the VSV L protein, which reacts with pppAACAG greater than with pppACGAA to form the covalent L-pRNA complex (the intermediate in the PRNTase reaction) (Ogino & Banerjee, 2007). On the other hand, the CHPV L protein did not complex with [α -³²P]ATP (lane 5), indicating that the SDS-resistant complex formation is specific to polynucleotides, similar to the VSV L protein (Ogino & Banerjee, 2007). Finally, we examined effects of the mutations into the PRNTase active site of the CHPV L protein on the formation of the SDS-resistant L-RNA complexes with the mRNA- and leader- RNA sequences. As shown in Fig. 3b (lanes 3–5), the R1211A, H1217A, and R1218A mutations greatly reduced their activities of the L-RNA complex formation with the pppAACAG mRNA-start sequence, but the H1226A mutation did not (lane 6). In contrast, all mutations (R1211A, H1217A, R1218A, and H1226A) did not affect the L-RNA complex formation with the pppACGAA leader RNA-start sequence (Fig. 3c, lanes 3–6). Therefore, the complex formation between the CHPV L protein and the pppACGAA leader RNA-start sequence does not appear to be involved in the RNA capping reaction. Thus, we conclude that the conserved HR motif in the CHPV L protein is essential for the PRNTase activity at the step of the enzyme-pRNA intermediate (L-pRNA complex) formation. The H1217 residue in the HR motif is strongly suggested to be a covalent RNA attachment site in the PRNTase domain of the CHPV L protein, because the histidine residue in the HR motif of the VSV L protein has been identified as the covalent RNA attachment site (Ogino *et al.*, 2010). On the other hand, since the R1211 residue (a counterpart of the R1221 residue in the VSV L protein) is conserved only in the L proteins of some rhabdoviruses (vesiculo-, lyssa-, and ephemero- viruses), it is suggested to be involved in some important step of the enzyme-pRNA intermediate formation that is specific for above rhabdoviruses, *e.g.*, sequence-specific recognition of the 5'-triphosphorylated RNA substrate. Interestingly, the mRNAs of above rhabdoviruses contain conserved mRNA-start sequences (5'-AACAA) (Bourhy *et al.*, 1993; Hoffmann *et al.*, 2002; Marriott, 2005; McWilliam *et al.*, 1997), whereas mRNAs of other NNS RNA viruses have unique sets of mRNA-start sequences. It should also be noted that all cap-deficient mutants (R1211A, H1217A, and R1218A) of the CHPV L protein retained very weak activities of binding with the mRNA-start sequence (Fig. 3b, lanes 3–5). It is, thus, conceivable that the CHPV L protein contains, in addition to the HR motif, other RNA attachment sites, which are probably not involved in RNA capping. Further studies are needed to define the roles of such putative RNA attachment sites in RNA synthesis, if any.

Here, we showed that the CHPV L protein selectively uses pppAACAG (the CHPV mRNA-start sequence) and GDP as the substrates to produce the GpppA cap structure, indicating that the CHPV L protein also carries out the unconventional mRNA capping mechanism involving the PRNTase activity. Furthermore, as in the case of the VSV L protein, the HR motif in the CHPV L protein was shown to be crucial for the PRNTase activity at the step of the enzyme-pRNA intermediate formation. Our findings provide the second example of an RNA transfer enzyme with the conserved HR motif that catalyzes unconventional capping of rhabdoviral mRNA, thus, offer a common target for developing anti-rhabdoviral agents.

Acknowledgments

We thank Anthony C. Marriott (University of Warwick, UK) for the plasmid encoding the CHPV L protein. This work was supported by a grant from the National Institutes of Health (AI26585, to A.K.B.).

References

- Bourhy H, Kissi B, Tordo N. Molecular diversity of the Lyssavirus genus. *Virology* 1993;194:70–81. [PubMed: 8386891]
- Chadha MS, Arankalle VA, Jadi RS, Joshi MV, Thakare JP, Mahadev PV, Mishra AC. An outbreak of Chandipura virus encephalitis in the eastern districts of Gujarat state, India. *Am J Trop Med Hyg* 2005;73:566–570. [PubMed: 16172482]
- Chang SH, Hefti E, Obijeski JF, Bishop DH. RNA transcription by the virion polymerases of five rhabdoviruses. *J Virol* 1974;13:652–661. [PubMed: 4362867]
- Furuichi Y, Shatkin AJ. Viral and cellular mRNA capping: past and prospects. *Adv Virus Res* 2000;55:135–184. [PubMed: 11050942]
- Hoffmann B, Schutze H, Mettenleiter TC. Determination of the complete genomic sequence and analysis of the gene products of the virus of Spring Viremia of Carp, a fish rhabdovirus. *Virus Res* 2002;84:89–100. [PubMed: 11900842]
- Marriott AC. Complete genome sequences of Chandipura and Isfahan vesiculoviruses. *Arch Virol* 2005;150:671–680. [PubMed: 15614433]
- McWilliam SM, Kongsuwan K, Cowley JA, Byrne KA, Walker PJ. Genome organization and transcription strategy in the complex GNS-L intergenic region of bovine ephemeral fever rhabdovirus. *J Gen Virol* 1997;78:1309–1317. [PubMed: 9191923]
- Ogino T, Banerjee AK. Unconventional mechanism of mRNA capping by the RNA-dependent RNA polymerase of vesicular stomatitis virus. *Mol Cell* 2007;25:85–97. [PubMed: 17218273]
- Ogino T, Banerjee AK. Formation of guanosine(5')tetraphospho(5')adenosine cap structure by an unconventional mRNA capping enzyme of vesicular stomatitis virus. *J Virol* 2008;82:7729–7734. [PubMed: 18495767]
- Ogino T, Yadav SP, Banerjee AK. Histidine-mediated RNA transfer to GDP for unique mRNA capping by vesicular stomatitis virus RNA polymerase. *Proc Natl Acad Sci U S A*. 2010 in press.
- Rao BL, Basu A, Wairagkar NS, Gore MM, Arankalle VA, Thakare JP, Jadi RS, Rao KA, Mishra AC. A large outbreak of acute encephalitis with high fatality rate in children in Andhra Pradesh, India, in 2003, associated with Chandipura virus. *Lancet* 2004;364:869–874. [PubMed: 15351194]
- Shuman S. Structure, mechanism, and evolution of the mRNA capping apparatus. *Prog Nucleic Acid Res Mol Biol* 2001;66:1–40. [PubMed: 11051760]




Fig. 2. The HR motif is essential for the PRNTase activity of the CHPV L protein

(a) A schematic structure of the CHPV L protein (2,092 amino acids) is shown with six amino acid sequence blocks (I–VI) conserved in the NNS RNA viral L proteins. The positions of the putative RNA-dependent RNA polymerase (RdRp) and cap methyltransferase (MTase) domains are indicated on the top. The local sequence containing the HR motif in the CHPV L protein is compared with that in the VSV L protein (H1227, the covalent RNA attachment site).

(b) Wild-type (WT) and mutant CHPV L proteins with indicated amino acid substitutions (0.7 μ g) were analyzed by SDS-7.5% PAGE followed by staining with Coomassie Brilliant Blue. The positions of marker proteins are indicated on the left.

(c) The WT and mutant CHPV L proteins (0.3 μg) were subjected to the RNA capping reactions with pppAACAG and [α - ^{32}P]GDP as substrates. Lane 1 indicates no L protein.




Fig. 3. The HR motif in the CHPV L protein is required for the formation of the enzyme-pRNA intermediate in the PRNTase reaction

(a) The recombinant CHPV L protein (0.3 μg) was incubated with the indicated mono- or poly-ribonucleotide labeled with ³²P (shown by asterisks), and then analyzed by SDS-7.5% PAGE followed by autoradiography. Note that all nucleotide probes had same specific radioactivities. Lane 1 indicates no L protein. The position of covalent L-RNA complexes is indicated by the arrowhead.

(b, c) The WT and mutant CHPV L proteins (0.3 μg) were incubated with pppAACAG (mRNA-start sequence, b) or pppACGAA (leader RNA-start sequence, c) labeled with ³²P

(shown by asterisks). The resulting covalent L-RNA complexes were analyzed by SDS-7.5% PAGE followed by autoradiography. Lane 1 indicates no L protein.