



HHS Public Access

Author manuscript

Neuropsychopharmacology. Author manuscript; available in PMC 2011 April 01.

Published in final edited form as:

Neuropsychopharmacology. 2010 October ; 35(11): 2292–2304. doi:10.1038/npp.2010.107.

Altered dopamine modulation of inhibition in the prefrontal cortex of cocaine-sensitized rats

Sven Kroener, Ph.D. and Antonieta Lavin, Ph.D.

Department of Neurosciences, Medical University of South Carolina, 173 Ashley Avenue, Charleston, SC 29425

Abstract

A functionally hypoactive prefrontal cortex (PFC) is thought to contribute to decreased cognitive inhibitory control over drug-seeking behavior in cocaine addicts. Alterations in PFC dopamine (DA) and GABA transmission are involved in the development of behavioral sensitization to cocaine, and repeated exposure to cocaine decreases DA D2 receptor (D2R) function in the PFC. We used recordings in PFC slices from adult rats to investigate how repeated cocaine treatment followed by 2 weeks of withdrawal affects DA modulation of GABA transmission and interneuron firing. In agreement with previous results in drug-naïve animals (Seamans, *et al* 2001; Trantham-Davidson, *et al* 2004) we found that in saline-treated control animals DA (20 μ M) modulated evoked inhibitory synaptic currents (eIPSCs) in a biphasic, time- and receptor-dependent manner. Activation of D2Rs transiently reduced, while D1 receptor activation persistently increased the amplitude of eIPSCs. In cocaine-sensitized animals the D2R-dependent modulation of eIPSCs was abolished and the time course of DA effects was altered. In both saline- and cocaine-treated animals the effects of DA on eIPSCs were paralleled by distinct changes in spontaneous IPSCs. In cocaine-treated animals the alterations in DA modulation of eIPSCs and sIPSCs correlated with a lack of D2R-specific reduction in action potential-independent GABA release, which might normally oppose D1-dependent increases in GABA transmission. Recordings from interneurons furthermore show that D2R activation can increase current-evoked spike firing in saline- but not in cocaine-treated animals. Altered DA regulation of inhibition during cocaine withdrawal could disturb normal cortical processing and contribute to a hypoactive PFC.

Keywords

prefrontal cortex; GABA; D2 receptor; D1 receptor; electrophysiology; addiction

Users may view, print, copy, download and text and data- mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use: http://www.nature.com/authors/editorial_policies/license.html#terms

Correspondence should be addressed to: Sven Kroener, PhD, School of Behavioral and Brain Sciences, University of Texas at Dallas, 800 West Campbell Rd., GR41, Richardson TX 75080, phone: 972-883-2039, kroener@utdallas.edu.

Disclosure: The authors declare that except for income received from their primary employers, no financial support or compensation was received from any individual or corporate entity over the past 3 years for research or professional service and there are no personal financial holdings that could be perceived as constituting a potential conflict of interest.

Introduction

Addiction to cocaine involves neurobiological changes in limbic and cortical circuits that mediate a variety of cognitive and emotional processes. Several lines of evidence suggest that a functionally “hypoactive” frontal cortex results in decreased cognitive inhibitory control to regulate drug-seeking behavior. Areas of the prefrontal cortex (PFC) are activated in addicted subjects during intoxication and exposure to drug-associated cues, but they are deactivated during withdrawal (Goldstein and Volkow, 2002; Bolla, *et al* 2004). Similarly, in rats inhibition of the medial frontal cortex blocks reinstatement of drug seeking (McFarland and Kalivas, 2001; McLaughlin and See, 2003; McFarland, *et al* 2004), and the suppression of drug seeking during extinction requires activation of the infralimbic cortex (Peters, *et al* 2008). Interestingly, cue-induced drug-seeking selectively increases expression of the immediate early gene *c-Fos* in interneurons of the PFC, while at the same time it reduces markers of activity in pyramidal cells (Miller and Marshall, 2004). Similarly, cocaine-sensitized animals showed an increase in extracellular GABA in the medial PFC (Jayaram and Steketee, 2005). These results suggest that in cocaine-sensitized animals interneurons may be preferentially activated, while the overall PFC output (the activity of pyramidal cells) is attenuated. Dopamine (DA) projections from the VTA provide a modulatory input to the PFC that is essential for optimal cognitive performance and this projection may be involved in sensitization processes (Brozoski, *et al* 1979; Steketee, 2003). Animals with a history of cocaine self-administration show deficits in tasks that require an intact medial PFC, an effect that is accompanied by a significant decrease in D2 receptors (Briand, *et al* 2008). Activation of D2 receptors (D2R) in the PFC of sensitized animals is less effective due to elevated levels of AGS3, a member of a family of G protein regulators that decreases signaling via *G α* (Bowers, *et al* 2004). Dopaminergic fibers contact both pyramidal cells and GABAergic interneurons in the PFC of rats and primates (LeMoine and Gaspar, 1998; Sesack, *et al* 1995), and D2R activation modulates interneuron activity and GABAergic synaptic transmission in drug-naïve animals (Seamans, *et al* 2001; Trantham-Davidson, *et al* 2004; Tseng and O’Donnell, 2007).

We used whole-cell recordings in PFC slices from adult animals treated repeatedly with cocaine and then subjected to long-term withdrawal, in order to assess changes in the DAergic modulation of PFC interneurons and GABAergic transmission. We hypothesized that in cocaine-sensitized animals the effect of D2R activation is blunted, so that D1 receptor (D1R)-dependent increases in GABAergic synaptic transmission dominate. Our results support this idea, showing that the D2R-dependent transient reduction in the amplitude of synaptically evoked inhibitory synaptic currents (eIPSCs) seen in saline-treated rats is abolished in cocaine-sensitized animals. In contrast, both in saline and cocaine-treated animals D1R-dependent modulation of IPSCs was unaffected, leading to increased eIPSC amplitudes, which was paralleled by an increase in the frequency of spontaneous IPSCs. In saline-treated animals the effects of D1R stimulation were at least partly countered by a D2R-dependent reduction of GABA release measured as action potential-independent release of miniature IPSCs. In cocaine-treated animals this D2R-dependent reduction in GABA release was not observed. Moreover, current clamp recordings from interneurons also show that D2R modulation of spike firing is disrupted in cocaine sensitized animals.

Taken together these results show opposing D1R and D2R-dependent mechanisms that contribute to DA modulation of GABAergic synaptic transmission in the cortex of saline-treated animals. Chronic exposure to cocaine and withdrawal that induces behavioral sensitization selectively disrupts D2R-dependent signaling. Altered DA regulation of inhibition during cocaine withdrawal could disturb normal cortical processing and contribute to a hypoactive PFC.

Materials and Methods

All procedures were carried out in accordance with the NIH Guide for the Care and Use of Laboratory Animals, and were approved by the Medical University of South Carolina Animal Care and Use Committee. Subjects were adult male Sprague Dawley rats (Charles River Laboratories, Wilmington, MA), that were older than 50 days before the start of the experiment. Animals were housed in pairs in a temperature-controlled colony room on a 12 h light/dark cycle (lights on at 7:00 A.M.), and food and water were available *ad libitum*. Animals were allowed to acclimatize to the colony room for 3–5 d after their arrival before they were divided into 2 treatment groups: Animals in the control group received daily intraperitoneal (IP) injections of saline for 7 days while animals in the cocaine group received daily injections of cocaine hydrochloride in saline (Sigma) using a dosing regimen previously described to produce locomotor sensitization (Pierce, *et al* 1996; Bowers, *et al* 2004). On the first and last days of cocaine administration, the rats were injected with 15 mg/kg cocaine IP. On the intervening days, the animals were injected with 30 mg/kg cocaine IP. Both groups of animals were subjected to a 2 week withdrawal following the last injection of saline or cocaine. On the first day of drug administration and after the withdrawal period the motor activity of animals in both groups was monitored in a photocell apparatus (Omnitech, Columbus, OH, USA) as described elsewhere (Pierce, *et al* 1996). Briefly, animals were placed in the photocell cages for 1 hr after which time they were removed from the cages, injected with cocaine or saline, respectively, and returned to the photocell cages for an additional 2 hrs. Consistent with previous studies (Pierce, *et al* 1996) the cocaine treatment produced locomotor sensitivity. The total distance traveled during the first hour after cocaine injection (15 mg/kg) was significantly increased after 2 weeks of withdrawal when compared with activity on the first day of cocaine treatment (day 1: 1500.7 ± 260.0 cm; day 23: 3523.3 ± 741.3 cm; paired Student's-test, $P < 0.01$). Animals were sacrificed the following day for the preparation of brain slices. The experimenter was not blind to the treatment condition of the animal on the day of the electrophysiological recordings.

Brain slice preparation and electrophysiology

Saline or cocaine-treated rats (72–100 days old) were deeply anesthetized with an IP injection of chloral hydrate (400 mg/kg) and transcardially perfused with an ice-cold high sucrose substituted ACSF (in mM): 230 sucrose, 1.9 KCl, 1.2 Na₂HPO₄, 33 NaHCO₃, 6 MgCl₂, 0.5 CaCl₂, 10 glucose, and 0.4 ascorbic acid. Coronal slices (350 μm) were cut on a vibratome (Leica, VT1000S or VT1200S, Nussloch, Germany). Slices were incubated at room temperature for at least 1 h in a solution consisting of (in mM) 125 NaCl, 2.5 KCl, 1.25 Na₂HPO₄, 10 glucose, 25 NaHCO₃, 6 MgCl₂, and 1 CaCl₂. Finally, slices were

transferred to a submersion-type recording chamber and recordings were performed at 33 ± 1 °C using a recording ACSF consisting of (in mM): 125 NaCl, 2.5 KCl, 25 NaHCO₃, 1.3 MgCl₂, 2 CaCl₂, 10 glucose and 0.4 ascorbic acid at a rate of 1–2 ml/min. All ACSF solutions were constantly aerated with a mixture of 95% O₂–5% CO₂ to maintain ~7.2 pH. Whole-cell recordings were obtained from neurons in layers 5 and 6 of the prelimbic cortex (between 3.7 and 2 mm from bregma) using differential interference contrast optics (Axioskop, Zeiss; Germany) and videomicroscopy (Dage MTL, Michigan City, IND). For voltage-clamp recordings electrodes (3–5 MOhm open tip resistance) were filled with (in mM): 130 CsCl, 20 TEA, 10 HEPES, 2 MgCl₂, 0.5 EGTA, 4 Na-ATP, 0.3 Na-GTP, 14 phosphocreatine, and 2 QX-314. For current-clamp recordings, electrodes were filled with a solution containing (in mM): 125 K-gluconate, 10 HEPES, 20 KCl, 4 ATP-Mg, 0.3 GTP-Na, 14 phosphocreatine and 0.4 Alexa 594.

Data Collection and Analysis

Recordings were obtained with a Multiclamp 700B amplifier (Molecular Devices, Mississauga, CAN). Membrane potential was not corrected for changes in junction potential after break-in. Signals were low-pass filtered at 3 kHz and digitized at 10 kHz. Data were stored on PC for off-line analysis. Data acquisition and analysis were performed using software written in LabView (National Instruments, Austin, TX) or Axograph-X (J. Clemens, AUS). In voltage-clamp experiments we examined both evoked and spontaneous inhibitory post-synaptic currents (IPSCs). The membrane potential was held at –80 mV and GABA-mediated events were pharmacologically isolated by adding 20 μM 6-cyano-7-nitroquinoxaline-2,3-dione (CNQX, to block AMPA receptors) and 50 μM (±)2-amino-5-phosphonopentanoic acid (APV, to block NMDA receptors) to the bath. Action-potential independent miniature IPSCs (mIPSCs) were recorded under the same conditions but in the presence of 1 μM tetrodotoxin (TTX; Sigma) to block sodium channels. Synaptic stimulation of evoked IPSCs (eIPSCs) occurred every 30 seconds via a bipolar stimulation electrode positioned within 200 μm of the soma. Under baseline conditions the input-output curves for evoked IPSCs did not differ between the two experimental groups (i.e. cocaine- or saline-injected rats) (Figure S1). In current clamp recordings series of hyperpolarizing and depolarizing current steps (500 ms duration; 10–30 pA increments at 0.3 Hz) were delivered from resting membrane potential to evoke subthreshold responses and spike firing. Intrinsic membrane properties and the evoked firing pattern were used to distinguish subtypes of GABAergic interneurons. In order to determine changes in neuronal excitability following pharmacological manipulations the series of current pulses was repeated every 3 minutes for the duration of the experiment. Comparisons of changes in the number of evoked spikes were made at a current level that reliably produced repetitive firing under control conditions. In both voltage-clamp and current clamp recordings access resistance was continuously monitored by applying small hyperpolarizing or depolarizing current steps through the recording pipette at the beginning of the sweep and cells that changed more than 20% during the recording were excluded from analysis. The morphology of non-pyramidal cells was verified at the end of the experiment via confocal imaging of Alexa 594. After collection of baseline data, DA, or specific DA receptor agonists were bath-applied for 2–3 minutes. We used the agonist (±)-Quinpirole dihydrochloride to activate D2-like receptors and (±)-6-chloro-PB hydrobromide (SKF-81297) to activate D1-like receptors. When the

effects of DA antagonists were examined, the D1 antagonist R-[+]-7-chloro-8-hydroxy-3-methyl-1-phenyl-2,3,4,5-tetrahydro-1H-3-benzazepine (SCH23390), or the D2 antagonist (\pm)-Sulpiride were bath-applied at least 10 minutes before application of DA and continued to be present throughout the remainder of the experiment. All DAergic agents were obtained from Sigma (St. Louis, MO). The frequency and amplitude of IPSCs were measured using MiniAnalysis (Synaptosoft, Decatur, GA). Spontaneous or miniature events were detected when they crossed a threshold set at two times the RMS baseline noise. The detected events were confirmed as synaptic events by eye. For sIPSCs and mIPSCs we collected 9 second long sweeps every 30 seconds. Two sweeps (representing one minute) were averaged and changes in the amplitude of eIPSCs or the frequency of sIPSCs, respectively, were plotted over time. For statistical comparisons the number of evoked responses or spontaneous events was averaged across two separate 5 minute intervals as indicated in the Results section and compared to baseline values. For statistical comparisons in current clamp recordings at least 3–5 responses under baseline and drug condition were averaged and compared. Comparisons were performed using two-tailed, paired *t*-tests unless indicated otherwise. Differences of $\alpha = 0.05$ were considered statistically significant. All data are presented as means \pm SEM.

Results

Dopamine has a temporally biphasic effect on the amplitude of eIPSCs in saline-treated animals. Figure 1A shows the effect of a high dose (20 μ M) of dopamine on isolated eIPSCs in pyramidal neurons. On average, dopamine modulated eIPSC amplitude in a biphasic, time-dependent manner, producing an initial decrease (2–20 min after application), followed by a prolonged increase in the amplitude of the eIPSC. In order to quantify this effect we averaged the responses over 5 minute periods either briefly after DA bath application (“early DA”, between 5–10 minutes after DA was first perfused into the bath) and at a later time point when the increase in eIPSC amplitude had reached a plateau (“late DA”, between 25–30 minutes after DA application). In this and all following experiments the averaged responses from these time periods were then compared with the averaged response of the baseline period prior to drug application. As seen in Figure 1A, in saline-treated rats 20 μ M DA reduced the amplitude of the eIPSC during the “early DA” period by about -24.4 ± 14.2 % ($p < 0.01$; $n = 8$) and increased the eIPSC amplitude during the “late DA” period on average by $+64.2 \pm 29.7$ % ($p < 0.05$). In contrast, in cocaine-treated animals the initial reduction in eIPSC amplitude was absent and instead an immediate increase of eIPSC amplitude was observed during the early phase of the DA effect ($+30.1 \pm 10.6$ %; $p < 0.05$) that persisted through the late phase of the DA effect ($+51.1 \pm 23.6$ %; $p < 0.05$; $n = 7$; Figure 1B).

Receptor-specific modulation of evoked IPSCS

The biphasic modulation of eIPSCs in our adult saline-treated animals is consistent with previous observations in young drug-naïve animals. In these studies concentrations of extracellular DA high enough to activate both receptor subtypes in the PFC ($> 1 \mu$ M DA) lead to a long-lasting enhancement of eIPSCs via activation of D1Rs, whereas the initial reduction in eIPSC amplitude required activation of D2Rs (Seamans, *et al* 2001; Trantham-Davidson, *et al* 2004). Accordingly, we found that in saline-treated animals bath application

of the D2R agonist quinpirole (5 μ M, n=6) reduced the amplitude of the eIPSC during the period of the early measurement (5–10 min after drug application), which partly recovered by the time of the second late measurement (25–30 min after drug application) (quinpirole early -37.7 ± 16.4 %, $p < 0.01$, quinpirole late -23.4 ± 24.7 %; $p < 0.01$; n=6, Figure 2A). Similar results were also observed when DA (20 μ M) was applied in the presence of the D1 antagonist SCH-23390 (10 μ M, n=3; data not shown). In contrast, in cocaine-sensitized animals bath application of the selective D2 agonist quinpirole (5 μ M) failed to induce a reduction in eIPSC amplitude (Figure 2B). Quinpirole did not significantly affect eIPSCs at either the early or late time point (quinpirole early $+0.7 \pm 10$ %; quinpirole late -12.3 ± 8 %; n=11). On the other hand, D1R-dependent signaling appeared to be unaffected in cocaine-sensitized animals, resulting in increased eIPSC amplitudes. Using different approaches to stimulate D1Rs this effect consistently became significant by the time of the second late measurement (Figure 2C–E): Bath application of the selective D1R agonist SKF-81297 (5 μ M; n=6) increased eIPSC amplitude by 31.4 ± 22 % ($p > 0.05$) during the period of the early measurement, and by 85.7 ± 30 % ($p < 0.05$) during the period of the late measurement. Similarly, application of 20 μ M DA in the presence of the selective D2R antagonist sulpiride (5 μ M) increased eIPSC amplitudes by 27.9 ± 18 % ($p > 0.05$) during the early measurement and by 26.0 ± 5 % ($p < 0.01$; n=11) during the period of the late measurement, respectively. Finally, a low dose (100 nM) of DA by itself, which previously had been shown to predominantly activate D1Rs under similar recording conditions in drug-naïve animals (Trantham-Davidson, *et al* 2004), also increased eIPSC amplitudes in cocaine-sensitized rats (Figure 2E; early DA $+20.9 \pm 12.6$, $p > 0.05$; late DA $+33.7 \pm 11.3$, $p = 0.5$; n=8). Taken together these results suggest that in cocaine-sensitized animals D2 mediated modulation of IPSCs is compromised but D1R-dependent mechanisms are unaffected.

Dopamine modulation of spontaneous IPSCs

In a first effort to assess which components of synaptic transmission are altered in cocaine-sensitized animals we analyzed the frequency and amplitude distribution of spontaneous IPSCs (sIPSCs) in saline and cocaine-treated animals (Figure 3). Our results show that in slices from both groups of animals bath application of a high dose of DA (20 μ M) led to a significant increase in the frequency of sIPSCs. The development of this effect over time was noticeably different in saline- and cocaine-treated animals. Figure 3B and 3C, show the time-course of the change in sIPSC frequency and the relative changes in the amplitude distribution of sIPSCs between 5–10 minutes and 25–30 minutes after DA bath application, respectively, compared with baseline measures. In slices from saline-treated animals the frequency of sIPSCs gradually rose and reached a plateau value after about 20 min (Figure 3B; early DA $+15.4 \pm 6$ %, $p > 0.05$; late DA $+29.0 \pm 12$ %, $p < 0.05$; n=8). We plotted changes in the frequency distribution of sIPSC amplitudes to see whether DA might selectively modulate sIPSCs of a certain size (Figure 3B, right column). We found that DA consistently increased the frequency of events of all amplitudes, with the exception of very small events (< 10 pA) that were decreased during the period of the early DA effect. However, across all events within a measurement period the DA-induced increase in sIPSC frequency had no effect on mean sIPSC amplitude (Figure 3B inset). In cocaine-sensitized animals (Figure 3C) the frequency of sIPSCs increased rapidly after DA application and remained significantly elevated throughout the time of the recordings (early DA $+44.7 \pm 13$ %,

$p=0.05$; late DA $+24.9 \pm 7\%$, $p<0.05$; $n=6$). As in saline animals the mean amplitude of sIPSCs did not significantly change during any of the three measurement periods (Figure 3C inset), but unlike in slices from saline animals there occurred no reduction in small amplitude events during the early measurement of DA effects in cocaine-treated animals. Instead, there was a disproportional increase in the number of large amplitude (>70 pA) events (Figure 3, right column). One potential mechanism for the differential DA modulation of small and large amplitude IPSCs, respectively, in saline- and cocaine-treated animals was explored in a series of action potential-independent miniature IPSC recordings (see below).

The increase in sIPSC frequency seen in cocaine-sensitized animals in response to $20 \mu\text{M}$ DA depended on D1R activation, as similar increases were seen in response to SKF-81297 ($5 \mu\text{M}$; $n=6$). SKF-81297 increased the frequency of sIPSCs by $23.1 \pm 4\%$ ($p<0.01$; $n=6$) during the early measurement and by $36.0 \pm 11\%$ ($p<0.05$; $n=6$) during the period of the late measurement, respectively (Figure 4C,D), and this again was mimicked by the application of a low dose of DA (Figure 4D). In contrast, D2-receptor activation with $5 \mu\text{M}$ quinpirole (Figure 4A,B,D) decreased the number of sIPSCs in saline-treated animals (early quinpirole $-15.6 \pm 7\%$, $p>0.05$; late quinpirole $+32.7 \pm 7\%$, $p<0.05$, $n=8$), but had no consistent effect in slices from cocaine-treated animals (early quinpirole $+7.0 \pm 8\%$, $p>0.05$; late quinpirole $-9.9 \pm 22\%$, $p>0.05$; $n=6$).

Dopamine and D2 receptor-dependent modulation of miniature IPSCs

Spontaneous IPSCs represent both action potential-dependent and -independent release of GABA. In contrast, miniature IPSCs (mIPSCs) are recorded in the presence of TTX ($1 \mu\text{M}$) to eliminate the contribution of action potential-mediated release events. Changes in GABAergic release could also contribute to the D2R-dependent reduction in eIPSC amplitude and the lack thereof in cocaine-sensitized animals. We therefore studied the effects of DA and D2R stimulation on mIPSCs in order to determine whether the increase in the frequency of sIPSCs was due to a selective effect on interneuron excitability and spike firing, or reflected at least in part a modification of GABA release as measured with mIPSCs. Miniature IPSCs were recorded in the presence of TTX ($1 \mu\text{M}$) and changes in the frequency of mIPSCs were compared between cocaine and saline-treated animals (Figure 5). In saline animals quinpirole ($5 \mu\text{M}$) significantly reduced the number of mIPSCs 5–10 minutes after drug application (Figure 5A; $-13.3 \pm 6\%$; $p<0.05$; $n=7$). DA D2 receptor activation not only reduced the frequency but also the average amplitude of mIPSCs, suggesting the presence of an additional postsynaptic mechanism through which DA can modulate inhibitory synaptic currents (Seamans et al., 2001). Quinpirole-induced changes in mIPSC frequency and amplitude occurred independent of changes in access resistance.

In cocaine-sensitized rats a similar change in the number of events was not observed at any time of the recording (Figure 5B; $-0.7 \pm 9\%$; $p>0.05$; $n=7$). Bath application of the full agonist DA ($20 \mu\text{M}$) led to a significant increase in the frequency of mIPSCs in slices from cocaine-sensitized animals (Figure 5D; baseline 4.7 ± 0.4 events/s, DA early 6.3 ± 0.5 events/s; $p<0.01$; DA late 6.7 ± 1.0 events/s $p<0.05$; $n=7$), but only a moderate increase in slices from saline-treated animals (Figure 5C; baseline 5.8 ± 0.4 events/s, DA early 6.2 ± 1.7

events/s; $p > 0.05$ early; DA late 6.2 ± 1.4 events/s $p > 0.05$; $n = 4$). This DA-induced increase in the frequency of mIPSCs, together with the absence of an effect of quinpirole on sIPSCs, suggests that a D1R-dependent increase in GABA release contributes to the rapid increase in the number of sIPSCs described above (c.f. Figure 3), but that in normal animals this is counteracted by a D2-mediated reduction in release. Thus taken together, the data from sIPSC and mIPSC recordings suggest that in control animals opposing D1/D2 mechanisms determine the direction and time course of DA modulation of inhibition, and that both a D2R-dependent reduction in GABA release and the desensitization of postsynaptic GABA receptors could contribute to the reduction of eIPSCs. In contrast, in cocaine-sensitized animals the efficacy of D2R stimulation appears to be greatly reduced, resulting in a change in the time-course of DA modulation of IPSCs and enhanced inhibition via D1-dependent mechanisms, most likely due to parallel increases in the frequency of action potential-dependent and -independent synaptic events.

Cocaine-sensitization reduces D2 receptor-dependent modulation of action-potential firing in PFC interneurons

Recently, it was shown that in PFC interneurons from adult drug-naïve animals D2R activation can increase action potential firing evoked by somatic current injection (Tseng and O'Donnell, 2007). Therefore, we examined whether withdrawal from cocaine can potentially also affect cortical inhibition via D2-dependent modulation of spike firing in interneurons. To this end we obtained current clamp recordings from identified non-pyramidal neurons in the PFC (Figure 6A) and we examined changes in membrane excitability in response to D2R activation with quinpirole. Based on their firing pattern, afterhyperpolarization (AHP), and spike frequency adaptation characteristics (Cauli, *et al* 1997; Kroener, *et al* 2007), cells were classified either as fast-spiking interneurons (FS; $n = 14$) or non-fast spiking cells (NFS; $n = 12$) (Figure 6). Fast-spiking cells were characterized by action potentials of short duration, which were followed by fast, monophasic AHPs of large amplitude (table 1). Fast-spiking cells were able to sustain high steady-state firing frequencies with relatively little spike-frequency adaptation (Figure 5B). Non-fast spiking cells displayed conspicuous spike frequency adaptation (Figure 6B), and their spike kinetics and AHP amplitude were significantly different from those of FS cells both in tissue from saline-treated and cocaine treated animals (table 1). No differences in basic membrane properties were observed when comparing cells belonging to the same class in saline-treated or cocaine-treated rats. However, both FS and NFS cells showed similar differences in their response to quinpirole ($5 \mu\text{M}$) when comparing cells in slices from saline-treated and cocaine-treated animals (Figure 6C,D): In saline animals quinpirole produced a small but consistent increase in the number of current-evoked spikes in both FS cells (baseline 8.2 ± 1.4 ; quinpirole 11.0 ± 1.5 ; $p < 0.05$) as well as NFS cells (baseline 6.5 ± 0.38 ; quinpirole 7.6 ± 0.32 ; $p < 0.05$). In contrast, in cocaine animals quinpirole had no consistent effect on membrane excitability or the number of evoked spikes in either FS or NFS cells (Figure 6C, D). Taken together our results suggest that in cocaine-sensitized animals the ability of D2R stimulation to modulate the activity of interneurons and to regulate GABAergic transmission onto pyramidal cells is greatly reduced. In contrast, D1R-dependent mechanisms, which can increase the amplitude of evoked IPSCs and the frequency of spontaneous IPSCs are not affected.

Discussion

We show that repeated non-contingent exposure and withdrawal from cocaine alter DA modulation of inhibition in the PFC. In control animals DA and D2R stimulation result in a transient reduction of eIPSCs, which is absent in cocaine-sensitized animals. These changes in eIPSCs are paralleled by a lack of D2R modulation of GABAergic release as measured with mIPSCs, and they manifest in changes in the time course of DA modulation of sIPSCs. In addition, D2R-mediated increases in current-evoked spike firing in interneurons from saline-treated animals were absent in cocaine-sensitized animals. Therefore, the time- and concentration-dependent interplay of D1Rs and D2Rs that characterizes DA modulation in control animals is disrupted in sensitized animals, resulting in an unopposed D1R-mediated increase in GABAergic inhibition. This loss of synergism may alter normal PFC network processing, thus contributing to the functional hypofrontality seen in cocaine addicts.

Dopamine-GABA interactions in the PFC as targets for cocaine-induced neuroadaptations

Numerous studies have shown the importance of prefrontal cortical DA and GABA neurotransmission in the development or reinstatement of addiction to cocaine. Injections of DA receptor antagonists into the PFC prevent, while agonists can induce reinstatement, and DA release in the PFC is increased during reinstatement (Sorg, *et al* 1997; McFarland and Kalivas, 2001; McFarland, *et al* 2004; Sun and Rebec, 2005; Williams and Steketee, 2005). Repeated exposure to cocaine decreases D2R function in the PFC, leading to persistent cognitive deficits (Briand, *et al* 2008). Injections of quinpirole into the mPFC block sensitization (Beyer and Steketee, 2002), while injections of a D2 antagonist enhance motor activity (Steketee and Walsh, 2005). Moreover, non-contingent cocaine administration produces a persistent decrease in D2R-dependent regulation of cortical excitability (Nogueira, *et al* 2006). GABA in the PFC is released from local interneurons and projections from the VTA (Carr and Sesack, 2000). GABAergic interneurons are a prominent target of cortical DA innervation (Sesack, *et al* 1995) and they possess both D1- and D2-type DA receptors (LeMoine and Gaspar 1998; Santana, *et al* 2009). Local infusion of the GABA_B agonist baclofen blocks the initiation of cocaine-induced behavioral sensitization (Steketee and Beyer, 2005). Once sensitization is induced, the levels of extracellular GABA in the PFC are increased (Jayaram and Steketee, 2005, 2006), and cue-induced drug-seeking selectively increases the expression of c-Fos in interneurons of the prelimbic cortex, while simultaneously reducing activity in pyramidal cells (Miller and Marshall, 2004).

Mechanisms of DA modulation of inhibition in control- and cocaine-sensitized animals

Reports on the effects of DA on GABAergic transmission in the PFC of drug-naïve animals paint a complex and often contradictory picture, and both a reduction (Abekawa, *et al* 2000; Gonzalez-Islas and Hablitz 2001; Seamans, *et al* 2001; Gao, *et al* 2003; Trantham-Davidson, *et al* 2004) and an increase (Retaux, *et al* 1991; Grobin and Deutch, 1998; Zhou and Hablitz 1999; Gullledge and Jaffe, 2001; Seamans, *et al* 2001; Gao, *et al* 2003; Trantham-Davidson, *et al* 2004; Kroener, *et al* 2007) in either the amplitude of eIPSCs, the frequency of sIPSCs, or the extracellular levels of GABA as measured with microdialysis have been described, hampering the interpretation of changes seen in cocaine-sensitized animals.

We found that in adult saline-treated animals DA had time-dependent effects on synaptically evoked IPSCs that were specific for D1R and D2R activation, respectively, replicating previous results in young drug-naïve rats (Seamans, *et al* 2001; Trantham-Davidson, *et al* 2004). Under normal conditions high DA (>1 μ M) activates both D1Rs and D2Rs, but the initial D2R-mediated reduction in eIPSC amplitude occludes the prolonged D1R-mediated enhancement (Figure 1; Seamans, *et al* 2001; Trantham-Davidson, *et al* 2004). The D2R-dependent reduction in eIPSC amplitude in saline-treated animals could result from alterations in GABA release at the presynaptic terminal as indicated by a reduction in the frequency of mIPSCs. Surprisingly, in saline-treated animals a high dose of DA did not result in a similar reduction in the frequency of mIPSCs, thus leaving the need for an additional mechanism to explain the transient reduction of eIPSCs in control animals. In young drug naïve animals D2R activation has previously also been shown to decrease the response of postsynaptic GABA_A receptors on pyramidal neurons (Seamans, *et al* 2001) thus working to further diminish the amplitude of eIPSCs. Here we also demonstrate data that is consistent with this idea as we show that in saline-treated animals not only the frequency, but also the amplitude of mIPSCs was reduced by application of the D2 agonist. In contrast this effect did also not occur in cocaine-treated animals. Thus under normal conditions D2 receptor activation appears to reduce both release of GABA via a presynaptic mechanism and may reduce the sensitivity of postsynaptic GABA receptors to decrease the amplitude of the IPSC. On the other hand, direct D1R activation (either via DA in the presence of a D2 antagonist or through a specific D1 agonist) or low extracellular concentrations of DA increased the amplitude of eIPSCs and the frequency of sIPSCs consistent with previous reports in drug-naïve animals (Zhou and Hablitz 1999; Gulledege and Jaffe, 2001; Seamans, *et al* 2001; Trantham-Davidson, *et al* 2004; Kroener, *et al* 2007). Because D1R activation also increases the membrane excitability of fast-spiking interneurons these changes reflect at least in part a presynaptic modulation of excitability and spontaneous action-potential firing (Gorelova, *et al* 2002; Kroener, *et al* 2007; Tseng and O'Donnell, 2007).

Repeated cocaine exposure reduced all measures of D2R activity tested here, shifting the time course of DA-induced changes and generally resulting in increases in inhibition. In cells from cocaine-sensitized animals eIPSCs showed an immediate increase in amplitude, which was paralleled by an increase in the frequency of sIPSCs and mIPSCs following application of DA or a D1 agonist. The D2R-specific reduction in mIPSC frequency was not observed in cocaine-sensitized animals. In acute brain slices action potential-independent “miniature” release represents a significant portion of the total number of spontaneous synaptic events (Simkus and Stricker, 2002; Kroener, *et al* 2007; present data). The lack of D2R modulation of mIPSCs may thus explain the rapid increase in sIPSCs in cocaine animals. This interpretation is further supported by the observation that in cocaine sensitized animals the full agonist DA increased the frequency of mIPSCs, and the fact that during the early DA effect the amplitude distribution of sIPSCs was shifted to the left, indicating a relative greater contribution of small amplitude mIPSCs to the overall number of synaptic events. Dopamine D1Rs in the terminals of interneurons provide an anatomical substrate for presynaptic modulation of GABA release by DA (Gulledege and Jaffe, 2001; Paspalas and Goldman-Rakic, 2005; Muly, *et al* 1998). Moreover, the D1R-dependent changes in

membrane excitability and action potential-dependent release discussed above likely also contributed to the increase in the frequency of sIPSCs and the increased amplitude of eIPSCs. The reduction in GABA release via D2Rs is functionally similar to stimulation of presynaptic GABA_B autoreceptors with baclofen, which prevents the initiation of sensitization to cocaine (Steketee and Beyer, 2005). Reductions in the effectiveness of D2R or GABA_B receptor signaling in the PFC of sensitized animals may result from elevated levels of AGS3 and these alterations have been shown to persist though up to 8 weeks of withdrawal (Bowers, *et al* 2004; Frankowska, *et al* 2008). AGS3 is a member of a family of G-protein regulators, which sequesters G α and thereby limits signaling via G α -coupled receptors. Finally, D2R activation increased the number of current-evoked spikes in interneurons from saline-treated, but not from cocaine-sensitized animals, providing further evidence for a generalized reduction in Gi-receptor signaling in cocaine-sensitized animals. However, the results in control animals also indicate an obvious paradox in that it appears that D2R activation can differentially modulate action potential-mediated release of GABA with opposing results (Retaux, *et al* 1991). D2R activation decreases the amplitude of eIPSCs (Figure 1; Seamans, *et al* 2001; Trantham-Davidson, *et al* 2004), yet it can also increase current-evoked firing in interneurons (Figure 6; Tseng and O'Donnell, 2007; Tseng, *et al* 2008) and DA release in vivo (Grobin and Deutch, 1998). We replicate two of these findings in adult (>72d) saline-treated rats, thus ruling out that these discrepancies solely reflect differences in the developmental state of animals used in previous studies. One potential caveat is that the response to somatic current injection may not adequately reflect the mechanisms that govern spike firing resulting from integration of synaptic inputs. We have recently shown that DA can decrease cortical network activity, even though it consistently increased somatic current-evoked spike firing in individual pyramidal cells (Kroener, *et al* 2009). Similarly, our results from mIPSC recordings may not fully reflect the effects of DA and D2 receptor activation on synaptic integration, and furthermore these effects may differ from the postsynaptic effects of DA that regulate spike firing. Thus clearly more research is needed to unveil the mechanisms through which D2Rs regulate inhibition in the cortex under normal conditions.

Functional implications of altered DA modulation of inhibition

Drug addiction has been conceptualized as a syndrome of impaired response inhibition and salience attribution (Goldstein and Volkow, 2002). Prefrontal control over top-down processes is reduced in addiction (Bolla, *et al* 2004; Briand, *et al* 2008). Behavioral data indicate that activating the DAergic projection from the VTA to the PFC initiates reinstatement via a glutamatergic projection to the nucleus accumbens core (McFarland *et al* 2004). On the other hand blocking all PFC activity through injections of GABA agonists reduces drug seeking (McFarland and Kalivas, 2001; McLaughlin and See, 2003; McFarland *et al* 2004). Our data using non-contingent delivery of cocaine indicate that changes in the dopaminergic modulation of inhibition contribute to the behavioral sensitization following withdrawal from cocaine. The most parsimonious interpretation of our data is that addiction to cocaine changes the balance between D1 and D2 receptor contribution in DA modulation of inhibition. It has been hypothesized that preferential activation of D1Rs and D2Rs, respectively, correlates with different PFC activity states that determine the ability of competing motivational representations to drive PFC output to the nucleus accumbens

(Seamans, *et al* 2001; Kalivas, *et al* 2005; Durstewitz and Seamans, 2008). Interneurons that receive DAergic innervation are well positioned to mediate a switch between different network states (Lapish, *et al* 2007). At the circuit level, interneurons gate inputs to a network and pattern its outputs (McBain and Fisahn, 2001; Compte, *et al* 2000; Lapish, *et al* 2007). Dopamine increases the temporal precision of interneuron firing in the PFC (Tierney, *et al* 2008), and thus cocaine-induced alterations in DAergic control will disrupt optimal signal processing in the PFC (Kroener *et al.*, 2009). If D2 regulation of PFC activity is disturbed as a result of repeated cocaine exposure the shift towards D1R signaling may result in attenuated PFC output to subcortical targets (Franklin and Druhan, 2000) and/or a breakdown in optimal processing of salient stimuli that guide behavior (Durstewitz and Seamans, 2008), which can be characterized as hypofrontality and reduced motivation to respond to non-drug-related stimuli. Under these conditions only very strong motivational stimuli (e.g., drug-associated cues) will prevail in activating PFC networks and initiating goal-directed behavior towards this limited set of stimuli (Kalivas, *et al* 2005). Our data provide physiological evidence for this hypothesized shift in network activation.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

This research was supported by grants from the National Institutes of Health (DA 014698 to AL; and R21NS056124-02 and P50DA015369-04 subcomponent 84875 to SK). The authors wish to thank Dr. Glen Andrews for support during preliminary experiments.

References

- Abekawa T, Ohmori T, Ito K, Koyama T. D1 dopamine receptor activation reduces extracellular glutamate and GABA concentrations in the medial prefrontal cortex. *Brain Res.* 2000; 867:250–254. [PubMed: 10837822]
- Beyer CE, Steketee JD. Cocaine sensitization: modulation by dopamine D2 receptors. *Cereb Cortex.* 2002; 12:526–535. [PubMed: 11950770]
- Bolla K, Ernst M, Kiehl K, Mouratidis M, Eldreth D, Contoreggi C, *et al.* Prefrontal cortical dysfunction in abstinent cocaine abusers. *J Neuropsychiatry Clin Neurosci.* 2004; 16:456–464. [PubMed: 15616172]
- Bowers MS, McFarland K, Lake RW, Peterson YK, Lapish CC, Gregory ML, *et al.* Activator of G protein signaling 3: a gatekeeper of cocaine sensitization and drug seeking. *Neuron.* 2004; 42:269–281. [PubMed: 15091342]
- Briand LA, Flagel SB, Garcia-Fuster MJ, Watson SJ, Akil H, Sarter M, *et al.* Persistent alterations in cognitive function and prefrontal dopamine D2 receptors following extended, but not limited, access to self-administered cocaine. *Neuropsychopharmacology.* 2008; 33:2969–2980. [PubMed: 18305460]
- Brozoski TJ, Brown RM, Rosvold HE, Goldman PS. Cognitive deficit caused by regional depletion of dopamine in prefrontal cortex of rhesus monkey. *Science.* 1979; 205:929–932. [PubMed: 112679]
- Carr DB, Sesack SR. GABA-containing neurons in the rat ventral tegmental area project to the prefrontal cortex. *Synapse.* 2000; 38:114–123. [PubMed: 11018785]
- Cauli B, Audinat E, Lambollez B, Angulo MC, Ropert N, Tsuzuki K, *et al.* Molecular and physiological diversity of cortical nonpyramidal cells. *J Neurosci.* 1997; 17:3894–3906. [PubMed: 9133407]

- Compte A, Brunel N, Goldman-Rakic PS, Wang XJ. Synaptic mechanisms and network dynamics underlying spatial working memory in a cortical network model. *Cereb. Cortex.* 2000; 10:910–923. [PubMed: 10982751]
- Durstewitz D, Seamans JK. The dual-state theory of prefrontal cortex dopamine function with relevance to catechol-o-methyltransferase genotypes and schizophrenia. *Biol Psychiatry.* 2008; 64:739–749. [PubMed: 18620336]
- Franklin TR, Druhan JP. Involvement of the nucleus accumbens and medial prefrontal cortex in the expression of conditioned hyperactivity to a cocaine-associated environment in rats. *Neuropsychopharmacology.* 2000; 23:633–644. [PubMed: 11063919]
- Frankowska M, Wydra K, Faron-Górecka A, Zaniewska M, Ku mider M, Dziedzicka-Wasylewska M, et al. Alterations in gamma-aminobutyric acid(B) receptor binding in the rat brain after reinstatement of cocaine-seeking behavior. *Pharmacol Rep.* 2008; 60:834–843. [PubMed: 19211975]
- Gao WJ, Wang Y, Goldman-Rakic PS. Dopamine modulation of perisomatic and peridendritic inhibition in prefrontal cortex. *J Neurosci.* 2003; 23:1622–1630. [PubMed: 12629166]
- Goldstein RZ, Volkow ND. Drug addiction and its underlying neurobiological basis: neuroimaging evidence for the involvement of the frontal cortex. *Am J Psychiatry.* 2002; 159:1642–1652. [PubMed: 12359667]
- Gonzalez-Islas C, Hablitz JJ. Dopamine inhibition of evoked IPSCs in rat prefrontal cortex. *J Neurophysiol.* 2001; 86:2911–2918. [PubMed: 11731547]
- Gorelova N, Seamans JK, Yang CR. Mechanisms of dopamine activation of fast-spiking interneurons that exert inhibition in rat prefrontal cortex. *J Neurophysiol.* 2002; 88:3150–3166. [PubMed: 12466437]
- Grobin AC, Deutch AY. Dopaminergic regulation of extracellular gamma-aminobutyric acid levels in the prefrontal cortex of the rat. *J Pharmacol Exp Ther.* 1998; 285:350–357. [PubMed: 9536031]
- Gulledge AT, Jaffe DB. Multiple effects of dopamine on layer V pyramidal cell excitability in rat prefrontal cortex. *J Neurophysiol.* 2001; 86:586–595. [PubMed: 11495934]
- Jayaram P, Steketeer JD. Effects of cocaine-induced behavioural sensitization on GABA transmission within rat medial prefrontal cortex. *Eur J Neurosci.* 2005; 21:2035–2039. [PubMed: 15869498]
- Jayaram P, Steketeer JD. Cocaine-induced increases in medial prefrontal cortical GABA transmission involves glutamatergic receptors. *Eur J Pharmacol.* 2006; 531:74–79. [PubMed: 16409999]
- Kalivas PW, Volkow N, Seamans JK. Unmanageable motivation in addiction: a pathology in prefrontal-accumbens glutamate transmission. *Neuron.* 2005; 45:647–650. [PubMed: 15748840]
- Kroener S, Phillips PEM, Chandler LJ, Seamans JK. Dopamine modulates recurrent synaptic activity and enhances the signal-to-noise ratio in the prefrontal cortex. *PLoS One.* 2009; 4:e6507. [PubMed: 19654866]
- Kroener S, Krimer LS, Lewis DA, Barrionuevo G. Dopamine increases inhibition in the monkey dorsolateral prefrontal cortex through cell type-specific modulation of interneurons. *Cereb Cortex.* 2007; 17:1020–1032. [PubMed: 16772311]
- Lapish CC, Kroener S, Durstewitz D, Lavin A, Seamans JK. The ability of the mesocortical dopamine system to operate in distinct temporal modes. *Psychopharmacology.* 2007; 191:609–625. [PubMed: 17086392]
- LeMoine C, Gaspar P. Subpopulations of cortical GABAergic interneurons differ by their expression of D1 and D2 dopamine receptor subtypes. *Brain Res Mol Brain Res.* 1998; 58:231–236. [PubMed: 9685656]
- McBain CJ, Fisahn A. Interneurons unbound. *Nat Rev Neurosci.* 2001; 2:11–23. [PubMed: 11253355]
- McFarland K, Davidge SB, Lapish CC, Kalivas PW. Limbic and motor circuitry underlying footshock-induced reinstatement of cocaine-seeking behavior. *J Neurosci.* 2004; 24:1551–1560. [PubMed: 14973230]
- McFarland K, Kalivas PW. The circuitry mediating cocaine-induced reinstatement of drug-seeking behavior. *J Neurosci.* 2001; 21:8655–8663. [PubMed: 11606653]
- McLaughlin J, See RE. Selective inactivation of the dorsomedial prefrontal cortex and the basolateral amygdala attenuates conditioned-cued reinstatement of extinguished cocaine-seeking behavior in rats. *Psychopharmacology.* 2003; 168:57–65. [PubMed: 12845418]

- Miller CA, Marshall JF. Altered prelimbic cortex output during cue-elicited drug seeking. *J Neurosci.* 2004; 24:6889–6897. [PubMed: 15295023]
- Muly EC III, Szigeti K, Goldman-Rakic PS. D1 receptor in interneurons of macaque prefrontal cortex: distribution and subcellular localization. *J Neurosci.* 1998; 18:10553–10565. [PubMed: 9852592]
- Nogueira L, Kalivas PW, Lavin A. Long-term neuroadaptations produced by withdrawal from repeated cocaine treatment: role of dopaminergic receptors in modulating cortical excitability. *J Neurosci.* 2006; 26:12308–12313. [PubMed: 17122056]
- Paspalas CD, Goldman-Rakic PS. Presynaptic D1 dopamine receptors in primate prefrontal cortex: target-specific expression in the glutamatergic synapse. *J Neurosci.* 2005; 25:1260–1267. [PubMed: 15689564]
- Peters J, LaLumiere RT, Kalivas PW. Infralimbic prefrontal cortex is responsible for inhibiting cocaine seeking in extinguished rats. *J Neurosci.* 2008; 28:6046–6053. [PubMed: 18524910]
- Pierce RC, Bell K, Duffy P, Kalivas PW. Repeated cocaine augments excitatory amino acid transmission in the nucleus accumbens only in rats having developed behavioral sensitization. *J Neurosci.* 1996; 16:1550–1560. [PubMed: 8778304]
- Rétaux S, Besson MJ, Penit-Soria J. Opposing effects of dopamine D2 receptor stimulation on the spontaneous and the electrically evoked release of [3H]GABA on rat prefrontal cortex slices. *Neuroscience.* 1991; 42:61–71. [PubMed: 1677746]
- Santana N, Mengod G, Artigas F. Quantitative analysis of the expression of dopamine D1 and D2 receptors in pyramidal and GABAergic neurons of the rat prefrontal cortex. *Cereb Cortex.* 2009; 19:849–860. [PubMed: 18689859]
- Seamans JK, Gorelova N, Durstewitz D, Yang CR. Bidirectional dopamine modulation of GABAergic inhibition in prefrontal cortical pyramidal neurons. *J. Neurosci.* 2001; 21:3628–3638. [PubMed: 11331392]
- Sesack SR, Snyder CL, Lewis DA. Axon terminals immunolabeled for dopamine or tyrosine hydroxylase synapse on GABA-immunoreactive dendrites in rat and monkey cortex. *J Comp Neurol.* 1995; 363:264–280. [PubMed: 8642074]
- Simkus CR, Stricker C. Properties of mEPSCs recorded in layer II neurones of rat barrel cortex. *J Physiol.* 2002; 545:509–520. [PubMed: 12456830]
- Sorg BA, Davidson DL, Kalivas PW, Prasad BM. Repeated daily cocaine alters subsequent cocaine-induced increase of extracellular dopamine in the medial prefrontal cortex. *J Pharmacol Exp Ther.* 1997; 281:54–61. [PubMed: 9103480]
- Steketee JD. Neurotransmitter systems of the medial prefrontal cortex: potential role in sensitization to psychostimulants. *Brain Res Brain Res Rev.* 2003; 41:203–228. [PubMed: 12663081]
- Steketee JD, Beyer CE. Injections of baclofen into the ventral medial prefrontal cortex block the initiation, but not the expression, of cocaine sensitization in rats. *Psychopharmacology.* 2005; 180:352–358. [PubMed: 15696327]
- Steketee JD, Walsh TJ. Repeated injections of sulpiride into the medial prefrontal cortex induces sensitization to cocaine in rats. *Psychopharmacology.* 2005; 179:753–760. [PubMed: 15619114]
- Sun W, Rebec GV. The role of prefrontal cortex D1-like and D2-like receptors in cocaine-seeking behavior in rats. *Psychopharmacology.* 2005; 177:315–323. [PubMed: 15309375]
- Tierney PL, Thierry AM, Glowinski J, Deniau JM, Gioanni Y. Dopamine modulates temporal dynamics of feedforward inhibition in rat prefrontal cortex in vivo. *Cereb Cortex.* 2008; 18:2251–2262. [PubMed: 18222936]
- Trantham-Davidson H, Neely LC, Lavin A, Seamans JK. Mechanisms underlying differential D1 versus D2 dopamine receptor regulation of inhibition in prefrontal cortex. *J Neurosci.* 2004; 24:10652–10659. [PubMed: 15564581]
- Tseng KY, Lewis BL, Hashimoto T, Sesack SR, Kloc M, Lewis DA, et al. A neonatal ventral hippocampal lesion causes functional deficits in adult prefrontal cortical interneurons. *J Neurosci.* 2008; 28:12691–12699. [PubMed: 19036962]
- Tseng KY, O'Donnell P. Dopamine modulation of prefrontal cortical interneurons changes during adolescence. *Cereb Cortex.* 2007; 17:1235–1240. [PubMed: 16818475]

- Williams JM, Steketee JD. Time-dependent effects of repeated cocaine administration on dopamine transmission in the medial prefrontal cortex. *Neuropharmacology*. 2005; 48:51–61. [PubMed: 15617727]
- Zhou FM, Hablitz JJ. Dopamine modulation of membrane and synaptic properties of interneurons in rat cerebral cortex. *J Neurophysiol*. 1999; 81:967–976. [PubMed: 10085325]

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 1. Dopamine effects on the amplitude of synaptically evoked IPSCs

A, In saline-treated animals bath application of 20 μM DA produced a temporally biphasic effect on the amplitude of eIPSCs, resulting in an initial decrease followed by a prolonged increase in the amplitude of the eIPSC. *Top*, Representative traces of eIPSCs at 3 different time points. Traces are averaged sweeps of 10 responses during baseline, or “early” (5–10 min after DA bath application) and “late” (25–30 min after DA application) time periods used to measure drug effects. *Bottom*, Group data showing the time course of eIPSC amplitude changes in saline-treated animals (n=8). Changes in amplitude are relative to the averaged response of the baseline period. The large vertical bar indicates the duration of DA application. Small horizontal bars indicate the periods from which averaged measures were used for statistical comparisons of “early” (black bar) and “late” (grey bar) drug effects. The numbers above the bars indicate the mean (\pm SEM) percentage change in amplitude during these periods. *B*, *Bottom*: In cocaine-treated animals (n=7) the initial reduction in eIPSC amplitude was absent and instead the eIPSC amplitude showed a rapid and persistent increase. *Top*: Representative traces of eIPSCs for the same time periods as indicated in *A*.




Figure 2. Pharmacology of DA effects on eIPSCs in saline-treated and cocaine-sensitized animals

A. Bath application of the D2-type receptor agonist quinpirole (5 μ M) induced a rapid reduction in eIPSC amplitude in saline-treated animals. The graph shows the time course of eIPSC amplitude changes relative to the averaged response of the baseline period. In all graphs the large vertical bar indicates the period of drug application. Horizontal bars indicate the periods from which averaged measures were used for statistical comparisons of “early” (black) and “late” (grey) drug effects, and the numbers indicate the mean (\pm SEM) percentage change in amplitude during these 5 minute periods. **B.** Activation of D2-type receptors in cocaine-treated animals (n=11) failed to induce a similar reduction in eIPSC

amplitude as seen in saline animals. The differential responses to quinpirole in the two treatment groups parallel the effects of DA on the transient reduction in eIPSCs amplitude in saline- and cocaine-treated animals, respectively. *C, D*. In contrast to D2 receptor activation, D1 receptor activation is unaltered in cocaine-treated rats. Bath application of the D1 agonist SKF-81297 (*C*) or of DA (20 μ M) in the presence of the D2 receptor antagonist sulpiride (*D*) evoked a rapid and persistent increase in the amplitude of the eIPSCs ($n=7$). Sulpiride was present in the bath for the duration of the recording (black line). *E*. Bath application of a low dose (100 nM) of DA also results in an immediate and long-lasting increase in eIPSC amplitude. Low concentrations of DA ($< 1\mu$ M) were previously reported to preferentially activate D1 receptors (Trantham-Davidson, *et al* 2004). See text for details.




Figure 3. Dopamine modulation of spontaneous IPSCs in saline- and cocaine-treated animals
 A) Example traces showing voltage-clamp recordings of pharmacologically isolated sIPSCs from PFC pyramidal cells under baseline conditions (left column) and after application of 20 μM DA. Most synaptic events were blocked by bath application of the GABAA antagonist bicuculline (20 μM) at the end of the experiments. Both in saline- and in cocaine-treated animals DA led to an increase in the frequency of sIPSCs, albeit with different time courses (B, C). *B*, Time course and amplitude histograms of the effect of 20 μM DA on sIPSCs in saline-treated animals (n=8). *Left*, The graph shows minute by minute changes in the

frequency of sIPSCs compared to the baseline average. The frequency rose slowly and reached a relatively stable level after about 20 minutes. The periods of measurement of early (5–10 min, light grey) and late (25–30 min, dark grey) DA effects are indicated by horizontal bars. The increase in the average frequency of sIPSCs was significant for the late measurement ($p < 0.05$). *Right*, Plotting the relative change in the number of sIPSCs across a range of amplitudes (10 pA bins) revealed that DA consistently increased events of all amplitudes with the exception of very small events (< 10 pA) which showed a decrease in frequency during the period of the early DA effect. *Inset*, Changes in the overall frequency and the relative distribution of sIPSCs did not affect the amplitude of the average sIPSC. *C*, Time course and amplitude histograms of the effect of 20 μ M DA on sIPSCs in cocaine-treated animals ($n=6$). *Left*, In cocaine-treated animals the frequency of sIPSCs rapidly increased. The increase in the average frequency of sIPSCs was significant at both the early and late measurement ($p < 0.05$). *Right*, Plotting the relative change in the number of sIPSCs across a range of amplitudes showed a consistent increase of events across all amplitudes following DA application. *Inset*, Similar to the effects in saline animals any shift in the relative distribution of events was not large enough to affect the average sIPSC amplitude.




Figure 4. Pharmacology of dopamine modulation of spontaneous IPSCs in saline- and cocaine-treated animals

A, B, Time course of changes in sIPSC frequency following D2 receptor stimulation with quinpirole (5 μM) in saline- (*A*) and cocaine-treated (*B*) animals. In saline animals (*n*=6) there was a slowly developing reduction in the number of sIPSCs that became significant by the time of the late measurement period (see *D*). In cocaine-treated animals there was no discernible effect of D2 receptor stimulation. *C*, In contrast, D1 receptor stimulation with SKF-81297 produced a rapid and persistent increase in the frequency of sIPSCs in cocaine-treated animals (*n*=6). Similar increases were seen with bath application of 100 nM DA (see *D*). *D*, Summary graph showing the average changes in sIPSC frequency during the early (5–10 min following drug application) and late (25–30 min following drug application) measurement periods for the data shown in *A–C*, as well as results obtained with bath application of 100 nM DA (*n*=8) (* *p*<0.05, ** *p*<0.01, paired *t*-tests).




Figure 5. Dopamine and D2 receptor-dependent modulation of action potential independent miniature IPSCs in saline- and cocaine-treated animals

A, In saline-treated animals (n=7) quinpirole (5 μM) significantly reduced the frequency and the average amplitude of mIPSCs. *Top*, Representative traces showing voltage clamp recordings of mIPSCs in PFC pyramidal cells before (left column) and after quinpirole application. *A2*, Time course of quinpirole effects on the frequency of mIPSCs. The inset shows a comparison of the average frequency of events from all cells during baseline and 5–10 minutes after quinpirole application. *A3* Time course of quinpirole effects on the average

amplitude of mIPSCs. The reduction in mIPSC amplitude became significant during the end of our recordings (23–27 min post quinpirole application; $-34.7 \pm 10\%$; $p < 0.05$) before experiments were terminated by the experimenter. *B*, In cocaine-treated animals ($n=7$) quinpirole did not consistently change the frequency or amplitude of mIPSCs. *Top*, Representative traces showing voltage clamp recordings of mIPSCs before (left column) and after quinpirole application. *B2*, Time course of quinpirole effects and average frequency of events from all cells during baseline and 5–10 minutes after quinpirole application (*inset*). *B3*, Time course of quinpirole effects on average mIPSC amplitude. *C, D*, Activation of both D1 and D2 receptors with the full agonist DA (20 μM) did not significantly change the frequency of mIPSCs in saline-treated animals (*C*, $n=4$), but as shown in *D* led to a significant increase in the number of mIPSCs in cocaine-sensitized animals ($n=7$; * $p < 0.05$, ** $p < 0.01$, paired t-tests).




Figure 6. D2 receptor-dependent modulation of current-evoked action potential firing in interneurons from saline- and cocaine-treated animals

A, Examples of fast-spiking (FS) and non-fast spiking (NFS) interneurons recorded in the prelimbic cortex of adult rats. Cells were filled with Alexa 594 during whole-cell recording and reconstructed using z-stacked confocal images. Scale bar is 50 μm. B, Example sweeps showing that D2 receptor stimulation with quinpirole (5 μM) increases action potential firing evoked by somatic current injection in both NFS and FS cells from saline-treated animals (left column), but not from cocaine-treated animals. Scale bars represent 200 ms and 20 mV.

C, Summary plot for the effects of quinpirole on evoked spike firing in FS cells (open symbols) and NFS cells (grey squares) in saline- and cocaine-treated animals. Individual values for each cell are averages of at least 3 repetitions for baseline and quinpirole condition, respectively. *D*, Same data as in *C* plotted to show the average change in the number of action potentials for FS and NFS cells in saline- and cocaine-treated animals (* $p < 0.05$ paired t-tests).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Electrophysiological properties of 3 classes of neurons in layers 5/6 of rat prefrontal cortex from saline- and cocaine-treated animals (means \pm SEM). In animals from both experimental groups FS and NFS cells differed with regard to the duration of their action-potentials at half-width and the amplitude of their AHPs. The average properties of FS cells and NFS cells, respectively, did not differ when comparing cells belonging to the same class of neurons from saline- and cocaine-treated animals.

	Saline		Cocaine		NFS (n=5)	Saline		Cocaine		Significant differences
	FS (n=6)	NFS (n=7)	FS (n=8)	NFS (n=7)		Pyramidal (n=13)	Pyramidal (n=6)			
Input resistance (MegaOhm)	223.4 \pm 34.7	262.7 \pm 66.6	202.6 \pm 29.0	241.6 \pm 29.3	123.9 \pm 13.5	162 \pm 35.6	—			—
Rheobase current (pA)	145.0 \pm 32.6	94.3 \pm 20.2	142.5 \pm 30.3	109.0 \pm 13.9	137.6 \pm 12.0	115 \pm 11.2	—			—
AP threshold (mV)	-39.9 \pm 2.3	-44.6 \pm 1.0	-40.7 \pm 1.2	-40.4 \pm 1.5	-41.9 \pm 1.1	-42.3 \pm 1.3	—			—
AP duration at half-width (ms)	0.67 \pm 0.03	1.21 \pm 0.08	0.71 \pm 0.07	1.18 \pm 0.09	1.58 \pm 0.16	1.3 \pm 0.1	SAL: NFS>FS COC: NFS>FS			
AHP amplitude (mV)	17.4 \pm 1.17	7.2 \pm 1.16	19.7 \pm 1.16	7.5 \pm 1.4	6.5 \pm 1.09	6.7 \pm 1.05	SAL: FS>NFS COC: FS>NFS			

Abbreviations: AHP, afterhyperpolarization; AP, action potential. FS, fast-spiking cells; NFS, non-fast-spiking cells. Significant differences indicated by the use of Student's t-tests (** $p < 0.01$).