



Published in final edited form as:

J Neurosci. 2010 December 1; 30(48): 16170–16179. doi:10.1523/JNEUROSCI.2323-10.2010.

Modulation of BK channel gating by the $\beta 2$ subunit involves both membrane-spanning and cytoplasmic domains of Slo1

Urvi S. Lee, Jingyi Shi, and Jianmin Cui*

Department of Biomedical Engineering and Cardiac Bioelectricity and Arrhythmia Center, Washington University, St. Louis, MO 63130

Abstract

Large conductance, Ca^{2+} - and voltage-sensitive K^+ (BK) channels regulate neuronal functions such as spike frequency adaptation and transmitter release. BK channels are composed of four Slo1 subunits, which contain the voltage sensing and pore-gate domains in the membrane and Ca^{2+} binding sites in the cytoplasmic domain, and accessory β subunits. Four types of BK channel β subunits ($\beta 1$ – $\beta 4$) show differential tissue distribution and unique functional modulation, resulting in diverse phenotypes of BK channels. Previous studies show that both the $\beta 1$ and $\beta 2$ subunit increase Ca^{2+} sensitivity, but different mechanisms may underline these modulations. However, the structural domains in Slo1 that are critical for Ca^{2+} dependent activation and targeted by these β subunits are not known. Here, we report that the N-termini of both the transmembrane (including S0) and cytoplasmic domains of Slo1 are critical for $\beta 2$ modulation based on the study of differential effects of the $\beta 2$ subunit on two orthologs, mSlo1 and dSlo1. The N-terminus of the cytoplasmic domain of Slo1 including the AC region of the RCK1 domain and the peptide linking it to S6, both of which have been previously shown to mediate the coupling between Ca^{2+} binding and channel opening, is specifically required for the $\beta 2$ but not for the $\beta 1$ modulation. These results suggest that the $\beta 2$ subunit modulates the coupling between Ca^{2+} binding and channel opening, and, although sharing structural homology, the BK channel β subunits interact with structural domains in the Slo1 subunit differently to enhance channel activity.

Keywords

BK; Slo1; β subunit; Ca^{2+} sensitivity; AC Region; S0

Introduction

BK channels activate in response to depolarizing membrane potentials and increases in intracellular Ca^{2+} (Marty, 1981). The opening of BK channels effectively repolarizes the membrane and shuts Ca^{2+} entry through voltage-dependent Ca^{2+} channels, thereby reducing intracellular Ca^{2+} . As a consequence, BK channels are negative feedback regulators of membrane excitability and Ca^{2+} entry in neurons (Lancaster and Nicoll, 1987; Storm, 1987) and smooth muscle cells (Brayden and Nelson, 1992).

Functional BK channels are tetramers of the pore-forming Slo1 subunit which is composed of seven membrane-spanning segments (S0–S6), where S1–S4 form the voltage sensing domain (VSD) and S5–S6 form the pore-gate domain, and a large cytoplasmic domain (Fig.

*Correspondence should be addressed to: Jianmin Cui, Department of Biomedical Engineering, Washington University in St. Louis, Campus Box 1097, 1 Brookings Drive, St. Louis, MO 63130, Tel: (314)935-8896, Fax: (314)935-7448, jcui@biomed.wustl.edu.

1) (Meera et al., 1997). This cytoplasmic domain is composed of two RCK (regulator of K⁺ conductance) domains and contains two putative Ca²⁺ binding sites, D367 in RCK1 and the Ca²⁺ bowl in RCK2 (Fig. 1) (Schreiber and Salkoff, 1997; Xia et al., 2002; Wu et al., 2010; Yuan et al., 2010).

BK channels also associate with β subunits that are composed of two transmembrane (TM) segments with cytoplasmic termini and a large extracellular loop. There are four types of β subunits, β 1– β 4, that have tissue-specific distributions and impart unique effect on voltage and Ca²⁺ dependent activation of BK channels (Orio et al., 2002). The β 1 and β 2 subunits have the highest sequence homology and both increase apparent Ca²⁺ sensitivity (Brenner et al., 2000), but different mechanisms may be responsible for their functions; the β 1 subunit was shown to primarily modify the voltage dependent activation, while the β 2 subunit shows less effects on the voltage sensor, although both affect the Ca²⁺ binding affinity and the allosteric coupling between sensors, and sensors and gate (Cox and Aldrich, 2000; Orio and Latorre, 2005; Yang et al., 2008).

Previous studies have identified the N-terminus, S0 segment, and voltage sensing residues in Slo1 play important roles for the function of the β 1 subunit (Wallner et al., 1996; Morrow et al., 2006; Yang et al., 2008). However, the structural components in Slo1 that are important for the function of the β 2 subunit are not known. In this study, we take advantage of differential effects of the β 2 subunit on two BK channel orthologs to make chimeras between them and test which structural component confers the properties of its parent channel in response to the modulation by the β 2 subunit. These experiments identified two regions of the Slo1 subunit that are critical for the β 2 dependent increase of Ca²⁺ sensitivity. The first region contains the extracellular N-terminus and S0 segment. The second region, which is specifically important for the β 2 dependent increase of Ca²⁺ sensitivity, contains the linker connecting S6 and the cytoplasmic domain, the C-Linker, and the N-terminus of RCK1, the AC region (Fig. 1). This segment has been previously shown to mediate the allosteric coupling between Ca²⁺ binding and the opening of the activation gate (Niu et al., 2004; Krishnamoorthy et al., 2005; Yang et al., 2010). Our results in this study support the role of this segment in Ca²⁺ dependent activation of BK channels and suggest that the β 2 subunit enhances the allosteric coupling between Ca²⁺ binding and the opening of the activation gate.

Materials and Methods

Mutagenesis and Expression

All chimeras and point mutations were generated from the *mbr5* splice variant of mouse Slo1 (mSlo1) (Butler et al., 1993) and the *A1C2E1G310* splice variant of drosophila Slo1 (dSlo1) (Adelman et al., 1992) in the pBSC-MXT vector. The protein sequences of mSlo1 and dSlo1 used in the design of each chimera is shown in Table 1. Human β 1 and β 2 (KCNMB1 and KCNMB2; GenBank Accession No. U25138 and AF209747, respectively) cDNAs were subcloned into pcDNA3.1(+). The β 2 with N-terminus deleted (β 2ND) subunit was created by removing amino acids from position 2 through 20. All chimeras and mutations were made using overlap-extension PCR (Shi et al., 2002) with *Pfu* polymerase (Stratagene, La Jolla, CA). The PCR-amplified regions for all constructs were verified by sequencing. cRNA was transcribed *in vitro* using T3 polymerase (Ambion, Austin, TX) for all Slo1 constructs and T7 polymerase (Ambion, Austin, TX) for β 1 and β 2ND subunits. 0.05–20 ng of Slo1 cRNA or a mixture of 5–15 ng Slo1 and 25–50 ng β subunit cRNAs were injected into each *Xenopus laevis* oocyte. Oocytes were incubated in ND96 solution (in mM: 96 NaCl, 2 KCl, 1.8 CaCl₂, 1 MgCl₂, 5 HEPES, pH 7.6) at 18°C for 3–6 days before recording.

Electrophysiology

Macroscopic currents were recorded from inside-out patches formed with borosilicate pipettes of ~0.9–1.5 MΩ resistance. The data was acquired using an Axopatch 200-B patch-clamp amplifier (Axon Instruments, Union City, CA) and Pulse acquisition software (HEKA Elektronik, Lambrecht/Pfalz, Germany). Recordings were digitized at 20 μs intervals and low-pass filtered at 10 kHz with the 4-pole Bessel filter built in the amplifier. Capacitive transients and leak currents were subtracted using a P/5 protocol. Experiments were conducted at room temperature (20°C~22°C). The pipette solution contained (in mM) 140 KMeSO₃, 20 Hepes, 2 KCl, and 2 MgCl₂, pH 7.2. The internal solution contained (in mM) 140 KMeSO₃, 20 Hepes, 2 KCl, and 1 N-(2-hydroxyethyl)ethylenediamine-*N, N, N*-triacetic acid (HEDTA), pH 7.2. CaCl₂ was added to the internal solution to give the appropriate free [Ca²⁺]_i, which was measured with a calcium-sensitive electrode (Orion Research, Cambridge, MA). 50 μM 18-crown-6-tetracarboxylic acid (Sigma-Aldrich, St. Louis, MO) was added to internal solutions to chelate Ba²⁺. For nominal 0 μM [Ca²⁺]_i, the same internal solution was used except that HEDTA was substituted by 5 mM EGTA and no CaCl₂ was added. The free [Ca²⁺] in nominal 0 μM [Ca²⁺]_i solution is 0.5 nM. Experiments were conducted over a period of time where multiple different [Ca²⁺]_i were made and their concentrations varied. For the sake of simplicity the 2 μM Ca²⁺ is referred to as ~2 μM in the text where the concentration varied from 1.8 to 2.3 μM and the saturating [Ca²⁺]_i is referred to as ~100 μM where the concentration varied from 87.1 and 118.4 μM.

Data Analysis

The relative conductance was determined by measuring tail current amplitudes at indicated voltages for all channels with and without β subunits. The conductance-voltage (G-V) relationships for all channels with and without β subunits were fitted with the Boltzmann

equation: $\frac{G}{G_{max}} = \frac{1}{1 + e^{-ze(V-V_{1/2})/kT}}$ where G/G_{max} is the ratio of conductance to maximum conductance in the same [Ca²⁺]_i, z is the number of equivalent charges, $V_{1/2}$ is the voltage at which the channel is 50% activated, e is the elementary charge, k is Boltzmann's constant, and T is the absolute temperature. Curve fittings were done with Igor Pro software (WaveMetrics, Inc., Lake Oswego, OR) using the Levenberg-Marquardt algorithm to perform nonlinear least squares fits. Statistics were performed using Origin 6.1 (OriginLab Corp., Northampton, MA); independent/unpaired t-test was performed. A p value of <0.05 was considered significant.

Monod-Wyman-Changeux (MWC) model

MWC model fits were performed using the following equation:

$$P_{open} = \frac{1}{1 + L_0 \cdot e^{\frac{-zeV}{kT}} \cdot \left[\frac{1 + \frac{[Ca^{2+}]}{K_C}}{1 + \frac{[Ca^{2+}]}{K_O}} \right]^4}$$

where P_{open} is the channel's open probability, L_0 is the steady-state equilibrium constant from open to close channels ($[C_0]/[O_0]$) in the absence of Ca²⁺ binding at 0 mV, z , e , k and T are same as in the Boltzmann equation (see above), K_C and K_O are the dissociation constants of Ca²⁺ in the closed and open states, respectively. MWC model code was written and executed in MATLAB V7.4 (The MathWorks, Inc.).

Since BK channels are activated by both voltage and Ca²⁺, the ideal condition for measuring Ca²⁺ sensitivity of channel activation would be in the absence of voltage sensor movements.

Such measurements have been done at very negative voltages (< -130 mV) where the voltage sensor of BK channels is kept at the resting state (Horrigan and Aldrich, 2002). Ca^{2+} dependent activation of BK channels under such a condition could be fitted by the MWC model, and the results show that the parameters K_C and K_O have very similar values as those obtained by the MWC model fitting to the G-V relations (Cox et al., 1997; Horrigan and Aldrich, 2002; Sweet and Cox, 2008; Yang et al., 2010) which is the method we used in this study. The lack of a large influence of voltage on Ca^{2+} affinity measurement is not surprising because it has been shown that voltage and Ca^{2+} activate the channel through distinct mechanisms that have little interaction (Cui and Aldrich, 2000; Horrigan and Aldrich, 2002; Sweet and Cox, 2008). In the MWC model used in this study, each Slo1 subunit is assumed to contain a single Ca^{2+} binding site although previous studies have proposed two putative Ca^{2+} binding sites in each Slo1 subunit (Schreiber and Salkoff, 1997; Xia et al., 2002). A model composed of two Ca^{2+} binding sites would not provide any additional information since we are interested in the effect on the overall Ca^{2+} sensitivity of the BK channel. In spite of this simplification, the MWC model used in this study provides an appropriate account of the Ca^{2+} sensitivity changes in BK channels.

Results

The $\beta 2$ ND subunit modulates Ca^{2+} sensitivity of mSlo1 and dSlo1 differently

Besides increasing Ca^{2+} sensitivity of mSlo1 channels, the $\beta 2$ subunit also inactivates the channel by a “ball and chain” mechanism as such that the N-terminus of the $\beta 2$ subunit blocks the open channel (Ding et al., 1998; Hicks and Marrion, 1998; Wallner et al., 1999; Xia et al., 1999, 2003) (Fig. 2A). To study increases in Ca^{2+} sensitivity conferred by the $\beta 2$ subunit in the absence of inactivation, we deleted the amino acids 2–20 of the human $\beta 2$ subunit, which resulted in the $\beta 2$ N-terminus deleted ($\beta 2$ ND) subunit that no longer causes inactivation (Fig. 2B) (Wallner et al., 1999; Xia et al., 1999; Brenner et al., 2000; Xia et al., 2003). We coexpressed the $\beta 2$ ND subunit with mSlo1 and dSlo1 and observed the changes to their properties (Fig. 2). The association of $\beta 2$ ND increases the rate of activation of mSlo1 (Brenner et al., 2000; Orio and Latorre, 2005; Orio et al., 2006; Lee and Cui, 2009) and decreases the rate of deactivation of both mSlo1 and dSlo1 (Fig. 2B). Since the association of $\beta 2$ ND with mSlo1 shifts G-V relation to more negative voltages the deactivation time course was measured at a more negative voltage with $\beta 2$ ND than without $\beta 2$ ND. However, mSlo1 with $\beta 2$ ND still deactivates more slowly than mSlo1 alone (τ_{Dact} : 5.7 ± 0.3 ms at -120 mV and 1.8 ± 0.1 ms at -50 mV, respectively when prepulsed to $+100$ mV, $p < 0.001$ with $n = 14$ and 55 , respectively). Measured at the same voltage, dSlo1 with $\beta 2$ ND also deactivates significantly more slowly than dSlo1 alone (τ_{Dact} : 1.5 ± 0.2 ms and 0.8 ± 0.1 ms at -50 mV, respectively when prepulsed to $+200$ mV, $p < 0.001$ with $n = 7$ and 23 , respectively). These changes in the macroscopic currents indicate a physical association between the $\beta 2$ ND subunit and Slo1 orthologs. Coexpression of the wild type (WT) $\beta 2$ subunit with both mSlo1 and dSlo1 confer inactivation in the channels (Fig. 2A), further indicating that the $\beta 2$ subunits associate with both Slo1 orthologs. However, the $\beta 2$ ND subunit has different effects on the Ca^{2+} dependent activation of the two Slo1 orthologs. The association of the $\beta 2$ ND subunit shifts the conductance-voltage (G-V) relation of mSlo1 channels to more negative potentials by ~ 100 mV at the saturating $[\text{Ca}^{2+}]_i$ of ~ 100 μM but has little effect on the G-V of dSlo1 channels ($V_{1/2}$ of dSlo1: -19.3 ± 2.5 mV and dSlo1+ $\beta 2$ ND: -23.8 ± 2.7 mV, $P > 0.05$; $V_{1/2}$ is the voltage where the G-V relation is half maximum) (Fig. 2C). At nominal 0 $[\text{Ca}^{2+}]_i$, the $\beta 2$ ND association does not alter the G-V relation of the mSlo1 channels ($V_{1/2}$ of mSlo1: 178.3 ± 1.1 mV and mSlo1+ $\beta 2$ ND: 181.9 ± 1.8 mV, $P > 0.05$) (Fig. 2C) (Orio and Latorre, 2005), indicating that $\beta 2$ ND modulates Ca^{2+} dependent activation of mSlo1. In contrast, the association of $\beta 2$ ND with dSlo1 has no significant effect at any $[\text{Ca}^{2+}]_i$ tested in our study. For instance at ~ 2 μM Ca^{2+}_i , $V_{1/2}$ of

dSlo1 and dSlo1+ β 2ND is 243.4 ± 6.5 mV and 241.5 ± 5.2 mV, respectively ($P > 0.05$) (Fig. 2C). The plot of $V_{1/2}$ versus $[Ca^{2+}]_i$ shows that the association of the β 2ND subunit results in a gradual increase in the negative shift of the mSlo1 G-V relation with increasing $[Ca^{2+}]_i$ but no shift of the dSlo1 G-V relation (Fig. 2D). In $[Ca^{2+}]_i \leq 1 \mu M$, few dSlo1 channels activate at voltages within the experimental limit (up to +300 mV) so that the G-V relations cannot be determined in these $[Ca^{2+}]_i$ s. The results in Figure 2 clearly show that the β 2ND subunit modulates Ca^{2+} dependent activation of mSlo1 but not dSlo1.

To further illustrate the differences in Ca^{2+} sensitivity imparted by β 2ND on mSlo1 and dSlo1 channels, a voltage dependent Monod-Wyman-Changeux (MWC) model (Cox et al., 1997) was used to fit the G-V relations at different $[Ca^{2+}]_i$ s (Fig. 6, Table 2). In mSlo1 channels, the association of β 2ND subunit increases the dissociation constant for Ca^{2+} binding in the closed state (K_C) by $>300\%$ and decreases the dissociation constant in the open state (K_O) by 14% (Table 2), same as previously reported (Orio and Latorre, 2005). Due to the disproportional change in the two dissociation constants, the ratio $c = K_O/K_C$ is smaller in mSlo1+ β 2ND than in mSlo1 channels, which indicates that Ca^{2+} favors the open state in the closed-open equilibrium more with the association of β 2ND (Table 2) (Cox et al., 1997). The association of β 2ND does not alter voltage dependence of mSlo1 channels, which is reflected in similar values of the parameters Z and L_0 , where Z is the number of gating charges in the voltage sensor of the channels and L_0 is the equilibrium constant of channel from the open to the closed state at 0 mV and in the absence of Ca^{2+} (Table 2). In dSlo1 channels, the association of β 2ND does not change K_C , K_O , or voltage dependence (Table 2). Collectively, these results show that the β 2ND subunit specifically enhances Ca^{2+} sensitivity in mSlo1 channels, but results in no such changes in dSlo1 channels, even though β 2ND is associated with dSlo1 subunit.

The N-termini of the transmembrane and cytoplasmic domains of mSlo1 and dSlo1 are necessary and sufficient to account for the different effects of the β 2ND subunit

Taking advantage of the differences in the response of the Slo1 orthologs to β 2ND modulation, a chimera strategy to exchange the secondary structural regions between mSlo1 and dSlo1 subunits was devised to find the structural components responsible for the β 2 dependent enhancement of Ca^{2+} sensitivity. To examine whether the association of β 2ND alters Ca^{2+} sensitivity of a channel, two criteria were used: first, Ca^{2+} sensitivity of a channel was estimated by the shift of its G-V relation in $\sim 100 \mu M$ $[Ca^{2+}]_i$ in response to the association of β 2ND, $\Delta V_{half} = V_{1/2(100Ca)}$ with β 2ND - $V_{1/2(100Ca)}$ without β 2ND; second, Ca^{2+} sensitivity of a channel was computed as the G-V shift $\Delta V_{1/2(0/2-100Ca)}$, in response to an increase of $[Ca^{2+}]_i$ from 0 or ~ 2 to $\sim 100 \mu M$, the change in Ca^{2+} sensitivity resulted from the association of β 2ND was $\Delta \Delta V_{half} = \Delta V_{1/2(0/2-100Ca)}$ with β 2ND - $\Delta V_{1/2(0/2-100Ca)}$ without β 2ND, which was then normalized to $\Delta \Delta V_{half}$ of the WT mSlo1 obtained under the same condition (Fig. 3). Using these two criteria, mSlo1 exhibited greater modulation by the association of β 2ND than dSlo1 (Fig. 3).

The chimeras developed for this study are shown in the bottom half of Figure 3, where the initial designs (chimeras C1 through m[dS0]) were composed of dSlo1 channels with increasing mSlo1 regions substituting the dSlo1 counterparts. In chimeras C1-C5, the addition of mSlo1 regions did not confer β 2ND modulation; rather the Ca^{2+} sensitivity of these channels and their responses to the association of β 2ND were similar to that of dSlo1 (Fig. 3). Even the chimera containing only the N-terminus of dSlo1 including the transmembrane segment S0 (m[dS0]), shows a $\sim 40\%$ reduced response to the β 2ND modulation, indicating that the N-terminus of mSlo1 is important for the β 2ND modulation of Ca^{2+} sensitivity (Fig. 3). The result of m[dS0] in Figure 3 is computed from the current measurements and G-V relations shown in Figure 4A and B. Figure 4C shows further studies of the Ca^{2+} sensitivity of this chimera where the $V_{1/2}$ of G-V relations of the channel

with (black unfilled circles) and without (black filled circles) β 2ND in various $[Ca^{2+}]_i$ from nominal 0 to saturating $\sim 100 \mu M$ are plotted versus $[Ca^{2+}]_i$. For comparison, similar data from dSlo1 (blue dotted and solid lines) and mSlo1 (orange dotted and solid lines) are also shown (Fig. 4C). The β 2ND subunit enhances Ca^{2+} sensitivity of mSlo1 channels and gradually shifts its G-V relation as $[Ca^{2+}]_i$ increases (Fig. 2) so that the $V_{1/2}-[Ca^{2+}]_i$ relations of mSlo1 channels with and without β 2ND bifurcate (Fig. 4C in orange). To the contrary, the β 2ND subunit does not alter Ca^{2+} sensitivity of dSlo1 channels so that the $V_{1/2}-[Ca^{2+}]_i$ relations of dSlo1 channels with and without β 2ND superimpose (Fig. 4C in blue). While the $V_{1/2}-[Ca^{2+}]_i$ relations of m[dS0] with and without β 2ND bifurcate (Fig. 4C in black), the difference between the two curves is smaller than that of mSlo1 at all non-zero $[Ca^{2+}]_i$, clearly showing that the N-terminus of mSlo1 is important for the β 2ND dependent enhancement of Ca^{2+} sensitivity.

To examine if the N-terminus of mSlo1 alone is responsible for the β 2ND dependent enhancement of Ca^{2+} sensitivity, we made the converse chimera of m[dS0], d[mS0], in which the N-terminus to the end of S0 of mSlo1 is transplanted into dSlo1. Since few d[mS0] channels open in $[Ca^{2+}]_i < \sim 2 \mu M$ (Fig. 5A and B), the change of Ca^{2+} sensitivity due to the association of β 2ND is only estimated by ΔV_{half} in Figure 3, which suggests that the N-terminus of mSlo1 is not sufficient to render dSlo1 the same response to β 2ND as mSlo1. Further studies of Ca^{2+} sensitivity of this chimera in various $[Ca^{2+}]_i$ show that the $V_{1/2}-[Ca^{2+}]_i$ relations of d[mS0] with and without β 2ND only bifurcate at high $[Ca^{2+}]_i$ ($> \sim 30 \mu M$) and the difference between the two curves is smaller than that of mSlo1, demonstrating that the N-terminus of mSlo1 is not the only structural component important for the β 2ND dependent enhancement of Ca^{2+} sensitivity (Fig. 5C).

Previous studies have shown that the AC region (the $\beta A-\alpha C$ of RCK1 domain) (Krishnamoorthy et al., 2005; Yang et al., 2010) and the C-Linker (Niu et al., 2004; Krishnamoorthy et al., 2005) are important in coupling Ca^{2+} binding to opening of the activation gate. To examine if this segment is also important for the β 2ND dependent enhancement of Ca^{2+} sensitivity, we studied the chimera channels in which this segment from dSlo1 is transplanted into mSlo1, m[dLinker+dAC]. Its response to β 2ND is estimated using ΔV_{half} (Fig. 3), which shows a dramatic reduction from that of mSlo1 and is similar to that of dSlo1, suggesting that the N-terminus of the cytoplasmic domain is important for the β 2ND dependent enhancement of Ca^{2+} sensitivity. Surprisingly, although the inactivation “ball” is deleted in β 2ND, its association with m[dLinker+dAC] causes inactivation of the channel, (Fig. 4A). The mechanism of this inactivation is not clear at this time and needs further study. Nevertheless, further studies of the Ca^{2+} sensitivity of this chimera in several $[Ca^{2+}]_i$'s shows that the $V_{1/2}-[Ca^{2+}]_i$ relations with and without β 2ND superimpose (Fig. 4C), confirming that β 2ND can no longer enhance Ca^{2+} sensitivity of this channel. The converse chimera in the dSlo1 background, d[mLinker+mAC] showed no increase in Ca^{2+} sensitivity with the association of β 2ND either (Fig. 5). Taken together, these results suggest that the N-terminus of the cytoplasmic domain of mSlo1 is also necessary but not sufficient to account for the difference in the β 2ND dependent increase of Ca^{2+} sensitivity between mSlo1 and dSlo1.

Either terminus of the transmembrane or cytoplasmic domain of mSlo1 is important for the β 2ND dependent increase of Ca^{2+} sensitivity, and as expected, the chimera containing both termini from dSlo1 in the mSlo1 background, m[dS0+dLinker+dAC], showed no increase in Ca^{2+} sensitivity in response to β 2ND, similar to dSlo1 (Fig. 3 and 4). We studied the converse chimera in dSlo1 background, d[mS0+mLinker+mAC], in order to examine if N-termini of both the transmembrane and cytoplasmic domains of mSlo1 are sufficient to render the β 2ND dependent increase of Ca^{2+} sensitivity in dSlo1. However, this chimera failed to express any functional channel with β 2ND subunit for reasons that are currently not

understood. Therefore, we modified this chimera by extending the AC region to include β G of the RCK1 domain that results in d[mS0+mLinker+mRCK1c]. The coexpression of this chimera with β 2ND showed increased Ca^{2+} sensitivity that is more like mSlo1 than dSlo1 (Fig. 3 and 5), suggesting that the N-termini of the transmembrane and cytoplasmic domains of mSlo1 are critical for the differential β 2ND dependent enhancement of Ca^{2+} sensitivity in dSlo1 and mSlo1.

The swapping of channel properties is further illustrated by fitting the G-V relations of chimeras m[dS0+dLinker+dAC] and d[mS0+mLinker+mRCK1c] to the MWC model (Fig. 6). The association of β 2ND with m[dS0+dLinker+dAC] produces a minor reduction in both K_C and K_O which does not change the “c” factor (Fig. 6, Table 2), and consequently there was a lack of G-V shift to negative potentials in the presence of Ca^{2+} (Fig. 4). However, the association of β 2ND with d[mS0+mLinker+mRCK1c] increases K_C and decreases K_O . These changes are in the same direction as those when β 2ND is associated with mSlo1, which results in reduction in “c” factor and enhanced G-V shifts in response to $[\text{Ca}^{2+}]_i$ increase (Fig. 6, Table 2).

Different molecular mechanisms for the β 1 and β 2 dependent enhancement of Ca^{2+} sensitivity

Similar to the β 2 subunit, the β 1 subunit also enhances the apparent Ca^{2+} sensitivity of BK channels to cause a G-V shift to more negative potentials in the presence of Ca^{2+} , and the two β subunits share a greater than 60% sequence homology (Brenner et al., 2000; Orio and Latorre, 2005). Furthermore, previous studies have shown that the β 1 subunit also specifically modulates Ca^{2+} sensitivity of the mSlo1 channel but not that of dSlo1, and the N-terminus of mSlo1 including S0 is critical for such modulation (Wallner et al., 1996; Morrow et al., 2006). To examine if the β 1 dependent enhancement of Ca^{2+} sensitivity is also sensitive to the difference between mSlo1 and dSlo1 in the N-terminus of the cytoplasmic domain, we studied the coexpression of the β 1 subunit with the chimera m[dLinker+dAC] (Fig. 7). In the saturating $[\text{Ca}^{2+}]_i$ of $\sim 100 \mu\text{M}$, the association of either the β 1 or β 2ND subunit shifts the G-V relation of mSlo1 to more negative potentials (Fig. 7A). The mutation m[dLinker+dAC] nearly abolishes the response of G-V relation to the association of β 2ND but not β 1 (Fig. 7A). Similar results are observed in various $[\text{Ca}^{2+}]_i$ (Fig. 7B), indicating that the differences in the structure of N-terminal part of the cytoplasmic domain between mSlo1 and dSlo1 are critical for β 2 modulation but inconsequential for β 1 function. On the other hand, for chimera m[dS0]/h[dS0] there was a lack of G-V shift in response to the association of both β 2ND and β 1 (Wallner et al., 1996), indicating that the N-terminus of the membrane-spanning domain is critical for the function of both the β 1 and β 2 subunits.

BK channels contain two high affinity Ca^{2+} binding sites located at D367 in the RCK1 domain and the Ca^{2+} bowl in the RCK2 domain (Fig. 1). Previous studies show that the two binding sites have slightly different affinities for Ca^{2+} and they contribute independently with only a small cooperativity to Ca^{2+} dependent activation (Xia et al., 2002; Qian et al., 2006; Sweet and Cox, 2008). A recent study shows that mutations in the AC region specifically affect Ca^{2+} dependent activation derived from the D367 site but not that from the Ca^{2+} bowl (Yang et al., 2010). These results indicate that the two Ca^{2+} binding sites may couple to the activation gate through distinct structural pathways. To examine which of these pathways is affected by the β subunits, we studied the effects of β 2ND and β 1 on each of the pathways by mutating the Ca^{2+} binding site in one pathway. Mutating either Ca^{2+} binding site of mSlo1 by D367A or 5D5N (5 consecutive Asp residues in the Ca^{2+} bowl are mutated to Asn) reduced overall G-V shift in response to a $[\text{Ca}^{2+}]_i$ increase from 0 to saturating $\sim 100 \mu\text{M}$ both in the absence and presence of β 2ND subunit (Fig. 8A). Nevertheless, the G-V shift is larger in the presence of β 2ND (gray bars in Fig. 8A). When

both Ca²⁺ binding sites are mutated, channel activation is no longer sensitive to Ca²⁺ either in the presence or absence of β 2ND (Fig. 8). These results indicate that the β 2 subunit modulates mSlo1 activation as long as at least one Ca²⁺ binding site is intact. Interestingly, regardless of one or both Ca²⁺ binding sites are intact, the association of β 2ND results in similar increase of G-V shift (Fig. 8B). To the contrary, while the β 1 subunit also modulates mSlo1 activation as long as at least one Ca²⁺ binding site is intact (white bars in Fig 8A), the mutation of either Ca²⁺ binding site reduces the ability of β 1 to increase G-V shift (Fig. 8B). This result suggests that although β 1 subunit is primarily known to affect the voltage dependent activation (Cox and Aldrich, 2000; Bao and Cox, 2005; Orio and Latorre, 2005; Yang et al., 2008), it may also be sensitive to the integrity of the Ca²⁺ binding site which is supported by Bao and Cox (Bao and Cox, 2005). Taken together, these experiments indicate that the β subunits modulate both pathways of Ca²⁺ dependent activation, but the β 1 and β 2 subunits may target different points in the Ca²⁺ dependent activation. Based on the recently published structures of the BK channel gating ring, the Linker and AC region reside closest to the pore (Wu et al., 2010; Yuan et al., 2010), which is the sole structural gateway to couple both Ca²⁺ binding sites to the activation gate (Fig. 9). The β 2 subunit may directly modify the allosteric mechanism in this region to affect the coupling between Ca²⁺ binding and channel opening. The β 1 subunit, on the other hand, may alter the coupling between Ca²⁺ binding and channel opening via different mechanisms, possibly involving the VSD. The difference of the two β subunits in response to Ca²⁺ binding site mutations may be the manifestations of such mechanistic differences. However, the exact nature of these allosteric mechanisms are not fully understood.

Discussion

The β subunits of BK channels modulate specific properties of the pore-forming Slo1 subunit. In the case with the β 2 subunit, the BK channels exhibit increased Ca²⁺ sensitivity by shifting the $V_{1/2}$ to more negative potentials with increasing [Ca²⁺]_i. In this study, we identified the regions in Slo1 targeted by the β 2 subunit modulation by taking advantage of different responses of two BK channel homologues, mSlo1 and dSlo1, to the β 2 subunit modulation. We found two regions in mSlo1 that are both necessary and sufficient to account for the enhanced Ca²⁺ sensitivity with the β 2 subunit as compared to dSlo1: the N-terminus of the channel to S0 and the N-terminus of the cytoplasmic domain, C-Linker/AC region. Both of these structural regions from mSlo1 are required for conferring Ca²⁺ sensitivity in dSlo1, but the replacement of either one in mSlo1 by the counterpart in dSlo1 reduced or abolished the β 2 dependent Ca²⁺ sensitivity increase.

The N-terminus of mSlo1 including S0 is important for both the β 1 and β 2 subunits to enhance Ca²⁺ sensitivity of channel activation (Fig. 3, 5) (Wallner et al., 1996; Morrow et al., 2006), which is consistent with the structural and functional role of S0. Using disulfide cross-linking, Liu et al. (Liu et al., 2008a; Liu et al., 2010) showed that S0 is an integral part of the voltage sensing domain of BK channels; its N-terminal extracellular end is close to that of S3 and S4. Consistently, mutations in S0 alter voltage dependence of channel activation, suggesting that it may interact with other membrane-spanning segments of the VSD (Koval et al., 2007). Similar disulfide cross-linking studies revealed that when the β 1 (Liu et al., 2008b; Liu et al., 2010), β 2 (Zakharov et al., 2009) and β 4 (Wu et al., 2009) subunits associate with Slo1, TM2 is located close to S0 while TM1 is located close to S2. These results suggest that S0 may directly interact with the TM2 of β subunits, while the VSD may serve as a scaffold for the conformation of β subunits. Both the β 1 and β 2 subunits are unable to enhance Ca²⁺ sensitivity of dSlo1 (Wallner et al., 1996), and a switch of the N-terminus and S0 of mSlo1 to that of dSlo1 reduces the function of both β subunits. These results may arise because, although the β subunits can associate with dSlo1 (Fig. 2),

the S0 of dSlo1 may alter either the direct TM2-S0 interaction or the structure of the VSD that leads to a conformational change in β subunits.

Contrary to the N-terminus of the channel, the C-Linker and AC region of mSlo1 are specifically important for the β 2, but not β 1, subunit to modulate Ca^{2+} sensitivity (Fig. 7). This result is consistent with previous findings that the cytoplasmic termini of the β 1 and β 2 subunits are important in determining their specific effects on channel gating (Orio et al., 2006; Wang and Brenner, 2006). Taken these results together, it is likely that the C-Linker and AC region of Slo1 may interact with the cytosolic domain of the β 2 subunit either directly or through an allosteric mechanism. Previous studies show that changes in the length of the C-Linker alter Ca^{2+} sensitivity of the channel (Niu et al., 2004), while mutations in the AC region revealed an allosteric network important for Ca^{2+} dependence of channel gating (Krishnamoorthy et al., 2005; Yang et al., 2010). Our results further support the important role of the C-Linker and AC region in mediating the coupling between Ca^{2+} binding and channel opening and show that the β 2 subunit may modulate Ca^{2+} sensitivity by altering such coupling. On the other hand, the switch of the C-Linker and AC region from mSlo1 to that of dSlo1 has no effect on the β 1 modulation of Ca^{2+} sensitivity (Fig. 7). This result could be due to that β 1 does not interact with the C-Linker or AC region so that it is not sensitive to the switch of these regions between mSlo1 and dSlo1. Alternatively, β 1 may interact with these regions but the interaction differs from how β 2ND interacts with these regions such that β 1 is insensitive to the differences between mSlo1 and dSlo1 in these regions. Due to the inherited limitations of chimera studies, our results cannot distinguish these two possibilities. Nevertheless, it is clear that the two β subunits interact with the structural domains in Slo1 differently, which is consistent with previous studies showing that the β 1 subunit alters the voltage sensor movements (Cox and Aldrich, 2000), and mutations in the VSD of Slo1 alter the modulation of the β 1 but not the β 2 subunit (Yang et al., 2008). These results provide a molecular mechanism for the previous results obtained by fitting of functional data to an allosteric model for channel activation, which show that the β 2 subunit alters Ca^{2+} binding affinities and allosteric coupling factors, while the β 1 subunit has less effects on parameters for Ca^{2+} dependent activation (Bao and Cox, 2005; Orio and Latorre, 2005).

These experimental results suggest a model, in which the TM1 and TM2 segments of the β 2 subunit associate with the VSD of mSlo1, while the cytoplasmic termini of the β 2 subunit may interact with the C-Linker and AC region in mSlo1 to alter the coupling between Ca^{2+} binding and channel opening, thereby enhancing Ca^{2+} sensitivity of channel gating (Fig. 9). The interaction between the β 2 subunit with S0 and other parts of the VSD may help the β 2 subunit adopt a correct conformation and a proper orientation toward the mSlo1 that are critical for its cytoplasmic termini to interact with the C-Linker and AC region. Therefore, mutations in either S0 and the N-terminus of the channel or the C-Linker and AC region may interrupt the interactions between the cytoplasmic termini of β 2 with the C-Linker and AC region to reduce or abolish the β 2 modulation of Ca^{2+} sensitivity.

Besides S0, other structures in Slo1 may also affect the conformation of the β 2 subunit and its orientation toward Slo1. For instance, for chimera m[dS0], in which the N-terminus including S0 of mSlo1 is replaced by the counterpart of dSlo1, the β 2 subunit still increases Ca^{2+} sensitivity although the effect is largely reduced (Fig. 3), indicating that the β 2 subunit remains in a conformation and orientation toward Slo1 that still allows some interaction with the C-Linker and AC region. On the other hand, chimera C5, in which an additional structure of mSlo1 is replaced by the counterpart of dSlo1, further reduces the β 2 modulation of Ca^{2+} sensitivity (Fig. 3). This result indicates that the S0-S1 linker and S1 of mSlo1 may also participate in the β 2 dependent enhancement of Ca^{2+} sensitivity (Fig. 3), possibly being involved in maintaining the conformation of the β 2 subunit. To test this

theory, N-terminus to S1 segment of mSlo1 were transplanted in dSlo1 to test if the C-Linker and AC region of mSlo1 is still required for the $\beta 2$ modulation of Ca^{2+} sensitivity. However, due to a large shift in its G-V relation to positive voltages, it is difficult to determine the modulation of Ca^{2+} sensitivity by the $\beta 2$ subunit (data not shown).

The $\beta 2$ subunit enhances Ca^{2+} sensitivity of mSlo1 channels, but has little effect on channel activation in the absence of Ca^{2+} (Fig. 2, 6; Table 2). Interestingly, for d[mS0+mLinker+mRCK1c] channels the association of the $\beta 2$ ND subunit results in a large shift of the G-V relation in the absence of intracellular Ca^{2+} (Fig. 5), indicating a change in the equilibrium between the open and closed conformation (Fig. 6; Table 2). This result indicates that the $\beta 2$ subunit not only alters the Ca^{2+} dependent mechanism to enhance Ca^{2+} sensitivity, but also affects the structure and mechanism important for channel opening in the absence of Ca^{2+} . These results are consistent with the previously reported results of an allosteric model fitting of BK channel gating, which showed that the enhancement of Ca^{2+} sensitivity by the $\beta 2$ subunit requires changes to the allosteric coupling factors and intrinsic channel opening (Orio and Latorre, 2005). Such an effect of the $\beta 2$ subunit may arise from its interactions with the membrane-spanning segments and VSD of Slo1 or with the C-Linker and RCK domains. It could also arise from the possibility that the association of the $\beta 2$ subunit alters the interactions between the membrane-spanning and cytoplasmic domains of Slo1 that are critical for BK channel activation. These mechanisms may also contribute to the modulation of the $\beta 2$ subunit on Ca^{2+} sensitivity.

Acknowledgments

The mSlo1 clone was kindly provided to us by Larry Salkoff (Washington University, St. Louis, MO). The $\beta 1$ and $\beta 2$ clones were kindly provided by Robert Brenner (University of Texas Health Science Center, San Antonio, TX). This work was supported by Epilepsy Foundation (U.S.L.) and National Institutes of Health grants R01-HL70393 (J.C.) and R01-NS060706 (J.C.). J. C. is the Spencer T. Olin Professor of Biomedical Engineering.

References

- Adelman JP, Shen KZ, Kavanaugh MP, Warren RA, Wu YN, Lagrutta A, Bond CT, North RA. Calcium-activated potassium channels expressed from cloned complementary DNAs. *Neuron*. 1992; 9:209–216. [PubMed: 1497890]
- Bao L, Cox DH. Gating and ionic currents reveal how the BKCa channel's Ca^{2+} sensitivity is enhanced by its beta1 subunit. *J Gen Physiol*. 2005; 126:393–412. [PubMed: 16186565]
- Brayden JE, Nelson MT. Regulation of arterial tone by activation of calcium-dependent potassium channels. *Science*. 1992; 256:532–535. [PubMed: 1373909]
- Brenner R, Jegla TJ, Wickenden A, Liu Y, Aldrich RW. Cloning and functional characterization of novel large conductance calcium-activated potassium channel beta subunits, hKCNMB3 and hKCNMB4. *J Biol Chem*. 2000; 275:6453–6461. [PubMed: 10692449]
- Butler A, Tsunoda S, McCobb DP, Wei A, Salkoff L. mSlo, a complex mouse gene encoding “maxi” calcium-activated potassium channels. *Science*. 1993; 261:221–224. [PubMed: 7687074]
- Cox DH, Aldrich RW. Role of the beta1 subunit in large-conductance Ca^{2+} -activated K^{+} channel gating energetics. Mechanisms of enhanced Ca^{2+} sensitivity. *J Gen Physiol*. 2000; 116:411–432. [PubMed: 10962017]
- Cox DH, Cui J, Aldrich RW. Allosteric gating of a large conductance Ca^{2+} -activated K^{+} channel. *J Gen Physiol*. 1997; 110:257–281. [PubMed: 9276753]
- Cui J, Aldrich RW. Allosteric linkage between voltage and Ca^{2+} -dependent activation of BK-type mslo1 K^{+} channels. *Biochemistry (Mosc)*. 2000; 39:15612–15619.
- Ding JP, Li ZW, Lingle CJ. Inactivating BK channels in rat chromaffin cells may arise from heteromultimeric assembly of distinct inactivation-competent and noninactivating subunits. *Biophys J*. 1998; 74:268–289. [PubMed: 9449328]

- Hicks GA, Marrion NV. Ca²⁺-dependent inactivation of large conductance Ca²⁺-activated K⁺ (BK) channels in rat hippocampal neurones produced by pore block from an associated particle. *J Physiol.* 1998; 508 (Pt 3):721–734. [PubMed: 9518728]
- Horrigan FT, Aldrich RW. Coupling between voltage sensor activation, Ca²⁺ binding and channel opening in large conductance (BK) potassium channels. *J Gen Physiol.* 2002; 120:267–305. [PubMed: 12198087]
- Jiang Y, Lee A, Chen J, Cadene M, Chait BT, MacKinnon R. Crystal structure and mechanism of a calcium-gated potassium channel. *Nature.* 2002; 417:515–522. [PubMed: 12037559]
- Koval OM, Fan Y, Rothberg BS. A role for the S0 transmembrane segment in voltage-dependent gating of BK channels. *J Gen Physiol.* 2007; 129:209–220. [PubMed: 17296928]
- Krishnamoorthy G, Shi J, Sept D, Cui J. The NH₂ terminus of RCK1 domain regulates Ca²⁺-dependent BK(Ca) channel gating. *J Gen Physiol.* 2005; 126:227–241. [PubMed: 16103277]
- Lancaster B, Nicoll RA. Properties of two calcium-activated hyperpolarizations in rat hippocampal neurones. *J Physiol (Lond).* 1987; 389:187–203. [PubMed: 2445972]
- Lee US, Cui J. {beta} subunit-specific modulations of BK channel function by a mutation associated with epilepsy and dyskinesia. *J Physiol.* 2009; 587:1481–1498. [PubMed: 19204046]
- Liu G, Zakharov SI, Yang L, Deng SX, Landry DW, Karlin A, Marx SO. Position and role of the BK channel alpha subunit S0 helix inferred from disulfide crosslinking. *J Gen Physiol.* 2008a; 131:537–548. [PubMed: 18474637]
- Liu G, Zakharov SI, Yang L, Wu RS, Deng SX, Landry DW, Karlin A, Marx SO. Locations of the beta1 transmembrane helices in the BK potassium channel. *Proc Natl Acad Sci U S A.* 2008b; 105:10727–10732. [PubMed: 18669652]
- Liu G, Niu X, Wu RS, Chudasama N, Yao Y, Jin X, Weinberg R, Zakharov SI, Motoike H, Marx SO, Karlin A. Location of modulatory beta subunits in BK potassium channels. *J Gen Physiol.* 2010; 135:449–459. [PubMed: 20385746]
- Marty A. Ca-dependent K channels with large unitary conductance in chromaffin cell membranes. *Nature.* 1981; 291:497–500. [PubMed: 6262657]
- Meera P, Wallner M, Song M, Toro L. Large conductance voltage- and calcium-dependent K⁺ channel, a distinct member of voltage-dependent ion channels with seven N-terminal transmembrane segments (S0-S6), an extracellular N terminus, and an intracellular (S9-S10) C terminus. *Proc Natl Acad Sci U S A.* 1997; 94:14066–14071. [PubMed: 9391153]
- Morrow JP, Zakharov SI, Liu G, Yang L, Sok AJ, Marx SO. Defining the BK channel domains required for beta1-subunit modulation. *Proc Natl Acad Sci U S A.* 2006; 103:5096–5101. [PubMed: 16549765]
- Niu X, Qian X, Magleby KL. Linker-gating ring complex as passive spring and Ca(2+)-dependent machine for a voltage- and Ca(2+)-activated potassium channel. *Neuron.* 2004; 42:745–756. [PubMed: 15182715]
- Orio P, Latorre R. Differential effects of beta 1 and beta 2 subunits on BK channel activity. *J Gen Physiol.* 2005; 125:395–411. [PubMed: 15767297]
- Orio P, Rojas P, Ferreira G, Latorre R. New disguises for an old channel: MaxiK channel beta-subunits. *News Physiol Sci.* 2002; 17:156–161. [PubMed: 12136044]
- Orio P, Torres Y, Rojas P, Carvacho I, Garcia ML, Toro L, Valverde MA, Latorre R. Structural determinants for functional coupling between the beta and alpha subunits in the Ca²⁺-activated K⁺ (BK) channel. *J Gen Physiol.* 2006; 127:191–204. [PubMed: 16446507]
- Qian X, Niu X, Magleby KL. Intra- and intersubunit cooperativity in activation of BK channels by Ca²⁺. *J Gen Physiol.* 2006; 128:389–404. [PubMed: 17001085]
- Schreiber M, Salkoff L. A novel calcium-sensing domain in the BK channel. *Biophys J.* 1997; 73:1355–1363. [PubMed: 9284303]
- Shi J, Krishnamoorthy G, Yang Y, Hu L, Chaturvedi N, Harilal D, Qin J, Cui J. Mechanism of magnesium activation of calcium-activated potassium channels. *Nature.* 2002; 418:876–880. [PubMed: 12192410]
- Storm JF. Action potential repolarization and a fast after-hyperpolarization in rat hippocampal pyramidal cells. *J Physiol.* 1987; 385:733–759. [PubMed: 2443676]

- Sweet TB, Cox DH. Measurements of the BKCa channel's high-affinity Ca²⁺ binding constants: effects of membrane voltage. *J Gen Physiol.* 2008; 132:491–505. [PubMed: 18955592]
- Wallner M, Meera P, Toro L. Determinant for beta-subunit regulation in high-conductance voltage-activated and Ca(2+)-sensitive K⁺ channels: an additional transmembrane region at the N terminus. *Proc Natl Acad Sci U S A.* 1996; 93:14922–14927. [PubMed: 8962157]
- Wallner M, Meera P, Toro L. Molecular basis of fast inactivation in voltage and Ca²⁺-activated K⁺ channels: a transmembrane beta-subunit homolog. *Proc Natl Acad Sci U S A.* 1999; 96:4137–4142. [PubMed: 10097176]
- Wang B, Brenner R. An S6 mutation in BK channels reveals beta1 subunit effects on intrinsic and voltage-dependent gating. *J Gen Physiol.* 2006; 128:731–744. [PubMed: 17130522]
- Wu RS, Chudasama N, Zakharov SI, Doshi D, Motoike H, Liu G, Yao Y, Niu X, Deng SX, Landry DW, Karlin A, Marx SO. Location of the beta 4 transmembrane helices in the BK potassium channel. *J Neurosci.* 2009; 29:8321–8328. [PubMed: 19571123]
- Wu Y, Yang Y, Ye S, Jiang Y. Structure of the gating ring from the human large-conductance Ca(2+)-gated K(+) channel. *Nature.* 2010
- Xia XM, Ding JP, Lingle CJ. Molecular basis for the inactivation of Ca²⁺- and voltage-dependent BK channels in adrenal chromaffin cells and rat insulinoma tumor cells. *J Neurosci.* 1999; 19:5255–5264. [PubMed: 10377337]
- Xia XM, Zeng X, Lingle CJ. Multiple regulatory sites in large-conductance calcium-activated potassium channels. *Nature.* 2002; 418:880–884. [PubMed: 12192411]
- Xia XM, Ding JP, Lingle CJ. Inactivation of BK channels by the NH₂ terminus of the beta2 auxiliary subunit: an essential role of a terminal peptide segment of three hydrophobic residues. *J Gen Physiol.* 2003; 121:125–148. [PubMed: 12566540]
- Yang H, Zhang G, Shi J, Lee US, Delaloye K, Cui J. Subunit-specific effect of the voltage sensor domain on Ca²⁺ sensitivity of BK channels. *Biophys J.* 2008; 94:4678–4687. [PubMed: 18339745]
- Yang J, Krishnamoorthy G, Saxena A, Zhang G, Shi J, Yang H, Delaloye K, Sept D, Cui J. An epilepsy/dyskinesia-associated mutation enhances BK channel activation by potentiating Ca²⁺ sensing. *Neuron.* 2010; 66:871–883. [PubMed: 20620873]
- Yuan P, Leonetti MD, Pico AR, Hsiung Y, Mackinnon R. Structure of the Human BK Channel Ca²⁺-Activation Apparatus at 3.0 Å Resolution. *Science.* 2010
- Zakharov, SI.; Wu, RS.; Liu, G.; Motoike, H.; Karlin, A.; Marx, SO. Locations of the Beta2 Transmembrane Helices in the BK potassium channel. 53rd Annual Biophysical Society Meeting; Boston, Massachusetts. 2009.




Figure 2. Differential modulation by $\beta 2ND$ on Ca^{2+} sensitivity of mSlo1 and dSlo1

A, Macroscopic currents of mSlo1 and dSlo1 with wt h $\beta 2$ at $\sim 100 \mu M [Ca^{2+}]_i$ (left). The voltage pulses are from -100 to +150 mV for 1s with 25 mV increments (shown only the first 20 ms), and the prepulse potential is -140 mV for 195 ms (shown only the last 5 ms). The dotted line represents the biexponential fit of the inactivation profile and the long dashed line represents the zero current line for each current trace. On the right, voltage dependence of the inactivation time constant. The fast and slow components of τ_{inact} were obtained by fitting current traces with a biexponential function from the peak amplitude to steady-state ($n = 10$ for mSlo1+wt $\beta 2$, except at +150 mV, $n = 9$; $n = 5$ for dSlo1+wt $\beta 2$, except at +150 mV, $n = 4$). **B,** Macroscopic currents of mSlo1 and dSlo1 with and without $\beta 2ND$ at $\sim 100 \mu M [Ca^{2+}]_i$. Voltage pulses are from -200 to 100 mV with 10 mV increments, and the repolarizing potential is -50 mV except for mSlo1+ $\beta 2ND$ that is -80 mV. The dashed line represents the zero current line for each current trace. **C,** Mean G-V relationship of mSlo1/dSlo1 with and without $\beta 2ND$ in 0, ~ 2 and $\sim 100 \mu M [Ca^{2+}]_i$, fitted with Boltzmann equation (smooth lines). The error bars in this and other figures show the standard error of the mean. **D,** $V_{1/2}$ vs. $[Ca^{2+}]_i$ plot of mSlo1/dSlo1 with and without $\beta 2ND$. The number of patches for each data set are: for the following $\sim (\mu M) [Ca^{2+}]_i$, 0, 1, 2, 5, 10, 30, 100, mSlo1: 118, 33, 29, 29, 42, 25, 56; mSlo1+ $\beta 2ND$: 44, 8, 8, 11, 9, 12, 28; dSlo1: 0, 0, 11, 19, 15, 9, 27; dSlo1+ $\beta 2ND$: 0, 0, 3, 4, 7, 8, 14.




Figure 3. Effects of chimeras of mSlo1/dSlo1 on the β2ND modulation of Ca^{2+} dependence
 The black bars show the change in Ca^{2+} sensitivity between with and without β2ND , normalized to mSlo1 (see Results for definitions). The gray bars show the Ca^{2+} response at $\sim 100 \mu\text{M } [\text{Ca}^{2+}]_i$ between with and without β2ND . The symbol * means $\sim 2 \mu\text{M } [\text{Ca}^{2+}]_i$ was used for comparison and ** mean the G-V could not be determined at 0 or $\sim 2 \mu\text{M } [\text{Ca}^{2+}]_i$. Bottom panel shows the schematic of different chimeras of mSlo1 and dSlo1 used in the top panel. The number of patches for each data are: for either $\sim 0/2$ and $100 \mu\text{M } [\text{Ca}^{2+}]_i$: C1: 3, 5; C1+ β2ND : 9, 11; C2: 4, 3; C2+ β2ND : 5, 4; C3: 7, 4; C3+ β2ND : 5, 5; C4: 5, 5; C4+ β2ND : 8, 3; C5: 7, 5; C5+ β2ND : 3, 4; m[dS0]: 9, 6; m[dS0]+ β2ND : 4, 7; m[dLinker+dAC]: 7, 20; m[dLinker+dAC]+ β2ND : 0, 20; m[dS0+dLinker+dAC]: 5, 14; m[dS0+dLinker+dAC]+ β2ND : 2, 9; d[mS0]: 3, 20; d[mS0]+ β2ND : 0, 12; d[mLinker+mAC]: 13, 7; d[mLinker+mAC]+ β2ND : 23, 13; d[mS0+mLinker+mRCK1c]: 11, 16; d[mS0+mLinker+mRCK1c]+ β2ND : 15, 14.




Figure 4. Effects of the N-terminus and the C-Linker/AC region of dSlo1 in mSlo1 background on the β 2ND modulation of Ca^{2+} sensitivity

A, Macroscopic currents of m[dS0], m[dLinker+dAC] and m[dS0+dLinker+dAC] with and without β 2ND at $\sim 100 \mu\text{M}$ $[\text{Ca}^{2+}]_i$. Voltage pulses are from -200 to 200 mV with 10 mV increments, and the repolarizing potentials is -50 mV except for m[dS0] and m[dLinker+dAC] with β 2ND, which is -80 mV. The dashed line represents the zero current line for each current trace. **B**, G-V relations of chimeras compared with that of mSlo1 (orange) at low (0 , ~ 2 or $\sim 10 \mu\text{M}$) and $\sim 100 \mu\text{M}$ $[\text{Ca}^{2+}]_i$. Due to apparent inactivation of the channels, the G-V relations were measured from both the peak currents and tail currents, and both methods resulted in the same $V_{1/2}$. **C**, $V_{1/2}$ vs. $[\text{Ca}^{2+}]_i$ plots of chimeras compared with that of mSlo1 and dSlo1 with and without β 2ND. The number of patches for each data are: for the following $\sim(\mu\text{M})$ $[\text{Ca}^{2+}]_i$, 0 , 1 , 2 , 5 , 10 , 30 , 100 , m[dS0]: 9 , 6 , 4 , 4 , 4 , 4 , 6 ; m[dS0] $+\beta$ 2ND: 4 , 3 , 3 , 4 , 3 , 5 , 7 ; m[dLinker+dAC]: 0 , 0 , 7 , 14 , 15 , 6 , 20 ; m[dLinker+dAC] $+\beta$ 2ND: 0 , 0 , 0 , 0 , 4 , 5 , 20 ; m[dS0+dLinker+dAC]: 0 , 0 , 5 , 4 , 4 , 4 , 14 ; m[dS0+dLinker+dAC] $+\beta$ 2ND: 0 , 0 , 2 , 3 , 4 , 2 , 9 .




Figure 5. Effects of the N-terminus and C-Linker/AC region of mSlo1 in dSlo1 background on the β 2ND modulation of Ca^{2+} sensitivity

A, Macroscopic currents of d[mS0], d[mLinker+mAC] and d[mS0+mLinker+mRCK1c] with and without β 2ND at ~100 μM $[\text{Ca}^{2+}]_i$. Voltage pulses are from -200 to 200 mV except for d[mS0+mLinker+mRCK1c]+ β 2ND, which are from -200 to 100 mV, with 10 mV increment. The repolarizing potential is -50 mV. The dashed line represents the zero current line for each current trace. **B**, G-V relations of chimeras with and without β 2ND at low (0 or ~2 μM) and ~100 μM $[\text{Ca}^{2+}]_i$ as compared with that of dSlo1 (blue). **C**, $V_{1/2}$ vs. $[\text{Ca}^{2+}]_i$ plots of chimeras compared with that of mSlo1 and dSlo1 with and without β 2ND. The number of patches for each data are: for the following $\sim(\mu\text{M})$ $[\text{Ca}^{2+}]_i$, 0, 1, 2, 5, 10, 30, 100, d[mS0]: 0, 0, 3, 10, 6, 4, 20; d[mS0]+ β 2ND: 0, 0, 0, 3, 2, 3, 16; d[mLinker+mAC]: 13, 5, 4, 4, 4, 4, 7; d[mLinker+mAC]+ β 2ND: 23, 5, 3, 3, 5, 5, 13; d[mS0+mLinker+mRCK1c]: 11, 2, 4, 7, 6, 16; d[mS0+mLinker+mRCK1c]+ β 2ND: 15, 5, 5, 5, 6, 14.




Figure 6. MWC model fittings of wt and chimera channels

G-V relationships (symbols) for mSlo1, dSlo1, m[dS0+dLinker+dAC] and d[mS0+mLinker+mRCK1c] with and without β 2ND fitted with the MWC model (lines). The $[Ca^{2+}]_i$ for each symbol is shown.




Figure 7. The role of the C-Linker and AC region of mSlo1 in the $\beta 1$ and $\beta 2ND$ modulation of Ca^{2+} sensitivity

A, G-V relations of mSlo1 and m[dLinker+dAC] with and without $\beta 1$ or $\beta 2ND$ in $\sim 100 \mu M$ $[Ca^{2+}]_i$. **B**, $V_{1/2}$ vs. $[Ca^{2+}]_i$ plots of mSlo1 and m[dLinker+dAC] with $\beta 1$ (left) or $\beta 2ND$ (right). The number of patches are: for the following $\sim (\mu M)$ $[Ca^{2+}]_i$, 0, 1, 2, 5, 10, 30, 100 : mSlo1+ $\beta 1$: 38, 13, 15, 11, 12, 8, 21; m[dLinker+dAC]+ $\beta 1$: 0, 0, 0, 10, 6, 9, 26.




Figure 8. Dependence of the $\beta 1$ and $\beta 2$ modulation of Ca^{2+} sensitivity on Ca^{2+} binding sites in mSlo1

A, Bar graph of the G-V shift in response to a $[\text{Ca}^{2+}]_i$ change from 0–100 μM of WT (wild type), single Ca^{2+} binding site mutations (D367A and 5D5N), double Ca^{2+} binding site mutation (D367A/5D5N) and double Ca^{2+} binding site plus the Mg^{2+} binding site mutation (D367A/5D5N/E399N) without (black) and with $\beta 1$ (unfilled) or $\beta 2\text{ND}$ (gray). **B**, Differences in $\Delta V_{1/2}$ from panel A of $\beta 1$ (unfilled) and $\beta 2\text{ND}$ (gray) with WT, single, double and triple binding site mutations. The number of patches for each data are: for ~ 0 and $\sim 100 \mu\text{M}$ $[\text{Ca}^{2+}]_i$: mD367A: 10, 9; mD367A+ $\beta 1$: 9, 9; mD367A+ $\beta 2\text{ND}$: 5, 7; 5D5N: 6, 11; 5D5N+ $\beta 1$: 8, 9; 5D5N+ $\beta 2\text{ND}$: 4, 8; mD367A/5D5N: 14, 6; mD367A/5D5N+ $\beta 1$: 6, 5; mD367A/5D5N+ $\beta 2\text{ND}$: 5, 5; mD367A/5D5N/E399N: 15, 15; mD367A/5D5N/E399N+ $\beta 1$: 5, 9; mD367A/5D5N/E399N+ $\beta 2\text{ND}$: 8, 12.




Figure 9. A structural model for the $\beta 2$ subunit modulation

Two opposing mSlo1 and $\beta 2$ (orange) subunits of BK channels are shown where cartoons for the VSD and S0 segment are constructed around the pore of the MthK channel (PDB ID: 1LNQ) (Jiang et al., 2002). The gating ring of the BK channel (PDB ID: 3MT5) (Yuan et al., 2010) is aligned to the MthK channel using UCSF Chimera v1.4.1. VMD v1.8.7 was used to show the aligned structure in cartoon and surface representation. The helices in the VSD are shown as cylinders and are positioned according to the $K_{V1.2}$ channel. The transmembrane (TM) segments of the $\beta 2$ subunit are positioned according to Zakharov et al (Zakharov et al., 2009). Green colored regions in the Slo1 subunit (S0, C-Linker and AC) are involved in the $\beta 2$ modulation of Ca^{2+} sensitivity.

Table 1

Residue information of mSlo1-dSlo1 chimeras

Chimera name	Residues from	
	dSlo1	mSlo1
C1	1–596	580–1169
C2	1–336	323–1169
C3	1–248	234–1169
C4	1–215	201–1169
C5	1–147	134–1169
m[dS0]	1–70	44–1169
m[dLinker+dAC]	340–432	1–325 & 419–1169
m[dS0+dLinker+dAC]	1–70 & 340–432	44–325 & 419–1169
d[mS0]	71–1164	1–43
d[mLinker+mAC]	1–339 & 433–1164	326–418
d[mS0+mLinker+mRCK1c]	71–346 & 559–1164	1–43 & 320–543

Table 2

Parameters for MWC model fits

	L_0	Z	K_c (μM)	K_o (μM)	$1/c$
mSlo1	9.20E+03	1.28	7.36	0.72	10.25
mSlo1+ β 2ND	9.41E+03	1.25	32.51	0.62	52.35
dSlo1	5.50E+05	0.88	22.00	0.51	43.48
dSlo1+ β 2ND	5.51E+05	0.78	23.95	0.53	45.28
m[dS0+dLinker+dAC]	9.98E+05	1.12	12.36	0.40	30.60
m[dS0+dLinker+dAC]+ β 2ND	9.80E+05	1.02	9.45	0.36	26.18
d[mS0+mLinker+miRCK 1c]	3.50E+04	1.07	30.76	3.30	9.32
d[mS0+mLinker+mRCK1c]+ β 2ND	7.90E+03	1.08	71.00	2.45	28.98