



Published in final edited form as:

J Neurosci. 2011 August 17; 31(33): 11825–11835. doi:10.1523/JNEUROSCI.1395-11.2011.

Fasting and 17 β -estradiol differentially modulate the M-current in NPY neurons

Troy A. Roepke¹, Jian Qiu¹, Arik W. Smith¹, Oline K. Rønnekleiv^{1,2}, and Martin J. Kelly^{1,2}

¹Department of Physiology and Pharmacology, Oregon Health and Science University, Portland, OR 97239

²Division of Neuroscience, Oregon National Primate Research Center, Beaverton, OR 97006

Abstract

Multiple K⁺ conductances are targets for many peripheral and central signals involved in the control of energy homeostasis. Potential K⁺ channel targets are the KCNQ subunits that form the channels underlying the M-current, a sub-threshold, non-inactivating K⁺ current that is a common target for G-protein coupled receptors. Whole-cell recordings were made from GFP (*Renilla*)-tagged NPY neurons from the arcuate nucleus of the hypothalamus using protocols to isolate and characterize the M-current in these orexigenic neurons. We recorded robust K⁺ currents in the voltage range of the M-current, which were inhibited by the selective KCNQ channel blocker XE991 (40 μ M), in both intact males and ovariectomized, 17 β -estradiol (E2)-treated females. Since NPY neurons are orexigenic and are active during fasting, the M-current was measured in fed and fasted male mice. Fasting attenuated the XE991-sensitive current by 3-fold which correlated with decreased expression of the KCNQ2 and KCNQ3 subunits as measured with quantitative real-time PCR. Furthermore, E2 treatment augmented the XE991-sensitive M-current by 3-fold in ovariectomized (vs. oil-treated) female mice. E2-treatment increased the expression of the KCNQ5 subunit in females but not KCNQ2 or KCNQ3 subunits. Fasting in females abrogated the effects of E2 on M-current activity, at least in part, by decreasing KCNQ2 and KCNQ3 expression. In summary, these data suggest that the M-current plays a pivotal role in the modulation of NPY neuronal excitability and may be an important cellular target for neurotransmitter and hormonal signals in the control of energy homeostasis in both males and females.

Keywords

Neuropeptide Y; fasting; 17 β -estradiol; KCNQ channels; XE991

Introduction

In the hypothalamic arcuate nucleus, two distinct, neuronal cell types differentially modulate energy homeostasis – the proopiomelanocortin (POMC) and neuropeptide Y (NPY) neurons. NPY neurons are orexigenic and respond to many peripheral and central hormonal signals that control energy homeostasis (Elmquist, 2001; Gao and Horvath, 2007; Kohno et al., 2007; Takahashi and Cone, 2005; Van den Pol et al., 2009). NPY neurons co-express Agouti-related Peptide, the endogenous antagonist to the melanocortin MC4 receptors, thus blocking the anorectic effects of the POMC peptide, α -MSH (Haskell-Luevano et al.,

2001;Lu et al., 2001). Ablation of NPY neurons in adults causes a rapid starvation (Gropp et al., 2005;Luquet et al., 2005) while icv injection of NPY into the paraventricular nucleus stimulates food intake (Stanley and Leibowitz, 1985). The anorectic, ovarian steroid 17 β -estradiol (E2) also suppresses NPY expression in the arcuate nucleus (Crowley et al., 1985;Pelletier et al., 2007) and may exert its effects on NPY neurons by modulating neuronal excitability.

NPY neurons are more excitable after fasting in a leptin-dependent manner (Takahashi and Cone, 2005). The activity of NPY neurons is modulated by peripheral signals (leptin, ghrelin, gastrin-releasing peptide) via targeting potassium and calcium channels (Van den Pol et al., 2009;Wang et al., 2008;Yang et al., 2010). These K⁺ channels hyperpolarize the membrane potential and inhibit neuronal excitability. Another K⁺ channel that modulates neuronal excitability, action potential kinetics and burst firing are the KCNQ channels (Delmas and Brown, 2005;Robbins, 2001). KCNQ channels constitute the M-current, a sub-threshold, non-inactivating, voltage-dependent, outward K⁺ current. The M-current, ubiquitously expressed in the brain and in neurons, is generated by expression of the KCNQ subunits, KCNQ2, -3, -4 and -5. Co-expression of KCNQ2/3 or KCNQ3/5 heteromultimers produces currents with similar kinetics to the native M-current (Robbins, 2001). Moreover, the M-current is a target of numerous neurotransmitters and neuropeptides that modulate neuronal excitability through G-protein coupled receptors including acetylcholine, serotonin, substance P and GnRH (Delmas and Brown, 2005;Xu et al., 2008). The role of the M-current in feeding behavior has, to our knowledge, not been reported.

Previously, we have shown that E2 (24 hr and long-term) increases the expression of KCNQ5 in guinea pig arcuate nucleus (Roepke et al., 2008;Roepke et al., 2007). Due to the heterogeneous nature of the arcuate nucleus, the exact identity of the neurons involved in the increase in KCNQ5 expression was unknown. However, since E2 reduces feeding and NPY expression (reviewed in Roepke, 2009) and the M-current dampens neuronal excitability, we hypothesized that an increase in M-current activity specifically in NPY neurons is a cellular mechanism employed by E2 to reduce food intake. Furthermore, the effects of fasting, which are known to increase NPY excitability, may attenuate the expression and activity of the M-current. Therefore, in this study, we examined the effects of fasting and E2 treatment on the expression and activity of the M-current using whole-cell patch clamp recordings, single-cell harvesting and quantitative real-time PCR (qPCR).

Materials & Methods

Animal care and experimental treatments

All animal treatments are in accordance with institutional guidelines based on National Institutes of Health standards, and were performed with Institutional Animal Care and Use Committee approval at the Oregon Health & Science University. Female and male GFP-NPY transgenic mice (provided by Dr. Brad Lowell, Harvard University)(Van den Pol et al., 2009) were selectively bred in-house, and maintained under controlled temperature (25 °C) and photoperiod conditions (12/12 hr light/dark cycle) with food and water *ad libitum*. Male mice were divided into three groups: fed, 24 hr fasted and 24 hr re-fed. The re-fed treatment entailed a 24 h fast followed by a 24 h re-feeding prior to experimentation. A 24 h fast has previously been reported to alter the neuronal excitability of the NPY neurons (Takahashi and Cone, 2005). NPY females were ovariectomized under isoflurane anesthesia, implanted with sterile capsules containing sesame oil or 35 μ l of 180 μ g/ml 17 β -estradiol (E2) in sesame oil and allowed to recover for 1 week. This E2 treatment produces low, diestrus levels of E2 (~30 pg/ml). After one week, the females were injected s.c. with either oil or 1 μ g of 17 β -estradiol benzoate (E2B) 24 h prior to experimentation. The estrogen replacement paradigm has been previously used to mimic the endogenous levels of E2 during proestrus

(average serum E2 levels = 77.6 ± 10.7 pg/ml (n=9) in our animals) (Bronson and Vom Saal FS, 1979;Zhang et al., 2009). A subset of E2-treated females was also subjected to a 24 hr fast starting at the time of injection. Average weight loss after a 24 hr fast was 2.9 ± 0.4 g (n=10) for females and 3.5 ± 0.2 g (n=21) for males. Refed males recovered on average 90% of their pre-fast weight. Fasting had no effect on animal activity, fur condition and response to handling.

Drugs

17 β -estradiol (E2) and 17 β -estradiol benzoate (E2B) was purchased from Steraloids (Newport, RI, USA) and dissolved in ethanol prior to oil dissolution. XE991 (10,10-bis(4-Pyridinylmethyl)-9(10H)-anthracenone dihydrochloride), D-AP5 (2R)-amino-5-phosphonovaleric acid), and CNQX (6-cyano-7-nitroquinoxaline-2,3-dione) were all purchased from Tocris (Ellisville, MO, USA). Bicuculline (1(S),9(R)-(-)-Bicuculline methiodide) was purchased from Sigma-Aldrich. Tetrodotoxin (TTX) was purchased from Alomone Laboratories, Inc. (Jerusalem, Israel) and dissolved in H₂O.

Preparation of basal hypothalamic slices

Slices were prepared as described previously (Ibrahim et al., 2003;Qiu et al., 2003;Qiu et al., 2010). Transgenic GFP-NPY mice were killed quickly by decapitation at 1000–1100 h. The brain was rapidly removed from the skull and a block containing the basal hypothalamus (BH) was immediately dissected. The BH block was submerged in cold (4 °C) oxygenated (95% O₂, 5% CO₂) high-sucrose cerebrospinal fluid (CSF; in mmol): 208 sucrose; 2 KCl; 26 NaHCO₃; 10 glucose; 1.25 NaH₂PO₄; 2 MgSO₄; 1 MgCl₂; 10 HEPES, pH 7.3; 300 mOsm. Coronal slices (250 μ m) from the BH were cut on a vibratome during which time (10 min) the slices were bathed in high-sucrose CSF at 4 °C. The slices were then transferred to an auxiliary chamber in which they were kept at room temperature (25 C) in artificial CSF (aCSF) consisting of (in mM): 124 NaCl; 5 KCl; 2.6 NaH₂PO₄; 2 MgCl₂; 2 CaCl₂; 26 NaHCO₃; 10 glucose, pH 7.3; 310 mOsm until recording (recovery for 2 h). A single slice was transferred to the recording chamber mounted on an Olympus BX51W1 upright microscope equipped with video-enhanced, infrared-differential interference contrast (DIC) and Exfo X-Cite 120 Series fluorescence light source (Mississauga, Ontario, CANADA). The slice was continually perfused with warm (35 °C), oxygenated aCSF at 1.5 ml/min. Targeted neurons were viewed using both IR-DIC and blue excitation light with an Olympus 40 \times water-immersion lens.

Visualized whole-cell patch recording

Normal aCSF and pipette solutions were used in electrophysiological recording (Ibrahim et al., 2003;Qiu et al., 2003;Qiu et al., 2010). Standard whole-cell patch recording procedures and pharmacological testing were as previously described (Qiu et al., 2010;Xu et al., 2008). Whole-cell voltage and current clamp recordings were performed using pipettes made of borosilicate glass and pulled using a P-97 Flaming/Brown Micropipette Puller (Sutter Instrument, Novato, CA, USA). Pipettes were filled with normal internal solution (in mM: 10 NaCl; 128 K-gluconate; 1 MgCl₂; 10 HEPES; 1 ATP; 11 EGTA; 0.25 GTP, pH 7.3; 300 mOsm) with a 3–5 M Ω resistance. An Axopatch 200A amplifier, Digidata 1322A Data Acquisition System and pCLAMP software (version 9.2, Molecular Devices, USA) were used for data acquisition and analysis. Input resistance, series resistance and membrane capacitance were monitored throughout the experiments. Only cells with stable series resistance (<30 M Ω , <20% change) and an input resistance >500 M Ω were used for analysis. The access resistance was 80% compensated and the calculated liquid junction potential (~10 mV) was corrected. To display reversal potential and rectification characteristics of the ligand-activated currents, I–V plots constructed by voltage steps from –50 to –140 mV at 10 mV increments applied at 1 s intervals from a holding potential of

–60 mV. From this protocol, the input resistance was determined from the I–V plot as the ratio of the voltage (–60 to –80 mV)/current (pA). To calculate the rheobase and elicit an action potential in current clamp, the resting membrane potential was held to –80 mV for all neurons with a small hyperpolarizing current, followed by injection of incremental current steps (~6 pA/step) for 500 ms before and after bathing the slice in XE991 (40 μ M). In voltage clamp, two protocols were employed to measure the deactivation and steady-state currents. The deactivation protocol measured the currents elicited during 500 ms voltage steps from –25 to –75 mV after a 300 ms prepulse to –20 mV, which included the membrane potential at which the maximal M-current could be obtained (Schweitzer, 2000; Xu et al., 2008). The amplitude of M-current relaxation or deactivation was measured as the difference between the initial (<10 ms) and sustained current (>475 ms) of the current trace in the control conditions (TTX only, 1 μ M, 5 min) minus the difference in the XE991 (40 μ M, +TTX, 10 min) conditions. To examine the steady-state current, a series of command voltage steps from –80 to +10 mV in 10 mV increments (1 s) were applied from a holding potential of –90 mV (500 ms) and returning to –60 mV holding potential (Leao et al., 2009). The XE991-sensitive current was determined by subtracting the steady state (SS) current (the last 50 ms of the voltage steps) of the whole K⁺ currents in the XE991-treated traces from steady state current in the control traces. Both voltage clamp protocols were repeated twice for each bath solution and averaged for analysis.

Cell harvesting of dispersed mouse NPY neurons, single cell RT-PCR and quantitative real-time PCR

These procedures were conducted as described previously (Ibrahim et al., 2003; Xu et al., 2008; Zhang et al., 2008; Zhang et al., 2009). Briefly, four 250 μ m basal hypothalamic slices were cut on a vibratome and placed in an auxiliary chamber containing oxygenated aCSF. The slices were allowed to recover for 1–2 hrs in the auxiliary chamber. Thereafter, the arcuate nucleus and median eminence was microdissected and incubated in 10 ml aCSF (pH 7.3, 300 mOsm) containing 1 mg/ml protease for ~15 minutes at 37° C. The tissue was then washed four times in low calcium CSF (1 mM CaCl₂) and two times in aCSF. The cells were isolated by trituration with flame-polished Pasteur pipettes, dispersed onto a 60 mm glass-bottomed Petri dish and perfused continuously with aCSF at a rate of 2.0 ml/min. The dispersed, bi-polar or uni-polar fluorescent cells were visualized using a Leitz inverted fluorescent microscope, patched and then harvested with gentle suction to the pipette using the Xenoworks manipulator system (Sutter Instruments, Novato, CA) to the pipette and expelled into siliconized microcentrifuge tubes. Each harvested cell or pools of cells were heat denatured and reverse transcribed as described previously (Zhang et al., 2008; Zhang et al., 2009). For quantitative PCR, six to eight pools of 5 cells (KCNQ2 & 3) or 10 cells (KCNQ5) each were harvested from individual ovariectomized, oil- and E2-treated (both fed and fasted) females or fed and fasted males (age 10–14 weeks). Perfused CSF was regularly collected along with the harvested cells and processed together with the cells and control samples. Single cell, and tissue RNA used as negative controls, were processed similarly but without reverse transcriptase (RT).

The single cell RT-PCR primers for KCNQ2 and KCNQ4 were as previously published (Xu et al., 2008). The single-cell RT-PCR primers for KCNQ3, KCNQ5 and NPY were as follows: KCNQ3 (94 bp product, accession number NM_152923.1, forward primer: GCTGCTGGAAACCTTTGC, reverse primer: ACGCCAGCCTTTGTATCG, 474–567 nt, annealing temperature: 60 °C); KCNQ5 (143 bp product, accession number NM_023872.2, forward primer: GATGGCAAGGAAGCCTGAG, reverse primer: TAGGAACCGGAGACTTCTG, 572–714 nt, annealing temperature: 57 °C) and NPY (182 bp product, accession number NM_023456, forward primer: ACTGACCCTCGCTCTATCTC, reverse primer: TCTCAGGGCTGGATCTCTTG, 106–

287 nt, annealing temperature: 64 °C). Each reaction was amplified for 50 cycles using a BIO-RAD C1000 Thermal Cycler (Bio-Rad, Hercules, CA). All scRT-PCR products were confirmed by sequencing. Negative (cell and tissue samples without RT and harvested CSF) and positive tissue controls were used for each PCR run. KCNQ subunit co-localization patterns were determined by counting some cells more than once e.g., each cell that had all three subunits also were included with those that co-expressed only KCNQ2+3, KCNQ3+5, or KCNQ2+5.

To determine the changes in KCNQ gene expression between fed and fasted males, and oil- and E2-treated females, pools of dispersed GFP-NPY neurons were harvested, reverse transcribed and analyzed using quantitative real-time PCR. Due to the technical limitations of single cell expression analysis of low-abundant genes, we could not measure KCNQ subunit transcripts in individual neurons reliably with quantitative real-time PCR. Therefore, pooling of neurons was required especially when targeting low-expression genes such as cation channel subunits. This has allowed us to do careful quantitative analysis of mRNA expression that often correlates with the whole-cell current activity (Zhang et al., 2009) as in the current study. The procedures for quantitative real-time PCR were previously described (Zhang et al., 2009). For KCNQ2 and KCNQ3, pools of 5 NPY neurons were sufficient while pools of 10 NPY neurons were needed for the analysis of KCNQ5 gene expression, which was most likely due to lower expression of the KCNQ5 subunit in NPY neurons or the fewer number of neurons that expressed this subunit. Negative (cell and tissue samples without RT and harvested CSF) and positive tissue controls were used for each single cell and quantitative PCR run. The primers for quantitative real time PCR for KCNQ3 were previously described above. For KCNQ2 and KCNQ5, the primers were as follow: KCNQ2 (171 bp product, accession number NM_133322, forward primer: GGTGCTGATTGCCTCCATTG, reverse: TCCTTGCTGTGAGCGTAGAC, 644–814 nt, annealing temperature: 60 °C); KCNQ5 (99 bp product, accession number NM_023872.2, forward primer: GGCACAATCACACTGACAAC, reverse primer: GAAATGCCAAGGAGTGCGAAG, 915–1013 nt, annealing temperature: 58 °C). The relative standard curve model was used to determine amplification efficiency using serial dilutions of basal hypothalamic cDNA. The calculated efficiencies for all four primers were >94%. The small difference between the efficiencies (less than 5%) does not affect the analysis of relative gene expression (Livak and Schmittgen, 2001; Pfaffl, 2001; Schmittgen and Livak, 2008). Relative KCNQ5 expression was also analyzed using quantitative real-time PCR in ovariectomized, oil- and E2-treated wild-type (C57BL/6) and α ERKO mice (C57BL/6 from Professor Pierre Chambon, CNRS/INSERM, Illkirch-Cedex, France).

To determine which estrogen receptor (ER) was expressed in NPY neurons, we also analyzed the expression of ER α and ER β in NPY pools (5 cells) from ovariectomized, female mice. The primers for ERs were as follows: ER α (107 bp product, accession number NM_007956, forward primer: GCGCAAGTGTTACGAAGTG, reverse primer: TTCCGGCCTTCCAAGTCATC, 919–1025 nt, annealing temperature: 60 °C) and ER β (113 bp product, accession number NM_010157, forward primer: AATGTCCACCCGCTAGGCATTTC, reverse primer: CTCCATGTCTTGCGTAGGTCTC, 298–410 nt, annealing temperature: 60 °C). The reference gene for all quantitative real-time PCR was β -actin as previously published (Zhang et al., 2009). All quantitative real-time PCR was conducted using the Power SyberGreen master mix on an ABI 7500 Fast Real-Time PCR System (Life Technologies Corp., Carlsbad, CA, USA).

Co-localization of NPY and ER α using immunohistochemistry

We performed immunohistochemistry to document ER α expression in mouse NPY neurons. Four ovariectomized female GFP-NPY mice were sedated by ketamine injection and decapitated 7–10 days post-ovariectomy. The brains were quickly extirpated and cooled in

aCSF (4 °C) for 1 min. One-mm blocks of the basal hypothalamus were sliced, fixed in 4% paraformaldehyde for 6 h with gentle shaking on ice and then cryopreserved in 20% sucrose overnight in the refrigerator. Tissue blocks were frozen at -55 °C, coronally sectioned on a cryostat at 20 µm and thaw-mounted on Superfrost Plus glass slides (Fisher Scientific, Pittsburgh, PA). Sections were rinsed in PBS (0.1 M phosphate buffer, pH 7.4, and 0.15 M NaCl; all rinses were in PBS for 30 min unless noted) and then incubated for 24 h at 4 °C in antiserum against ERα (1:10,000; C1355; Upstate; Lake Placid, NY). After rinsing, sections were incubated for 1 h at room temperature in biotinylated goat anti-rabbit IgG (1:1,000; Vector Laboratories; Burlingame, CA), washed and then incubated in Cy3-conjugated streptavidin (1:1,000; Jackson ImmunoResearch; West Grove, PA). Following a final rinse for 3 h, slides were coverslipped using a glycerol-glycine buffer (2:1, pH 8.6) containing 5% *n*-propyl gallate to reduce photobleaching (Giloh and Sedat, 1982). Both the primary and secondary antisera and tertiary marker were diluted in Tris-(hydroxymethyl)aminomethane (0.5%; Sigma Chemical Co., St. Louis, MO) in PBS containing 0.7% seaweed gelatin (Sigma), 0.4% Triton X-100 (Sigma) and 3% BSA (Sigma) adjusted to pH 7.6. Slides were analyzed and photographed using Nikon E800 microscope equipped with epifluorescence and a DS digital camera. A total of 10–12 arcuate sections from each female were examined. The number of GFP-NPY cells co-localizing ERα was counted per section.

Data analysis

Comparisons of the I–V plots between different feeding states and steroid treatments were performed using a two-way ANOVA analysis with the Bonferroni-Dunn *post hoc* test. Rheobase and quantitative real-time PCR results were analyzed by unpaired Student's *t*-test. Differences were considered statistically significant if the probability of error was <5%. All data are presented as mean ± SEM.

Results

Feeding states modulate the activity of the M-current in males

We initially determined the basal activity of the NPY neurons and the role of the M-current in modulating neuronal excitability in fed male mice. In current clamp, action potentials were evoked by the injection of incremental current steps (~6 pA/step) before and after the application of XE991 (40 µM), a selective blocker of KCNQ channels (Fig. 1A). Under control conditions, no action potentials were evoked after the first current injection of 6 pA (gray trace); however, the first current injection did produce an action potential after treatment with XE991. The first action potential evoked under control conditions (black trace) was after ~30 pA of current which produced two action potentials (or more in other cells) in the XE991 conditions. Therefore, the average rheobase was significantly reduced by XE991 from a mean of 27.4 ± 1.2 pA to 18 ± 0.9 pA ($n=15$, $p < 0.01$) (Fig. 1B). During a 10 min perfusion of XE991, the resting membrane potential (RMP) of each neuron was continuously recorded in current clamp. The average RMP of NPY neurons from the fed males was -73.9 ± 0.9 mV, and after a 10 min XE991 perfusion the membrane depolarized to -57.3 ± 1.1 mV ($p < 0.0001$). XE991 application also increased the input resistance in the fed males from 0.75 ± 0.05 GΩ to 1.43 ± 0.27 GΩ, ($p < 0.01$) indicating a reduction in membrane conductance. The majority of the neurons (10 of 15) were quiescent prior to XE991 application (Fig. 1D), yet became active after 8–9 minutes of XE991 perfusion. The remaining neurons were active but depolarized (and increased firing) after XE991. To determine if action potential generation during XE991 perfusion was dependent on presynaptic inputs (glutamatergic, GABAergic), we repeated these experiments in the presence of glutamate receptor (CNQX, AP5) and GABA-A receptor (bicuculline) blockers (Figure 1D). Blockade of the M-current by XE991 in NPY neurons did not increase firing, although the neurons were significantly depolarized from baseline to a similar level of

depolarization (14.9 ± 3.2 mV ($n=3$) with blockers compared to 16.8 ± 1.3 mV ($n=15$) without blockers). The level of depolarization did not reach the action potential threshold for hypothalamic arcuate neurons (Kelly et al., 1990; Loose et al., 1990) to fire independently of excitatory inputs. However, the neurons treated with the synaptic blockers did fire when sufficient depolarizing current was applied, and the response was greater in the XE-treated neurons (Figure 1E). Therefore, the data suggests that the M-current in NPY neurons contributes to the control of neuronal excitability and therefore is a potential target for modulation during different physiological states.

To measure the M-current in voltage clamp, we measured the deactivation or relaxation of the whole-cell K^+ currents elicited by an established protocol (Schweitzer, 2000; Xu et al., 2008) before and after application of XE991 ($40 \mu\text{M}$) in the presence of TTX ($1 \mu\text{M}$). The XE991-sensitive current (M-current) was calculated by subtracting the current relaxation (the difference between the instantaneous and steady state (arrows) Figure 2A) in XE991 from the control conditions. Because the M-current has been shown to decrease or “rundown” after cell dialysis in whole-cell recordings in other neuronal cell types (Shen et al., 2005), the whole cell K^+ currents evoked by the deactivation protocol were monitored over 30 minutes to determine the change in the relaxation currents (Figure 2B). In NPY neurons from fed males, the outward K^+ currents evoked by the protocol did not significantly decrease over the 30 min, which is beyond the experimental time needed for each recording, yet the KCNQ channel blocker XE991 was fully efficacious over this time period.

To determine the appropriate concentration to use in the bath perfusion, a sequential concentration-response was performed using $1 \mu\text{M}$, $10 \mu\text{M}$ and $40 \mu\text{M}$ of XE991 in five NPY neurons from fed males (Figure 2C). The lowest concentration of XE991 did not significantly inhibit the K^+ current evoked by the deactivation protocol, while $10 \mu\text{M}$ did significantly attenuate the current at the highest voltages (-35 & -30 mV). At $40 \mu\text{M}$, XE991 significantly attenuated the current at voltages more depolarized than -45 mV. The effects of XE991 at this concentration were not fully reversible within the experimental time period (>30 min, data not shown). However, this concentration ensured that all possible homomultimers and heteromultimers of KCNQ channel subunits (KCNQ2, -3, & -5), which have different sensitivity to XE991 (Robbins, 2001), were inhibited. For example, in cell expression systems the IC_{50} of XE991 for KCNQ5 homomultimers is approximately $65 \mu\text{M}$ (Schroeder et al., 2000), while the IC_{50} for KCNQ2/3 heteromultimers is $0.6 \mu\text{M}$ (Wang et al., 1998). Higher concentrations of XE991 (greater than $100 \mu\text{M}$) can block ERG (Ether-a-go-go Related Gene) channels (Elmehrik et al., 2007) and may affect Kv4.x (A-type) channels (Wang et al., 1998); although similar to other investigators (van den Pol et al., 2009), we did not measure a robust A-current in NPY neurons (data not shown). Therefore, the $40 \mu\text{M}$ concentration of XE991 was used for all subsequent experiments to ensure the maximum amount of KCNQ blockade without alteration of other K^+ currents.

Because NPY neurons play a pivotal role in the hypothalamic control of feeding behaviors, we determined the effects of fasting and re-feeding on the M-current in NPY neurons from adult male mice in the presence of $1 \mu\text{M}$ TTX to block Na^+ - spike dependent synaptic inputs. The deactivation protocol examines the voltage range (-30 to -75 mV) where the M-current has its most profound effect on excitability. In males, the XE991-sensitive current (the M-current) was significantly attenuated by a 24 hr fasting ($p < 0.01$, Fig. 2D). A 24 hr re-feeding restored a significant portion of the M-current in the NPY neurons. A 2 hr re-feeding had no effect on the M-current in fasted males (data not shown). The maximum XE-sensitive current was recorded at -35 mV (fed males = 43.6 ± 5.4 pA ($n=11$); fasted = 14.8 ± 3.3 pA ($n=17$); re-fed = 35.7 ± 2.6 pA ($n=10$), Fig. 2D). In addition, the RMP between fed and fasted males was significantly different (fed, -71.5 ± 1.6 mV vs. fasted, -64.8 ± 1.6

mV, $p < 0.01$) while the capacitance was not significantly different (fed: 17.9 ± 2.0 pF; fasted: 16.4 ± 0.7 pF). XE991 application significantly reduced the RMP in both states but only significantly increased the input resistance in fed males. The RMP in re-fed males was not significantly different from either of the other treatments, but XE991 significantly increased the input resistance (Table 1), which correlates with the increased conductance due to the M-current in the fed and re-fed states compared to the fasted state (Figure 2D).

To determine the steady-state conductance and kinetics of the M-current in NPY neurons, we used an established activation protocol (Leao et al., 2009) in the presence of $1 \mu\text{M}$ TTX (Fig. 3A). The steady state currents at the higher voltages were significantly higher in the fed and re-fed males compared to the fasted males ($p < 0.001$). The XE-sensitive current at $+10$ mV was 484 ± 37.6 pA ($n=11$) for fed males, 259 ± 63.3 pA ($n=15$) for fasted males, and 480 ± 53.6 pA ($n=10$) for re-fed males (Fig. 3B). The percentage of steady state current at $+10$ mV that was XE991-sensitive was $51.6 \pm 5.2\%$ in the fed state, $29.3 \pm 3.7\%$ in the fasted state and $38.7 \pm 3.3\%$ in the re-fed state (fed vs fasted, $p < 0.01$). The M-current is active at sub-threshold potentials and increases in amplitude around -50 to -40 mV (Robbins, 2001). The amplitude of the currents evoked by the activation protocol at these voltages was similar to the amplitude of the currents evoked by the deactivation protocol for all treatments (> -50 mV). A Boltzmann equation fit to the G/G_{max} plot (G_{max} at $+10$ mV) yielded a $V_{1/2}$ for activation of the M-current of -22.9 ± 2.4 mV in the fasted state. This $V_{1/2}$ is well within the range of the $V_{1/2}$ for M-currents expressed in heterologous cell systems (Robbins, 2001). Since the XE-sensitive current did not plateau at the maximum test potential of $+10$ mV, a Boltzmann fit was not done for the fed or re-fed states.

To determine if changes in the K^+ conductance were due mainly to the M-current and not other K^+ currents, the total conductances and the XE991-sensitive and XE991-insensitive currents were compared between each feeding state (Fig. 3C). The differences in the total conductance were directly correlated to the decrease in XE991-sensitive conductance in the fasted state. The ratio of time-dependent current (Inst-SS) to the steady state current (SS) at $+10$ mV was calculated to determine the relative effect of XE991 on outward currents between the feeding states (Fig. 3D). The ratio at the $+10$ mV step was significantly higher after XE991-treated compared to control in the fasted state ($p < 0.001$) indicating that the reduction in the control currents was mostly due to the decrease in the steady state M-current. Conversely, in the fed and re-fed males, blocking the M-current with XE991 did not significantly affect the ratio.

E2 increases activity of the M-current in ovariectomized females

Because E2 suppresses food intake and NPY neurons are orexigenic, we hypothesized that E2 may suppress feeding behavior by increasing the KCNQ5 expression and activity of the M-current in NPY neurons. To initially test this hypothesis, we used the deactivation protocol described in Figure 2 in the presence of $1 \mu\text{M}$ TTX. In oil-treated, ovariectomized females ($n=14$), the M-current (XE991-sensitive) was significantly less than in E2-treated, ovariectomized females ($n=15$, $p < 0.01$, Fig. 4). The relaxation current at -35 mV was 15.2 ± 2.5 pA and 42.5 ± 3.9 pA for oil-treated and E2-treated females, respectively (Fig. 4, $p < 0.001$). The RMP between oil- and E2-treated females was significantly different (-68.1 ± 1.8 mV vs. -72.4 ± 1.1 mV, $p < 0.05$) while the capacitance between the treatments was not significantly different (oil-treated: 18.5 ± 1.3 pF; E2-treated: 16.1 ± 0.9 pF). XE991 application significantly reduced the RMP in both treatments but only significantly increased input resistance in E2-treated females (Table 1), again indicating that the M-current contributed a significant portion of the resting conductance in the E2-treated NPY neurons. Interestingly, fasting (24 hr) abrogated the E2-induced potentiation of the M-current and significantly reduced the relaxation current to 14.7 ± 2.6 pA ($n=11$, $p < 0.001$) (Figure 4). The RMP in NPY neurons from fasted, E2-treated females was significantly

lower (-65.6 ± 1.5 mV) than in fed, E2-treated females. XE991 depolarized NPY neurons but did not significantly increase the input resistance in the fasted, E2-treated females (See Table 1).

To determine if E2 alters the steady-state conductances and kinetics of the M-current in NPY neurons, we used the activation protocol described in Figure 3. In the presence of 1 μ M TTX, the steady state XE991-sensitive currents were significantly greater in fed, E2-treated females ($n=12$) compared to both oil-treated ($n=14$) and fasted, E2-treated females ($n=11$; $p < 0.001$, Figure 5A). The current at +10 mV for the three treatments was 231 ± 24 pA for oil-treated females; 513 ± 82 pA for fed, E2-treated females; and 250 ± 31 pA for fasted, E2-treated females ($p < 0.001$, oil and fasted E2 compared to E2). The current at +10 mV for the fed, E2-treated, ovariectomized females was not significantly different from that of the fed, intact males. The percentage of steady state current at +10 mV that was XE991-sensitive was $28.9 \pm 2.8\%$ in oil-treated females, $44.1 \pm 3.1\%$ in fed, E2-treated females and $23.1 \pm 1.9\%$ in fasted, E2-treated females (oil vs. fed E2, $p < 0.01$; fed E2 vs. fasted E2, $p < 0.001$). As in the males, the M-current is active at sub-threshold potentials and increases significantly in amplitude at -50 to -40 mV. Using G/G_{\max} and a Boltzmann equation fit, the $V_{1/2}$ of the M-current in the oil-treated females was -22.2 ± 1.6 mV. Since the XE-sensitive current did not plateau at the maximum test potential of +10 mV, a Boltzmann fit was not done for the fed or fasted, E2-treated females.

As in the males, the total conductance and the XE991-sensitive and XE991-insensitive current were compared between steroid treatments and feeding states to determine if the changes in total conductance were due solely to the M-current (Fig. 5B). The E2-treated, XE991-sensitive conductances were significantly greater than the oil-treated or fasted, E2-treated ($p < 0.001$). However, unlike the males, the total conductances were not significantly different, which suggests that the contribution of the M-current to the total conductance is smaller in female NPY neurons compared to male NPY neurons. The ratio of time-dependent current (Inst-SS) to the steady state current (SS) at +10 mV was significantly different between control and XE991-treated conditions in both oil-treated and fed, E2-treated females ($p < 0.001$, Fig. 5C).

KCNQ expression in mouse NPY neurons

Previously, we have shown that KCNQ 2, 3, and 5 are all expressed in guinea pig NPY neurons (Roepke et al., 2007). Based on the present electrophysiological findings, we hypothesized that all three subunits would also be expressed in mouse NPY neurons. Using single cell RT-PCR collected from 8 ovariectomized females, the average percent expression was $72 \pm 2\%$; $65 \pm 2\%$, and $48 \pm 3\%$ for KCNQ2, KCNQ3 and KCNQ5, respectively. A representative co-expression pattern is illustrated in Figure 6A ($n=8$ animals, average 15 cells/animal, 110 total cells). Only GFP cells positive for NPY mRNA were used in the analysis and 99% of all GFP-labeled arcuate neurons were NPY neurons. The co-expression pattern of the three KCNQ subunits showed that a majority (~65%) of NPY neurons express more than one subunit. Amongst the harvested NPY neurons, 37% co-expressed KCNQ2 & -3; 33% co-expressed KCNQ2 & -5; and 25% co-expressed KCNQ3 & -5 (Figure 6B). Amongst the 110 NPY neurons, 15% co-expressed all three subunits; 50% expressed two transcripts; 32% expressed only one and the remaining 3% did not express any KCNQ transcripts. The majority of the cells expressing only one subunit expressed KCNQ2 or KCNQ3. One or more KCNQ subunits were detected in 97% of NPY neurons by single cell RT-PCR, and the whole-cell recording of the M-current revealed that the channels are expressed in the vast majority of cells. All three KCNQ subunits may be expressed in the majority of cells but at levels beyond the sensitivity of the single cell RT-PCR and therefore cannot be coupled directly to the electrophysiological analysis. We and

others have previously discussed this technical problem with other channels expressed in CNS neurons (Tkatch et al., 2000; Zhang et al., 2007).

Differentially regulation of KCNQ channel subunits by fasting and E2

To determine if regulation of KCNQ subunit gene expression is, at least, partially responsible for the reduction of the M-current, we examined the relative mRNA expression of KCNQ2, KCNQ3 and KCNQ5 (Figure 7A, 7B and 7C, respectively) in pooled NPY neurons from fed and fasted males using quantitative real-time PCR. The expression of KCNQ2 and KCNQ3 in fasted males was significantly reduced compared to fed males (Fig. 7D, $n=5$, $p < 0.01$) without any effect on KCNQ5 expression (Fig. 7D). The regulation of KCNQ channel mRNA expression and whole-cell M-current by fasting indicates that the M-current is an important neuronal current in the control of energy homeostasis.

Previously, we have shown that estrogen replacement in ovariectomized female guinea pigs increased the expression of KCNQ5 in the arcuate nucleus after both 24 hr and long-term treatment (Roepke et al., 2008; Roepke et al., 2007). To determine if regulation of KCNQ subunit gene expression is, at least, partially responsible for the increase in M-current activity in female arcuate NPY neurons, we also examined the relative mRNA expression of KCNQ subunits in pooled NPY neurons from oil- and E2-treated females using quantitative real-time PCR. The expression of KCNQ2 and KCNQ3 was not affected by E2 treatment while KCNQ5 was significantly increased by E2 treatment (Figure 7E; $n=5$, $p < 0.05$). As in the males (Fig. 7D), fasting significantly reduced the expression of KCNQ2 and KCNQ3 in NPY neurons from E2-treated females as compared to fed, E2-treated females (Fig. 7E, $n=5$, $p < 0.01$) but did not affect the elevated KCNQ5 expression observed in fed, E2-treated females (Fig. 7E, $n=5$, $p < 0.05$ compared to oil-treated). The increase in M-current activity after E2 replacement indicates that the M-current has a role in mediating the effects of E2 on NPY neuronal activity, which may be, in part, due to the up-regulation of KCNQ5 expression.

ER α expression and co-localization with KCNQ5 in NPY neurons

Because E2 increased the expression of KCNQ5 in NPY neurons, we wanted to determine if this was dependent on ER α . Tissue RNA was extracted from the microdissected arcuate nucleus obtained from WT and ER α KO oil- and E2-treated mice. As found in NPY neurons, E2 replacement significantly increased the mRNA expression of KCNQ5 in RNA extracted from the arcuate nucleus dissected from wild-type females but failed to alter KCNQ5 expression in ER α KO females (Figure 8A). Because E2 regulates KCNQ5 and NPY gene expression in the arcuate (Roepke, 2009; Roepke et al., 2007) but fails to increase KCNQ5 expression in NPY neurons in ER α KO mice, we determined the expression of ER α and ER β in single and pooled NPY neurons collected from ovariectomized females. In pools of NPY neurons ($n=12$, 5 cells per pool), ER α mRNA was detected in every pool using quantitative real-time PCR with an average C_T value of 33.6 ± 0.19 and a single product at 80.5 °C. ER β was not detected in any NPY pools (data not shown). Using single-cell RT-PCR, we found that ER α mRNA was expressed in 19% (23 of 122; average 18 cells/animal ($n=7$)) of NPY neurons while ER β was not expressed in any of these neurons (Figure 8B). Interestingly, 50% of ER α -positive NPY neurons co-expressed KCNQ5 mRNA, but only 18% of KCNQ5-positive NPY neurons co-expressed ER α . In contrast to mRNA expression, immunocytochemistry detected only a small percentage (<4%) of NPY neurons that co-localized ER α (Figure 8C), which is similar to a previous report (Simonian et al., 1999).

Discussion

For the first time, we show that energy states (fasting) regulate the expression and activity of the M-current in hypothalamic NPY neurons. In fasted males, the M-current was suppressed by ~3-fold in NPY neurons compared to fed males, in part, due to a reduction in KCNQ2/3 channel expression. This study is also the first to characterize the effects of E2 on M-current activity in a native neuronal cell type. E2 replacement in ovariectomized females potentiated the M-current compared to oil-treated females, in part, by increasing the mRNA expression of the KCNQ5 subunit. Fasting abrogated the effects of E2 on the M-current indicating that the activity of the M-current is critical for the control of food intake via NPY neurons.

The M-current controls resting membrane potential and is suppressed during fasting in NPY neurons

Arcuate NPY neurons are involved in the control of food intake and are responsive to multiple central and peripheral signals that are responsive to satiety and fasting (reviewed in (Gao and Horvath, 2007; Saper et al., 2002; Woods, 2009). In fed animals, NPY neurons are in a less active state with reduced firing activity (Takahashi and Cone, 2005) and NPY expression, but NPY release is increased during periods of food restriction or deprivation (Brady et al., 1995; Yoshihara et al., 1996). Indeed in our studies, fed NPY neurons were relatively quiescent and were more hyperpolarized compared to the fasted state. The depolarized state reported in fasted males may be due to pre-synaptic inputs, peripheral hormonal signals such as leptin or the modulation of endogenous cation conductances, e.g. the M-current.

Most arcuate NPY neurons express multiple KCNQ subunits and exhibit a robust outward K^+ current in fed males. One of the primary functions of the M-current is to control action potential firing, spike-frequency adaptation, depolarizing afterpotentials and afterhyperpolarization (AHP) currents (Gu et al., 2005; Peters et al., 2005; Tzingounis and Nicoll, 2008; Yue and Yaari, 2004). Therefore, we examined the effects of XE991, a selective blocker of KCNQ channels, on the excitability of NPY neurons from fed males. We found that by blocking the M-current, quiescent NPY neurons became depolarized and started firing continuously with a concomitant reduction in the rheobase. Although, action potential firing did not occur spontaneously in the presence of glutaminergic and GABAergic blockers, the membrane potential was similarly depolarized. Therefore, the M-current plays a role in action potential generation in arcuate NPY neurons as in other neuronal cell types (Hu et al., 2007; Leao et al., 2009).

Interestingly under our examination, NPY-GFP neurons in the fed state were more hyperpolarized in comparison to the initial description of this transgenic strain (Van den Pol et al., 2009). In the van den Pol study, NPY neurons exhibited a mean RMP of -56.6 mV and were highly active exhibiting regular, irregular and burst firing patterns. However, the neurons in the van den Pol study were mostly from juvenile mice (2–6 weeks of age) of unreported sex examined with subtly different external and internal solutions. Therefore, the differences in membrane properties and excitability between the two studies may be due to differences in experimental conditions and/or changes in feeding patterns between weanling-juvenile mice used in the latter study and the mature, adult mice used in the present experiments, as suggested in earlier studies (Leibowitz et al., 2005).

Fasting significantly reduces the current while increasing the neuronal excitability of the NPY neurons. This effect on M-current activity is partially restored by a 24 hr re-feeding but not by a shorter 2 hr re-feeding. The suppression of the M-current due to fasting appears to be an NPY-specific effect as a similar effect is not found in POMC neurons between fed and fasted male mice ($n=4, 5$ respectively, Roepke *et al.* unpublished findings). Moreover, the

suppression of the M-current in NPY neurons by fasting may involve peripheral signals such as serum leptin concentrations, which are higher during states of satiety, driving KCNQ channel expression. However, it is not known, at this time, if leptin affects KCNQ2/3 or M-current expression or that another hormone, peptide or neurotransmitter is responsible for altered expression in the different feeding states in these hypothalamic neurons.

The role of the M-current in the control of food intake by 17 β -estradiol

E2 is known to reduce NPY/AgRP expression in ovariectomized females (Bonavera et al., 1994; Crowley et al., 1985; Pelletier et al., 2007) and during proestrous when serum E2 levels are elevated (Olofsson et al., 2009). Short-term (3–48 hr) E2 replacement reduces NPY mRNA expression and protein immunoreactivity in the arcuate nucleus (Crowley et al., 1985; Pelletier et al., 2007), and long-term (18 days) E2 treatment reduces the *in vitro* release of NPY in the paraventricular nucleus (Bonavera et al., 1994). The suppression of NPY/AgRP mRNA expression in intact females is associated with a reduction in food intake and body weight, and E2 attenuates food intake during re-feeding (2 hr) after fasting, indicating that the interplay between fasting and E2 is complex (Olofsson et al., 2009). E2 replacement also attenuates the increase in food intake following NPY infusion into the lateral ventricle (Santollo and Eckel, 2008). These data suggests that the anorectic effects of E2 involve the suppression of NPY expression and neuronal function, which may involve the potentiation of the M-current.

NPY neurons are significantly hyperpolarized in the E2- versus oil-treated, ovariectomized females, in part, because of the potentiation of the M-current. This potentiation could be due, in part, to the KCNQ5 subunit expression. Furthermore, the KCNQ5 subunit contributes to the medium and slow AHP current (Tzingounis et al., 2010) in hippocampal CA3 neurons, which may also be altered in the E2-treated NPY neurons to reduce firing frequency. Ultimately, a reduction in NPY activity would lead to less NPY/AgRP release and subsequently an attenuation in feeding activity.

Fasting abrogated the effects of E2-treatment on M-current activity. This fasting effect was due, in part, to reduction of KCNQ2/3 expression in the native neurons resulting in a significant depolarization of the membrane potential, similar to the fasted males. The effects of fasting overwhelmed the effects of E2 (elevated KCNQ5 expression) at the neuronal level via the reduction of the other KCNQ subunits, especially KCNQ3. Co-expression of KCNQ3 with KCNQ5 in heterologous cell systems increases current amplitudes 2–3 fold over KCNQ5 homomultimers and also produces currents with similar amplitudes to the native M-current in hippocampal neurons and KCNQ2/3 heteromultimers in the heterologous cells (Lerche et al., 2000; Tzingounis et al., 2010). Thus, the more robust M-current measured in fed, E2-treated NPY neurons may be due to a greater association of KCNQ3/5 while the effects of fasting reduces this association via a reduction in KCNQ3.

We have previously shown that E2 will increase KCNQ5 expression in the guinea pig arcuate nucleus both after 24 hr and long-term treatment using unbiased microarray analysis followed by quantitative real-time PCR (Roepke et al., 2008; Roepke et al., 2007). In the mouse, E2 replacement also increased the expression of KCNQ5 both in the whole arcuate and specifically in NPY neurons. The gene regulation is lost in ER α KO transgenic mice suggesting that ER α is involved in KCNQ5 gene regulation. However, ER α mRNA and ER α protein are detected in only 19% and in <4%, respectively, of native NPY neurons and co-localizes with only approximately 20% of KCNQ5-expressing NPY neurons. Therefore, these data suggest that cellular localization of ER α in NPY neurons is below detectability or that ER α -mediated gene regulation is not the sole mechanism. Similarly, previous immunohistochemical studies have reported low levels or absence of co-localization between NPY neurons and ER α (Olofsson et al., 2009; Simonian et al., 1999). The estrogenic

regulation of KCNQ5 may be via ER-protein associations (Sp-1, Ap-1, etc.) because a putative estrogen response element (ERE) has not been identified within the KCNQ5 gene using an online promoter sequence search tool (Roepke et al., 2007). Consequently, the low level of ER α co-localization suggests that E2's effects on the M-current occurs via multiple mechanisms including nuclear receptor-mediated gene regulation, presynaptic inputs from other arcuate neurons and/or potentially via a membrane-mediated (G-protein coupled receptor) mechanism (Kelly and Rønnekleiv, 2008) that will be examined in future experiments.

In conclusion, NPY neurons are orexigenic and active during states of negative energy balance. Suppression of the M-current by gene regulation, and potentially other pathways, is a novel mechanism for the increase in NPY neuronal activity during fasting. Furthermore, E2 treatment augments the M-current, in part, via KCNQ5 gene regulation and is abrogated during fasting. The opposing actions of E2 and fasting on the M-current suggest that the M-current plays a pivotal role in NPY neuronal excitability and may be an important cellular target for neurotransmitter and hormonal signals to control energy homeostasis in both males and females.

Acknowledgments

This research was supported by United States Public Health Grants DK 68098, NS 43330 and NS 38809 and K99 DK 83457 for TAR. We thank Martha A. Bosch for her expert technical assistance with single-cell harvesting and quantitative real-time PCR.

References

- Bonavera JJ, Dube MG, Kalra PS, Kalra SP. Anorectic effects of estrogen may be mediated by decreased neuropeptide-Y release in the hypothalamic paraventricular nucleus. *Endocrinology*. 1994; 134:2367–2370. [PubMed: 8194462]
- Brady LS, Smith MA, Gold PW, Herkenham M. Altered expression of hypothalamic neuropeptide mRNA in food-restricted and food-deprived rats. *Neuroendocrinology*. 1995; 52:441–447. [PubMed: 2177853]
- Bronson FH, Vom Saal FS. Control of the preovulatory release of luteinizing hormone by steroids in the mouse. *Endocrinology*. 1979; 104:1247–1255. [PubMed: 571329]
- Crowley WR, Tessel RE, O'Donohue TL, Adler BA, Kalra SP. Effects of ovarian hormones on the concentrations of immunoreactive neuropeptide Y in discrete brain regions of the female rat: correlation with serum luteinizing hormone (LH) and median eminence LH-releasing hormone. *Endocrinology*. 1985; 117:1151–1155. [PubMed: 3893992]
- Delmas P, Brown DA. Pathways modulating neural KCNQ/M (Kv7) potassium channels. *Nat Rev Neurosci*. 2005; 6:850–862. [PubMed: 16261179]
- Elmedy P, Calloe K, Schmitt N, Schultz Hansen R, Grunnet M, Oleson S-P. Modulation of ERG channels by XE991. *Basic Clin Pharmacol Toxicol*. 2007; 100:316–322. [PubMed: 17448117]
- Elmqvist JK. Hypothalamic pathways underlying the endocrine, autonomic, and behavioral effects of leptin. *Physiol Behav*. 2001; 74:703–708. [PubMed: 11790432]
- Gao Q, Horvath TL. Neurobiology of feeding and energy expenditure. *Annu Rev Neurosci*. 2007; 30:367–398. [PubMed: 17506645]
- Giloh H, Sedat JW. Fluorescence microscopy: Reduced photobleaching of rhodamine and fluorescein protein conjugates by *n*-propyl gallate. *Science*. 1982; 217:1252–1255. [PubMed: 7112126]
- Gropp E, Shanabrough M, Borok E, Xu AW, Janoschek R, Buch T, Plum L, Balthasar N, Hampel B, Waisman A, Barsh GS, Horvath TL, Brüning JC. Agouti-related peptide-expressing neurons are mandatory for feeding. *Nat Neurosci*. 2005; 8:1289–1291. [PubMed: 16158063]
- Gu N, Vervaeke K, Hu H, Storm JF. Kv7/KCNQ/M and HCN/h, but not K_{ca}2/SK channels, contribute to the somatic medium after-hyperpolarization and excitability control in CA1 hippocampal pyramidal cells. *J Physiol*. 2005; 566.3:689–715. [PubMed: 15890705]

- Haskell-Luevano C, Cone RD, Monck EK, Wan Y-P. Structure activity studies of the Melanocortin-4 receptor by *in Vitro* mutagenesis: Identification of agouti-related protein (AGRP), melanocortin agonist and synthetic peptide antagonist interaction determinants. *Biochem.* 2001; 40:6164–6179. [PubMed: 11352754]
- Hu H, Vervaeke K, Storm JF. M-channels (Kv7/KCNQ channels) that regulate synaptic integration, excitability, and spike pattern of CA1 pyramidal cells are located in the perisomatic region. *J Neurosci.* 2007; 27:1853–1867. [PubMed: 17314282]
- Ibrahim N, Bosch MA, Smart JL, Qiu J, Rubinstein M, Rønnekleiv OK, Low MJ, Kelly MJ. Hypothalamic proopiomelanocortin neurons are glucose-responsive and express K-ATP channels. *Endocrinology.* 2003; 144:1331–1340. [PubMed: 12639916]
- Kelly MJ, Loose MD, Rønnekleiv OK. Opioids hyperpolarize β -endorphin neurons via μ -receptor activation of a potassium conductance. *Neuroendocrinology.* 1990; 52:268–275. [PubMed: 2170854]
- Kelly MJ, Rønnekleiv OK. Membrane-initiated estrogen signaling in hypothalamic neurons. *Mol Cell Endocrinol.* 2008; 290:14–23. [PubMed: 18538919]
- Kohno D, Nakata M, Maekawa F, Fujiwara K, Maejima Y, Kuramochi M, Shimazaki T, Okano H, Onaka T, Yada T. Leptin suppresses ghrelin-induced activation of neuropeptide Y neurons in the arcuate nucleus via phosphatidylinositol 3-kinase- and phosphodiesterase 3-mediated pathway. *Endocrinology.* 2007; 148:2251–2263. [PubMed: 17303662]
- Leao RN, Tan HM, Fisahn A. Kv7/KCNQ channels control action potential phasing of pyramidal neurons during hippocampal gamma oscillations in vitro. *J Neurosci.* 2009; 29:13353–13363. [PubMed: 19846723]
- Leibowitz SF, Sepiashvili K, Akabayashi A, Karatayev O, Davydova Z, Alexander JT, Wang J, Chang G-Q. Function of neuropeptide Y and agouti-related protein at weaning: relation to corticosterone, dietary carbohydrate and body weight. *Brain Res.* 2005; 1036:180–191. [PubMed: 15725416]
- Lerche C, Scherer CR, Seebohm G, Derst C, Wei AD, Busch AE, Steinmeyer K. Molecular cloning and functional expression of KCNQ5, a potassium channel subunit that may contribute to neuronal M-current diversity. *J Biol Chem.* 2000; 275:22395–22400. [PubMed: 10787416]
- Livak KJ, Schmittgen TD. Analysis of relative gene expression data using real-time quantitative PCR and the 2(-delta-delta C(T)) method. *Methods.* 2001; 25:402–408. [PubMed: 11846609]
- Loose MD, Rønnekleiv OK, Kelly MJ. Membrane properties and response to opioids of identified dopamine neurons in the guinea pig hypothalamus. *J Neurosci.* 1990; 10:3627–3634. [PubMed: 1977895]
- Lu X-Y, Nicholson JR, Akil H, Watson SJ. Time course of short-term and long-term orexigenic effects of agouti-related protein (86–132). *NeuroReport.* 2001; 12:1281–1284. [PubMed: 11338207]
- Luquet S, Perez FA, Hnasko TS, Palmiter RD. NPY/AgRP neurons are essential for feeding in adult mice but can be ablated in neonates. *Science.* 2005; 310:683–685. [PubMed: 16254186]
- Olofsson LE, Pierce AA, Xu AW. Functional requirement of AgRP and NPY neurons in ovarian cycle-dependent regulation of food intake. *Proc Natl Acad Sci USA.* 2009; 106:15932–15937. [PubMed: 19805233]
- Pelletier G, Li S, Luu-The V, Labrie F. Oestrogenic regulation of pro-opiomelanocortin, neuropeptide Y and corticotrophin-releasing hormone mRNAs in mouse hypothalamus. *J Neuroendocrinol.* 2007; 19:426–431. [PubMed: 17388940]
- Peters HC, Hu H, Pongs O, Storm JF, Isbrandt D. Conditional transgenic suppression of M channels in mouse brain reveals functions in neuronal excitability, resonance and behavior. *Nat Neurosci.* 2005; 8:51–60. [PubMed: 15608631]
- Pfaffl MW. A new mathematical model for relative quantification in real-time RT-PCR. *Nucleic Acid Res.* 2001; 29:2002–2007.
- Qiu J, Bosch MA, Tobias SC, Grandy DK, Scanlan TS, Rønnekleiv OK, Kelly MJ. Rapid signaling of estrogen in hypothalamic neurons involves a novel G protein-coupled estrogen receptor that activates protein kinase C. *J Neurosci.* 2003; 23:9529–9540. [PubMed: 14573532]
- Qiu J, Fang Y, Rønnekleiv OK, Kelly MJ. Leptin excites proopiomelanocortin neurons via activation of TRPC channels. *J Neurosci.* 2010; 30:1560–1565. [PubMed: 20107083]

- Robbins J. KCNQ potassium channels: physiology, pathophysiology, and pharmacology. *Pharmacol Ther.* 2001; 90:1–19. [PubMed: 11448722]
- Roepke TA. Oestrogen modulates hypothalamic control of energy homeostasis through multiple mechanisms. *J Neuroendocrinol.* 2009; 21:141–150. [PubMed: 19076267]
- Roepke TA, Malyala A, Bosch MA, Kelly MJ, Rønnekleiv OK. Estrogen regulation of genes important for K⁺ channel signaling in the arcuate nucleus. *Endocrinology.* 2007; 148:4937–4951. [PubMed: 17595223]
- Roepke TA, Xue C, Bosch MA, Scanlan TS, Kelly MJ, Rønnekleiv OK. Genes associated with membrane-initiated signaling of estrogen and energy homeostasis. *Endocrinology.* 2008; 149:6113–6124. [PubMed: 18755790]
- Santollo J, Eckel LA. Estradiol decreases the orexigenic effect of neuropeptide Y, but not agouti-related protein, in ovariectomized rats. *Behav Brain Res.* 2008; 191:173–177. [PubMed: 18453005]
- Saper CB, Chou TC, Elmquist JK. The need to feed: homeostatic and hedonic control of eating. *Neuron.* 2002; 36:199–211. [PubMed: 12383777]
- Schmittgen TD, Livak KJ. Analyzing real-time PCR data by the comparative C_T method. *Nature Protocols.* 2008; 3:1101–1108.
- Schroeder BC, Hechenberger M, Weinreich F, Kubisch C, Jentsch TJ. KCNQ5, a novel potassium channel broadly expressed in brain, mediates M-type currents. *J Biol Chem.* 2000; 275:24089–24095. [PubMed: 10816588]
- Schweitzer P. Cannabinoids decrease the K⁺ M-current in hippocampal CA1 neurons. *J Neurosci.* 2000; 20:51–58. [PubMed: 10627580]
- Shen W, Hamilton SE, Nathanson NM, Surmeier DJ. Cholinergic suppression of KCNQ channel currents enhances excitability of striatal medium spiny neurons. *J Neurosci.* 2005; 10:7449–7458. [PubMed: 16093396]
- Simonian SX, Spratt DP, Herbison AE. Identification and characterization of estrogen receptor α -containing neurons projecting to the vicinity of the gonadotropin-releasing hormone perikarya in the rostral preoptic area of the rat. *J Comp Neurol.* 1999; 411:346–358. [PubMed: 10404258]
- Stanley BG, Leibowitz SF. Neuropeptide Y injected in the paraventricular hypothalamus: a powerful stimulant of feeding behavior. *Proc Natl Acad Sci USA.* 1985; 82:3940–3943. [PubMed: 3858854]
- Takahashi KA, Cone RD. Fasting induces a large, leptin-dependent increase in the intrinsic action potential frequency of orexigenic arcuate nucleus neuropeptide Y/Agouti-related protein neurons. *Endocrinology.* 2005; 146:1043–1047. [PubMed: 15591135]
- Tkatch T, Baranaukas G, Surmeier DJ. Kv4.2 mRNA abundance and A-type K⁺ current amplitude are linearly related in basal ganglia and basal forebrain neurons. *J Neurosci.* 2000; 20:579–588. [PubMed: 10632587]
- Tzingounis AV, Heidenreich M, Kharkovets T, Spitzmaul G, Jensen HS, Nicoll RA, Jentsch TJ. The KCNQ5 potassium channel mediates a component of the afterhyperpolarization current in mouse hippocampus. *Proc Natl Acad Sci U S A.* 2010; 107:10232–10237. [PubMed: 20534576]
- Tzingounis AV, Nicoll RA. Contribution of KCNQ2 and KCNQ3 to the medium and slow afterhyperpolarization currents. *Proc Natl Acad Sci USA.* 2008; 105:19974–19979. [PubMed: 19060215]
- Van den Pol AN, Yao Y, Fu L-Y, Foo K, Huang H, Coppari R, Lowell BB, Broberger C. Neuromedin B and gastrin-releasing peptide excite arcuate nucleus neuropeptide Y neurons in a novel transgenic mouse expressing stong *renilla* green fluorescent protein in NPY neurons. *J Neurosci.* 2009; 29:4622–4639. [PubMed: 19357287]
- Wang HS, Pan ZM, Shi WM, Brown BS, Wymore RS, Cohen IS, Dixon JE, McKinnon D. KCNQ2 and KCNQ3 potassium channel subunits: Molecular correlates of the M-channel. *Science.* 1998; 282:1890–1893. [PubMed: 9836639]
- Wang J-H, Wang F, Yang M-J, Yu D-F, Wu W-N, Liu J, Ma L-Q, Cai F, Chen J-G. Leptin regulated calcium channels of NPY and POMC neurons by activation of different signal pathways. *Neuroscience.* 2008; 156:89–98. [PubMed: 18588949]
- Woods SC. The control of food intake: behavioral versus molecular perspectives. *Cell Metab.* 2009; 9:489–498. [PubMed: 19490904]

- Xu C, Roepke TA, Zhang C, Rønnekleiv OK, Kelly MJ. GnRH activates the M-current in GnRH neurons: an autoregulatory negative feedback mechanism? *Endocrinology*. 2008; 149:2459–2466. [PubMed: 18218690]
- Yang M-J, Wang F, Wang J-H, Wu W-N, Hu Z-L, Cheng J, Yu D-F, Long L-H, Fu H, Xie N, Chen J-G. PI3K integrates the effects of insulin and leptin on large-conductance Ca²⁺-activated K⁺ channels in neuropeptide Y neurons of the hypothalamic arcuate nucleus. *Am J Physiol Endocrinol Metab*. 2010; 298:E193–E201. [PubMed: 19671839]
- Yoshihara T, Honma S, Honma K-I. Effects of restricted daily feeding on neuropeptide Y release in the rat paraventricular nucleus. *Am J Physiol Endocrinol Metab*. 1996; 270:E589–E595.
- Yue C, Yaari Y. KCNQ/M channels control spike afterdepolarization and burst generation in hippocampal neurons. *J Neurosci*. 2004; 24:4614–4624. [PubMed: 15140933]
- Zhang C, Bosch MA, Levine JE, Rønnekleiv OK, Kelly MJ. Gonadotropin-releasing hormone neurons express K_{ATP} Channels that are regulated by estrogen and responsive to glucose and metabolic inhibition. *J Neurosci*. 2007; 27:10153–10164. [PubMed: 17881521]
- Zhang C, Bosch MA, Rick EA, Kelly MJ, Rønnekleiv OK. 17β-estradiol regulation of T-type calcium channels in gonadotropin-releasing hormone neurons. *J Neurosci*. 2009; 29:10552–10562. [PubMed: 19710308]
- Zhang C, Roepke TA, Kelly MJ, Rønnekleiv OK. Kisspeptin depolarizes GnRH neurons through activation of TRPC-like cationic channels. *J Neurosci*. 2008; 28:4423–4434. [PubMed: 18434521]




Figure 1. Blocking the M-current increases firing activity in NPY neurons from fed male mice
A, In current clamp, the rheobase was determined in both control and XE991 (40 μM) conditions. To elicit action potentials, the resting membrane potential (RMP) was held at -80 mV, followed by injection of small amounts of current (~ 6 pA/step, 500 ms). The black trace is the first current injection to evoke an AP in the control conditions and multiple spikes in the XE991 conditions. The gray trace is the first current injection which evokes a spike in XE991 conditions but not the control conditions. **B**, In fed males, the average rheobase in control conditions was significantly higher than after XE991 (40 μM , 10 min) treatment ($n=15$; **, $p<0.01$). **C**, The firing activity of NPY neurons in fed males was continuously recorded in control and XE991 (40 μM) conditions. A typical example of an

NPY neuron at rest (RMP = -75 mV) and the effects of XE991 to depolarize the cell (to -58.5 mV) and induce firing. **D**, The firing activity of NPY neurons in fed males was continuously recorded in control and XE991 (40 μ M) conditions but in the presence of glutamate receptor blockers (CNQX (10 μ M); AP5 (50 μ M)) and a GABA-A receptor blocker (bicuculline (20 μ M)). A typical example of an NPY neuron at rest (RMP = -74 mV) and the effects of XE991 to depolarize the cell (to -57.5 mV) without inducing firing. The average depolarization with and without blockers was 14.9 ± 3.2 mV ($n=3$, at 20–22 min) and 16.8 ± 1.3 mV ($n=15$, at 12–14 min), respectively. **E**, Action potentials were evoked by current injection before (control) and after XE991 perfusion from the neuron recorded in Figure 1D. The black trace is the first current injection to evoke a response in the control conditions and the gray trace is the first current injection during XE991 application.




Figure 2. Fasting reduces the activity of the M-current in NPY neurons

A, The slow deactivation of the M-current (I_M) in NPY neurons was recorded from fed, male mice using whole-cell patch clamp recording. From a holding potential of -60 mV, a voltage jump to -20 mV (300 ms) was followed by steps from -25 to -75 mV in 5 mV increments (500 ms). Currents were recorded under control conditions and after 10 min incubation with XE991 ($40 \mu\text{M}$), a selective KCNQ blocker in the presence of TTX ($1 \mu\text{M}$). The XE991-sensitive current was calculated by subtracting the current relaxation (the difference between the instantaneous and steady state (arrows)) in XE991-treated traces from the control traces. This XE991-sensitive current is graphed as an I-V plot (see Figure 2D) (Xu et al., 2008). **B**, The K^+ currents evoked by the deactivation protocol do not rundown over the 30 min recording period. T=0 is 3 minutes after whole-cell access. XE991 ($40 \mu\text{M}$) was perfused after 30 min to demonstrate the amount of M-current still active. The figure is from one cell, but rundown was examined in 2 other cells with the same results. **C**, To determine the most effective dose to measure the M-current across treatments, a sequential concentration response of XE991 (1, 10, $40 \mu\text{M}$) was applied to same neuron (n=5). I-V plots were analyzed by a two-way ANOVA ($p < 0.05$, $F = 4.67$, $df = 3$) followed by *Bonferroni-Dunn* comparison. a, $p < 0.05$; b, $p < 0.01$; c, $p < 0.001$ compared to control. **D**, In NPY neurons, the I-V plot of **A** from fed males (n=11); fasted males (n=17) and 24 hr re-fed males (n=10). I-V plots were analyzed by a two-way ANOVA ($p < 0.01$, $F = 7.3$, $df = 2$) followed by *Bonferroni-Dunn* comparison. a, $p < 0.05$; c, $p < 0.001$ compared to fasted.




Figure 3. Fasting reduces the activity of the M-current but not XE991-insensitive K^+ currents in NPY neurons

A, The activation of the M-current (I_M) in NPY neurons was recorded from fed, fasted or re-fed, male mice using the following protocol: In the presence of TTX ($1 \mu M$) and from a holding potential of -60 mV, the voltage was jumped to -90 mV (500 ms) followed by steps from -80 to $+10$ mV in 10 mV increments (1 s) and returned to -60 mV. Currents were recorded in control conditions and after 10 min incubation with XE991 ($40 \mu M$). The XE991-sensitive current was calculated by subtracting the steady state (SS) current (the last 50 ms of the voltage steps) in the XE991-treated traces from the control traces. This difference is graphed in the I-V plots as the amplitude of the XE991-sensitive current (Leão et al., 2009). **B**, The I-V curve of **A** is shown for fed males ($n=11$); fasted males ($n=15$) and 24 hr re-fed males ($n=10$). The currents at $+10$ mV for the three feeding states were 484 ± 37.6 pA for fed males; 259 ± 63.3 pA for fasted males; and 480 ± 53.6 pA for re-fed males. I-V plots were analyzed by a two-way ANOVA (ns) followed by *Bonferroni-Dunn* comparison. **a**, $p < 0.05$; **b**, $p < 0.01$; **c**, $p < 0.001$ compared to fasted. **C**, A plot of the conductances at 0 mV between each feeding state comparing total, XE991-sensitive and -insensitive currents. **D**, Summary of the ratio ((Inst-SS)/SS) of the time-dependent (Inst-SS) and steady state (SS) currents from control and XE991-treated conditions in response to a $+10$ mV step. **, $p < 0.01$.




Figure 4. 17β -estradiol increases the activity of the M-current in NPY neurons

A, Using the same protocol described in Figure 2, the I–V curve in the E2-treated females (n=15), the oil-treated females (n=14) and 24 hr fasted females (n=11). I–V plots were analyzed by a two-way ANOVA ($p < 0.01$, $F = 6.0$, $df = 2$) followed by *Bonferroni-Dunn* comparison. **a**, $p < 0.05$; **b**, $p < 0.01$; **c**, $p < 0.001$ compared to oil-treated and fasted E2-treated.




Figure 5. E2 increases M-current activity but not XE991-insensitive K⁺ currents in NPY neurons
A, The activation of the I_M in NPY neurons was recorded from ovariectomized, oil-treated (n=14); E2-treated (n=12) and fasted, E2-treated (n=11) female mice using the protocol described in Figure 3. The current at +10 mV for the three treatments was 231 ± 24 pA; 513 ± 82 pA; and 250 ± 31 pA, respectively. I–V plots were analyzed by a two-way ANOVA (p<0.001, F=11.0, df=2) followed by *Bonferroni-Dunn* comparison. **a**, p<0.05; **b**, p<0.01; **c**, p<0.001 compared to E2-treated. **B**, A plot of the conductances at 0 mV between each steroid treatment comparing control, XE991-sensitive and -insensitive currents. **C**, Summary of the ratio ((Inst-SS)/SS) of the time-dependent (Inst-SS) and steady state (SS) currents

from control and XE991-treated conditions in response to a +10 mV step. *, $p < 0.05$; **, $p < 0.01$.




Figure 6. KCNQ subunits are highly expressed in NPY neurons
A, a representative gel and **B**, a frequency distribution plot illustrating the average number of NPY neurons expressing KCNQ2, KCNQ3 and KCNQ5 subunits from ovariectomized females. The number in the column equals the number of females. The total number of cells analyzed was 110 with an average of 15 cells per female. KCNQ4 is not expressed in NPY neurons. The ladder for the KCNQ2 & 3 gels has been overexposed in order to visualize the lower molecular (100 bp) weight bands.




Figure 7. Fasting reduces KCNQ2 & KCNQ3 expression and 17 β -estradiol increases KCNQ 5 expression in NPY neurons

A–C, Quantitative real-time PCR assays for KCNQ2 (A), KCNQ3 (B), and KCNQ5 (C) transcripts using the SybrGreen method. Standard curves were prepared with BH cDNA serial dilutions as follows: 1:50, 1:100, 1:500, 1:1000, 1:5000, 1:7500, 1:10000. Cycle number was plotted against the normalized fluorescence intensity (Delta Rn) to visualize the PCR amplification. The cycle threshold (C_T , arrow) is the point in the amplification at which the sample values were calculated. The BH cDNA serial dilutions and one representative NPY pool (black squares) are illustrated in A, B, C. Insets in A, B, and C show standard curve regression lines, slopes from serial dilution data and the dissociation

(melting) curves for all three transcripts. The amplification efficiencies calculated from the slopes were 96% for KCNQ2, 100% for KCNQ3, and 94% for KCNQ5. The melting curves depict single-product melting at 83 °C, 78 °C, and 80 °C for KCNQ2,-3,-5, respectively. Single peaks illustrate that only one product was formed in NPY pools. β -Actin was used as control and its primer pair was also 100% efficient (data not shown). Negative (CSF, -RT cell and tissue samples) and positive tissue controls were analyzed on each plate along with the experimental samples. These efficiencies allowed us to use the $\Delta\Delta$ CT method for quantification. **D**, Using cell harvesting coupled with quantitative real-time PCR, fasting significantly reduced the expression of KCNQ2 and KCNQ3 in NPY neurons (3–4 pools of 5 (KCNQ2/3) or 10 (KCNQ5) cells each from 5 animals, n=5) from male mice. **, p<0.01. **E**, E2 treatment significantly increased the expression of KCNQ5 while fasting significantly reduced the expression of KCNQ2 and KCNQ3 in NPY neurons (3–4 pools of 5 (KCNQ2/3) or 10 (KCNQ5) cells each from 5 animals, n=5) from E2-treated female mice. *, p<0.05; **, p<0.01 compared to oil-treated. KCNQ2 and KCNQ3 expression in fed, E2-treated and fasted, E2-treated were significantly different (p<0.05).




Figure 8. 17 β -estradiol regulation of KCNQ5 expression is ER α -dependent and a small population of NPY neurons express ER α

A, E2 treatment in ovariectomized, wild-type females increased the mRNA expression of KCNQ5 in the arcuate nucleus but failed to regulate KCNQ5 expression in ER α knock-out mice. *, $p < 0.05$. The number in the column equals the number of animals per treatment. **B**, A representative gel illustrating the expression of ER α in NPY neurons harvested from ovariectomized females and the co-expression of KCNQ5. **C**, Immunocytochemistry showed a small population of GFP-NPY neurons co-localizing ER α : NPY-GFP neurons to left and ER α -immunoreactive neurons in middle with an overlay illustrating co-localization (indicated by arrows) to right. 99% of GFP-labeled neurons express NPY mRNA. 3V demarks the third ventricle. White bar in the overlay represents 25 μ m.

Table 1

The membrane potential and input resistance for each feeding state and steroid treatment

Treatment	Control		XE991	
	Membrane potential (mV)	Input resistance (G Ω)	Membrane potential (mV)	Input resistance (G Ω)
Males				
Fed	-71.5 \pm 1.6 **	0.64 \pm 0.12	-62.5 \pm 2.9 ^a	1.02 \pm 0.18 ^a
Fasted	-64.8 \pm 1.6	0.65 \pm 0.12	-57.1 \pm 1.9 ^b	0.81 \pm 0.09
Re-fed	-68.5 \pm 0.9	0.63 \pm 0.07	-58.3 \pm 1.5 ^d	0.97 \pm 0.11 ^b
Females				
Oil	-68.1 \pm 1.8 *	0.65 \pm 0.09	-62.7 \pm 1.3 ^a	1.19 \pm 0.28
E2	-72.4 \pm 1.1	0.68 \pm 0.08	-62.4 \pm 2.0 ^c	1.0 \pm 0.12 ^b
Fasted E2	-65.6 \pm 1.5 ***	0.63 \pm 0.06	-58.8 \pm 2.0 ^a	0.79 \pm 0.12

^a =p<0.05;

^b =<0.01;

^c =p<0.001;

^d =p<0.0001 denote control to XE991 comparisons.

* =p<0.05;

** =<0.01;

=p<0.001; denote fed vs. fasted comparisons in males and oil & fasted E2 compared to E2 in females. For males, n=11 for fed; n=17 for fasted; and n=10 for re-fed. For females, n=14 for oil; n=15 for fed E2; and n=11 for fasted E2. The cells from the rheobase measurement experiments (Figure 1) are not included in the Table since they were recorded under different conditions.