

Published in final edited form as:

*Antiviral Res.* 2012 June ; 94(3): 258–271. doi:10.1016/j.antiviral.2012.03.005.

## Quercetin inhibits rhinovirus replication *in vitro* and *in vivo*

Shyamala Ganesan<sup>1</sup>, Andrea N. Faris<sup>1</sup>, Adam T. Comstock<sup>1</sup>, Qiong Wang<sup>1</sup>, Suparna Nanua<sup>1</sup>, Marc B. Hershenon<sup>1,2</sup>, and Uma S. Sajjan<sup>1</sup>

<sup>1</sup>Department of Pediatrics and Communicable Diseases, University of Michigan, Ann Arbor, MI, USA 48109

<sup>2</sup>Department of Internal Medicine, <sup>2</sup>Department of Molecular and Integrative Physiology, University of Michigan, Ann Arbor, MI, USA 48109

### Summary

Rhinovirus (RV), which is responsible for the majority of common colds, also causes exacerbations in patients with asthma and chronic obstructive pulmonary disease. So far, there are no drugs available for treatment of rhinovirus infection. We examined the effect of quercetin, a plant flavanol on RV infection *in vitro* and *in vivo*. Pretreatment of airway epithelial cells with quercetin decreased Akt phosphorylation, viral endocytosis and IL-8 responses. Addition of quercetin 6 h after RV infection (after viral endocytosis) reduced viral load, IL-8 and IFN responses in airway epithelial cells. This was associated with decreased levels of negative and positive strand viral RNA, and RV capsid protein, abrogation of RV-induced eIF4GI cleavage and increased phosphorylation of eIF2 $\alpha$ . In mice infected with RV, quercetin treatment decreased viral replication as well as expression of chemokines and cytokines. Quercetin treatment also attenuated RV-induced airway cholinergic hyperresponsiveness. Together, our results suggest that quercetin inhibits RV endocytosis and replication in airway epithelial cells at multiple stages of the RV life cycle. Quercetin also decreases expression of pro-inflammatory cytokines and improves lung function in RV-infected mice. Based on these observations, further studies examining the potential benefits of quercetin in the prevention and treatment of RV infection are warranted.

### Keywords

infection; translation; transcription; inflammation

### Introduction

Rhinovirus (RV) is a single-stranded RNA virus from the *Picornaviridae* family which is responsible for the majority of upper respiratory tract infections (Makela et al., 1998; Winther, 2011). Although RV infections cause only mild disease and are self-limiting in normal individuals, they represent a significant economic burden, particularly in loss of working hours and missed school days (Proud, 2005). Accumulating evidence also suggests that RV is an important trigger of acute exacerbations in patients with chronic obstructive pulmonary disease (COPD) and asthma (Johnston et al., 1995; Seemungal et al., 2001;

© 2012 Elsevier B.V. All rights reserved.

\*Address correspondence to: Uma Sajjan, Ph.D., University of Michigan, 1150 W. Medical Center Dr., Room 3570, MSRBII, Box 5688, Ann Arbor, MI USA 48109-0688. Phone: (734) 936-4200; FAX: (734) 764-3200; usajjan@umich.edu.

**Publisher's Disclaimer:** This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Seemungal et al., 2000). Experimental RV infection of asthma and COPD patients results in the development of lower respiratory disease, airflow obstruction, and systemic and airway inflammation which are greater and more prolonged than the similarly-infected control subjects (Mallia et al., 2006, 2011; Message et al., 2008). The clinical outcomes of experimental RV infection correlate with viral load and inflammatory cytokines in sputum and bronchoalveolar lavage.

Current therapies for treating virus-related acute exacerbations of asthma and COPD are only partially effective and may also have side effects. Similarly, while currently available therapies may relieve symptoms, no FDA-approved drugs exist for the treatment of RV infections. Therefore, an antiviral drug which possesses anti-inflammatory activity and at the same time effectively inhibits viral replication may be beneficial in treating RV infections.

The RV life cycle offers many opportunities for intervention. The initial stage of the viral life cycle involves binding to a cell surface receptor, endocytosis, acidification of the endosome, viral uncoating, and release of viral RNA (vRNA) to the cytoplasm (Prchla et al., 1994). In the next stage, vRNA is translated as a single large polyprotein from an internal ribosomal entry site (IRES) which is located in the 5' untranslated region of RV genome. The polyprotein is processed through sequential enzymatic cleavages by two virus-encoded proteinases, 2A and 3C proteinase (Foeger et al., 2003; Hellen et al., 1989). Simultaneously, transcription of viral genome that is catalyzed by virus-encoded RNA polymerase begins (Meredith et al., 1999). In the final stage of viral replication, RNA and capsid proteins are assembled into mature viral particles, which are then released by cell lysis. During the replication process, RNA viruses, including RV, modify the cellular functions to their advantage. For example, RV proteinase 2A attacks the host translational machinery in susceptible HeLa cells by cleaving eukaryotic initiation factor (eIF)-4GI and eIF4GII, thereby shutting off cap-dependent host cell protein synthesis (Glaser and Skern, 2000; Gradi et al., 2003). Drugs which interfere with any of the above steps would be beneficial in treating RV infection.

Viruses also interfere with host defense mechanisms. Upon viral infection, host cells negatively regulate global protein translation in order to limit viral replication. Protein kinase R (PKR), which is activated by viral double-stranded (ds) RNA, phosphorylates eIF2 $\alpha$  phosphorylation, thereby inhibiting translation of both host and vRNA (Jacobs and Langland, 1996; Qian et al., 2004). Non-structural proteins of respiratory syncytial virus, influenza virus and hepatitis virus competitively inhibit eIF2 $\alpha$  binding to PKR, thereby inhibiting eIF2 $\alpha$  phosphorylation (Gale et al., 1997; Groskreutz et al., 2010; Li et al., 2006; Tan and Katze, 2001). Conversely, reversal of such inhibition would augment viral clearance.

Quercetin (3,3',4',5,7-pentahydroxyflavone) is a plant flavonoid present in many plants including broccoli, apples, berries, onions, and tea. Flavonoids share a common chemical structure consisting of two phenol rings linked through three carbons. Quercetin has potent antioxidant effects, combined with free radical species to form considerably less reactive phenoxy radicals. Quercetin also has inhibitory effects on several lipid, protein tyrosine and serine/threonine kinases including phosphatidylinositol (PI) 3-kinase, and induces nitric oxide synthase (Agullo et al., 1997; Davies et al., 2000; Huang et al., 1999; Jiang et al., 2006; Peet and Li, 1999). Quercetin has also been shown to exert antiviral activities *in vitro*. Quercetin or its derivatives reduce the infectivity or replication of herpes simplex virus type 1, poliovirus type 1, parainfluenza virus type 3, adenovirus, respiratory syncytial virus and influenza virus *in vitro* (Castrillo and Carrasco, 1987; Chiang et al., 2003; Kaul et al., 1985; Kim et al., 2010b; Neznanov et al., 2008). Among the several quercetin derivatives examined, 4',7-dihydroxy-3-methoxy-5,6-dimethylflavone exhibited potent antirhinoviral effects in

human skin fibroblasts (De Meyer et al., 1991). However, quercetin by itself had no antirhinoviral effect in this study. Later, quercetin was shown to inhibit the SARS-CoV protease that is required for SARS virus replication (Chen et al., 2006) as well as hepatitis C viral protein translation and release of newly formed viral particles (Gonzalez et al., 2009). Quercetin supplementation was also shown to reduce susceptibility to influenza A virus infection and severity of disease following stressful exercise in mice (Davis et al., 2008) and symptoms of upper respiratory tract infections in athletes (Nieman et al., 2007). This is of importance, because quercetin supplementation increases skeletal muscle mitochondrial biogenesis modestly in untrained athletes and this is associated with improved exercise performance (Nieman et al., 2010). Since mitochondria orchestrate antiviral signaling following recognition of cytosolic viral components (Tal and Iwasaki, 2011), there may exist a relationship between quercetin-induced mitochondrial biogenesis and reduced susceptibility to influenza infection. Although, association of physical stress/fatigue with susceptibility to RV infection is not established, RV causes severe exacerbations in patients with COPD and asthma (Mallia et al., 2006, 2011; Message et al., 2008). Understanding the mechanisms by which quercetin inhibits rhinovirus infectivity and replication is necessary to determine the suitability of quercetin to treat RV infections in these patients. In the present study, we examined the effects of quercetin treatment on RV endocytosis, viral genome transcription and translation, and pro-inflammatory responses *in vitro*. We also assessed the effect of quercetin *in vivo* using a novel mouse model of RV infection (Newcomb et al., 2008).

## Methods and Materials

### Rhinovirus

RV1B and RV39 were purchased from American Type Culture Collection (Manassas, VA) and viral stocks were generated as described previously (Newcomb et al., 2005). Fifty percent cell culture infectivity (CCID<sub>50</sub>) values of viral stocks were determined by the Spearman-Kärber method (Johnston and Tyrrell, 1997). To generate replication-deficient virus (UV-RV), the RV preparation was exposed to ultraviolet light for 15 min on ice at 100 m joules/cm<sup>2</sup> (Chattoraj et al., 2011).

### Cell culture and infection

BEAS-2B immortalized human bronchial epithelial cells (American Type Culture Collection) were cultured in bronchial epithelial cell growth medium (BEGM, Lonza, Walkersville, MD) in collagen-coated plates, as described previously (Chattoraj et al., 2011; Wang et al., 2009).

For RV infection, cells were grown in 24- or 6-well plates until 90% confluency. Cells were infected with RV at a multiplicity of infection (MOI) of 1.0 based on CCID<sub>50</sub> and incubated for 90 min at 33°C to allow binding and endocytosis of RV. Infection medium was replaced with fresh medium and incubated for 4.5 h at 33°C. Quercetin dihydrate (Sigma, St. Louis, MO) or DMSO (vehicle control) was added to the required concentration and further incubated for 16 to 18 h at 33°C. In some experiments, cells were pretreated with quercetin for 1 h, infected with RV for 1 h in the presence of quercetin, and the infection medium was replaced with fresh medium containing DMSO or quercetin. In selected experiments, cells were pretreated with quercetin or DMSO overnight, and shifted to quercetin medium for 8h. Cells were rinsed with fresh medium and then infected with RV1B in the absence of quercetin.

## Endocytosis of RV

Endocytosis of RV by airway epithelial cells was measured as described previously (Chattoraj et al., 2011; Newcomb et al., 2005). Briefly, BEAS-2B cells were pretreated with quercetin or DMSO and then incubated with RV or equal volumes of sham (cell lysate from uninfected HeLa cells) at 4°C for 30 min. Cells were washed and then incubated at 37°C for 30 min to facilitate endocytosis of bound RV. Cells were permeabilized and endocytosed RV was detected by antibody to RV capsid protein, VP2 (Mosser et al., 2002) and Alexafluor 488-conjugated secondary antibody and quantified by flow cytometry, correcting for sham controls. To visualize endocytosed RV, cells were infected, incubated for 30 min, fixed in cold methanol, blocked with 5% normal donkey serum and treated with antibodies to RV capsid protein and early endosomal antigen (EEA)-1 (Santa Cruz Biotechnologies, Santa Cruz, CA). Bound antibodies were detected by Alexafluor 488- (RV) or Alexafluor 594- (EEA-1) conjugated (RV) secondary antibody. Cell nuclei were counterstained with DAPI and the cells were visualized by confocal microscopy.

## Animals and treatment

Eight to ten week-old C57BL/6 mice were infected with sham or RV1B through the intranasal route and gavaged with 0.2 mg of quercetin daily for one, or four days. The control group consisted of similarly infected mice treated with vehicle (50% propylene glycol, PG). In some experiments mice were treated with quercetin or vehicle as above for 10 days. Two days after the last treatment, mice were infected with RV or sham. These experiments were approved by the Animal Care and Use Committee of the University of Michigan.

## Viral titer

After appropriate treatment, cells along with medium were subjected to one freeze-thaw cycle and centrifuged. Mice were sacrificed at 1, 4 and 7 days post-infection. Lungs were collected under aseptic conditions and homogenized in 1 ml of PBS. Lung homogenates were subjected to two freeze thaw cycles and centrifuged. The viral titer (CCID<sub>50</sub>) in supernatants of cell cultures or lung homogenates was measured as described previously (Sajjan et al., 2009).

## Cytokines

Protein levels of cytokines in cell culture or lung homogenate supernatants were measured by multiplex immunoassay (Biorad, Hercules, CA) or ELISA (R&D Systems, Minneapolis, MN).

## Measurement of airways responsiveness

Mice were anesthetized by intraperitoneal injection of ketamine (2.5–5 mg/100 g body weight) and pentobarbital (2mg/100 g body weight). A steel cannula was inserted into the trachea and connected to a miniature computerized ventilator (Buxco FinePointe Operating system, Wilmington, NC). Airways responsiveness to methacholine was measured by nebulizing increasing doses of methacholine, as described previously (Sajjan et al., 2009).

## Realtime PCR

Total RNA was isolated from the lungs or cell cultures by using RNeasy (Qiagen, Valencia, CA). cDNA was synthesized utilizing a first strand synthesis kit (Applied Biosystems, Carlsbad, CA). mRNA levels of IFN- $\alpha$ , IFN- $\beta$ , IFN- $\lambda_1$  and IFN- $\lambda_{2/3}$  were quantified by real-time PCR using gene specific primers and expressed as fold increase over the housekeeping gene glyceraldehyde-3-phosphate dehydrogenase (G3PDH). Positive strand RV RNA was quantified by qPCR or as described previously (Newcomb et al., 2008; Sajjan

et al., 2009). To detect negative strand RV RNA, 5 µg of total RNA was converted to cDNA using a primer specific to negative strand RV RNA and superscript III reverse transcriptase (Invitrogen, Carlsbad, CA) as described previously (Newcomb et al., 2008). cDNA was amplified by qPCR using nested primers and the products were analyzed by agarose gel electrophoresis.

### Western blotting

After relevant treatment, cells were lysed in RIPA buffer containing complete protease inhibitors (Roche Diagnostics, Indianapolis, IN) and phosphatase inhibitors (1mM sodium fluoride and 1mM sodium orthovanadate). Cell lysates were centrifuged and total protein in the supernatant was determined. Equal amounts of protein were subjected to SDS-PAGE and proteins transferred to nitrocellulose membranes. The membranes were blocked with either 5% bovine serum albumin or fat-free milk and probed with antibody to phospho-Akt, total Akt, and phospho-eIF2α (all from Cell Signaling Technology, Danvers, MA), eIF4GI (Abcam, Cambridge, MA), or β-actin (Sigma-Aldrich, St. Louis, MO). The bound antibody was detected with appropriate second antibody conjugated with horseradish peroxidase (Biorad) and chemiluminescent substrate (Pierce, Rockford, IL).

### Statistical analysis

Statistical analysis of significance was calculated by either one-way analysis of variance followed by Tukey's post-hoc test or an unpaired t test with Welch's correction using Sigma Stat software (Systat Software, San Jose, CA). Results are shown as mean ± SD.

## Results

### Pretreatment of airway epithelial cells with quercetin decreases RV-induced IL-8 responses

BEAS-2B cells were pretreated with 5, 10 or 25 µM quercetin or an identical volume of DMSO and infected with sham, RV1B, RV39, UV-irradiated (UV)-RV1B or UV-RV39. IL-8 protein levels were determined by ELISA. A concentration of 10 µM quercetin attenuated both RV and UV-RV-stimulated IL-8 responses by 75 to 80% (Figures 1A and 1B). Treatment of cells with 25 µM quercetin, did not decrease IL-8 further, and therefore in subsequent experiments we used 10 µM quercetin.

### Quercetin pretreatment inhibits RV endocytosis by airway epithelial cells

Previously, we demonstrated in cultured airway epithelial cells that RV binding and endocytosis is sufficient for the initial IL-8 response to infection (Newcomb et al., 2005). We also showed that endocytosis of RV requires activation of PI 3-kinase. Since quercetin inhibits PI-3 kinase activity (Davies et al., 2000), we determined whether quercetin pretreatment inhibits endocytosis of RV. Quercetin pretreatment significantly decreased endocytosis of both RV1B and RV39 by BEAS-2B cells (Figures 2A and 2B). We observed similar attenuation in the endocytosis of UV-RV. We also examined the colocalization of RV1B and EEA1, an early endosomal marker, by confocal microscopy. Cells inoculated with sham HeLa cell lysate showed diffuse signals for EEA1 in the cytoplasm (data not shown). Cells infected with RV1B showed clustering of RV particles. These cells also showed punctate staining for EEA1 as opposed to diffuse staining, an indication of recruitment of EEA1 to endosomes. Clustered RV particles frequently co-localized with endosomal EEA1, indicating endocytosis of RV in these cells. (Figure 2C). In contrast, cells pretreated with quercetin and infected with RV1B showed an EEA1 distribution similar to sham-treated cells and only faint signals for RV which did not co-localize with EEA1-positive endosomes. (The faint signals for RV may represent the bound, but not endocytosed

virus.) These results suggest that pretreatment with quercetin may block endocytosis of RV. Next we examined whether quercetin inhibits RV-induced phosphorylation of Akt, a downstream effector of PI 3-kinase. As observed previously, RV1B treated cells showed increased Akt phosphorylation compared to sham-infected cells (Figure 2D). Quercetin pretreatment completely inhibited RV-induced Akt phosphorylation. These results indicate that quercetin decreases RV endocytosis and PI 3-kinase/Akt signaling, leading to a reduction in IL-8 expression.

### **Endocytosis-independent effects of quercetin on RV1B-stimulated IFN and IL-8 responses in airway epithelial cells**

Next, to determine endocytosis-independent effects, BEAS-2B airway epithelial cell monolayers were infected with RV1B or RV39 and incubated for 6 h to allow time for viral endocytosis. Quercetin or DMSO was then added to the cell culture medium and incubated for another 18 h. Cells inoculated with sham served as negative controls. Cellular mRNA levels of IFN- $\beta$ , IFN- $\lambda$ 1 and IFN- $\lambda$ 2/3 and cell supernatant protein levels of IL-8 were determined. As observed earlier (Newcomb et al., 2005; Wang et al., 2009), infection with RV1B and RV39 each stimulated significantly higher IFN- $\beta$ , IFN- $\lambda$ 1 and IFN- $\lambda$ 2 than UV-irradiated virus (Figures 3A–3C). We also observed decrease in protein levels of IFN. In contrast, UV-RV1B or UV-RV39-infected cells showed significant increases in IL-8 similar to intact RV (Figure 3D), suggesting that IFNs are stimulated only by intact, replicating RV. Quercetin treatment inhibited RV-stimulated IFN as well as IL-8 expression, consistent with the notion that quercetin blocks viral replication.

### **Quercetin-treated cells show decreased viral load**

To determine whether decreased IFN and IL-8 responses were due to decreased viral load, we determined viral titer. Cells were infected with RV1B, RV39, UV-RV1B or UV-RV39 and DMSO or quercetin (10  $\mu$ M) added 6 h later. Cells were incubated for another 18 h. Viral titer was determined at 6 h post-infection (before the addition of quercetin) and 24 h after infection (18 h after adding quercetin). As expected, cells infected with UV-RV1B or UV-RV39 did not show the presence of infectious RV (not shown). Compared to 6 h after infection, cells infected with intact RV1B or RV39 showed an increased load of infectious RV at 24 h post infection, indicating replication of virus in BEAS2B cells (Figure 4A). Twenty-four h post-infection, cells treated with quercetin showed a viral load 3 logs lower than similarly infected, vehicle-treated cells. We also measured the viral load by assessing vRNA levels. As expected, RV-infected cells showed significantly increased vRNA copy numbers compared to UV-RV-infected cells (Figure 4B). In contrast, RV-infected cells treated with quercetin showed vRNA copy numbers similar to those infected with UV-RV. These results are again consistent with the notion that quercetin may inhibit viral replication.

### **Quercetin interferes with viral genome transcription**

During viral genome amplification, negative-strand vRNA is generated as an intermediate. Therefore, we monitored the negative-strand RNA in RV-infected BEAS2B cells by PCR. We observed amplification of negative-strand RNA in cells infected with RV1B or RV39, but not in cells infected with replication-deficient UV-RV (Figure 4C). In RV-infected, quercetin-treated cells, negative-strand RNA was either decreased or not detected, implying that quercetin may inhibit viral replication at the transcriptional level.

### **Quercetin attenuates RV-induced cleavage of eIF4GI**

During viral replication, viral proteases cleave eIF4GI to inhibit formation of the initiation complex required for host cap-dependent protein synthesis. This in turn facilitates translation of viral proteins. Cells were infected with RV1B, RV39, UV-RV1B or UV-RV39

as above and quercetin or DMSO was added 6 h post RV infection. After overnight incubation, cells infected with intact RV1B or RV39 showed cleavage of eIF4GI, but not cells infected with sham or UV-RV. Treatment with quercetin completely abolished RV1B- and RV39-induced eIF4GI cleavage (Figure 5A).

Increased eIF2 $\alpha$  phosphorylation limits mRNA translation globally and is essential to inhibit viral replication in infected cells. Therefore, we assessed eIF2 $\alpha$  phosphorylation in airway epithelial cells infected with RV1B. Cells infected with intact RV1B showed increased eIF2 $\alpha$  phosphorylation compared to cells inoculated with sham or replication-deficient UV-RV (Figure 5B). As we showed previously (Nanua et al., 2006), quercetin treatment further increased eIF2 $\alpha$  phosphorylation. Together these results imply that quercetin may decrease viral genome translation by inhibiting cleavage of eIF4GI and increasing eIF2 $\alpha$  phosphorylation.

Next, we determined whether quercetin decreased viral protein levels, examining the levels of the RV capsid protein VP2 by Western blot analysis. While sham-infected cells did not show VP2 capsid protein, RV-infected cells showed two bands corresponding to VP2 and VP2 along with VP4 (VP2+VP4), a precursor of VP2 (Figure 5C). Quercetin treated cells showed lower amounts of VP2 and VP2+VP4 bands, consistent with the notion that quercetin-induced changes in translation initiation may reduce the translation of viral RNA.

### Presence of quercetin during RV infection is necessary to limit viral replication

Next, we examined whether withdrawal of quercetin 8 h prior to viral infection limits viral replication and associated IL-8 and IFN responses. Cells were pretreated with quercetin overnight and shifted to quercetin free medium. Eight hours later, cells were rinsed with medium and infected with RV1B in the absence of quercetin and vRNA, viral titer, IL-8 and IFN production were determined. Although, cells pretreated with quercetin showed a slightly lower viral titer and vRNA than the vehicle-treated cells, the difference was not statistically significant (Figure 6A and 6B). Similarly, no difference in IFNs or IL-8 levels between quercetin and vehicle treated cells was observed (Figure 6C and 6D). This is not surprising because quercetin or its metabolite levels in the cells peak at 5 h post treatment and it is slowly reduced with time and completely depleted by 24h (Kim et al., 2009, 2010a). Therefore addition of quercetin during or immediately after viral infection may be necessary to effectively limit viral infection *in vitro*.

### Quercetin treatment decreases viral load *in vivo*

RV39, which binds to intercellular adhesion molecule (ICAM)-1, does not infect mouse cells. On the other hand, RV1B, which binds to low density lipoprotein receptor (LDLR) or LDLR-like receptors, infects mouse airway epithelial cells *in vitro* and *in vivo*. Mice infected with RV1B show the presence of negative-strand viral RNA in the lungs, transmissibility of RV infection from the lung homogenates of inoculated mice to cultured HeLa cells, and the induction of a robust lung interferon response (Newcomb et al., 2008; Sajjan et al., 2009; Wang et al., 2011). Therefore, we used RV1B to determine the effects of quercetin *in vivo*. Normal mice were infected with RV1B or sham by intranasal route. After two hours, mice were treated with quercetin or propylene glycol (vehicle). Mice were sacrificed 1 or 4 days later and assessed for lung viral load. As anticipated, mice infected with sham did not show infectious virus (data not shown). Mice infected with RV1B and treated with vehicle showed significant levels ( $9 \times 10^4$  CCID<sub>50</sub>/ml) of infectious virus 1 day post infection (Figure 7A). At 4 days after infection, there was no detectable infectious virus, as observed previously (Newcomb et al., 2008). On the other hand, RV1B-infected mice treated with quercetin showed less than 1 log infectious virus/lung, *i.e.*, 4 logs less virus compared to vehicle-treated mice. Similarly, quercetin-treated mice also showed

significantly lower vRNA levels at both 1 and 4 days post infection compared to vehicle-treated mice (Figure 7B).

To determine whether the observed decrease in viral load also depends on the inhibition of viral replication, we determined the level of negative-strand RNA in mouse lungs. RV1B-infected animals treated with vehicle showed detectable levels of negative strand vRNA 1 day after infection (Figure 7C) which was decreased in similarly infected mice treated with quercetin. At 4 days post-infection, neither vehicle- nor quercetin-treated mice showed detectable levels of negative-strand vRNA (data not shown). These results imply that, in this model, RV replication occurs only within the first 24 hours after infection, and that quercetin effectively inhibits this process.

### **Quercetin decreases RV-stimulated chemokines and cytokines *in vivo***

Previously, we have shown that RV infection stimulates lung chemokine and cytokine expression (Newcomb et al., 2008). To determine whether quercetin decreases RV-stimulated chemokine and cytokines levels, we determined the protein levels of CXCL-1 (KC) and CXCL-2 (MIP-2), TNF- $\alpha$  and CCL2 (MCP-1) by single or multiplex ELISA. Mice infected with RV and treated with vehicle showed significantly increased levels of all four cytokines/chemokines compared to mice infected with sham at 1 day but not 4 days post-infection, correlating with the presence of virus in the lungs (Figure 8A–D). Quercetin treatment decreased levels of all four cytokines in RV-infected mice, implying that quercetin attenuates RV-stimulated pro-inflammatory responses in mice by decreasing viral load.

Next, we measured the mRNA levels of representative type I (IFN- $\beta$ ) and III IFN (IFN- $\lambda_1$ ), the expression of which depends on generation of the double-stranded RNA intermediate during RV replication. As expected, mice infected with RV1B and treated with vehicle showed significantly higher levels of IFN- $\alpha$  and IFN- $\lambda_1$  than the sham-infected animals at 1 day post infection, which returned to normal levels by 4 days after RV infection (Figure 8E and 8F). This correlated with the presence of negative- strand vRNA in the lungs. RV1B-infected mice treated with quercetin showed significantly decreased levels of both IFNs compared to RV-infected, vehicle-treated mice, consistent with the decreased negative strand vRNA in quercetin-treated mice.

### **Quercetin decreases RV-induced airways cholinergic hyperresponsiveness**

In our previous studies, we demonstrated that RV infection increase airways responsiveness to methacholine challenge at 1 day post-infection (Newcomb et al., 2008), which returns to normal by 4 days after RV infection. We therefore determined the effects of quercetin on airways responsiveness to methacholine one day after RV infection. As expected, compared to sham-inoculated mice, RV-infected mice treated with vehicle showed increased airways responsiveness, particularly at methacholine doses of 10 and 20 mg/ml (Figure 9). In contrast, RV-infected mice treated with quercetin showed a level of airways responsiveness similar to sham-infected, quercetin-treated animals, which was significantly lower than that observed in RV-infected vehicle treated mice. Together, these results suggest that *in vivo* quercetin treatment decreases viral load and RV-stimulated chemokine and cytokine responses while improving lung function.

### **Quercetin treatment during RV infection is necessary for efficient viral clearance and reduce lung inflammation and improve lung function**

To determine whether quercetin pretreatment modulates innate immune responses to subsequent viral infections and thus aid viral clearance, mice were pretreated with quercetin for 10 days, followed by a washout period of 2 days. Mice were then infected with RV1B as above, sacrificed 24 h later and levels of proinflammatory cytokines, IFN expression,



vRNA, viral titers and lung function were determined. As observed above, mice infected with RV1B showed increased levels of pro-inflammatory cytokines and IFN responses and this was not affected by quercetin pretreatment (Figure 10). In addition, quercetin pretreated mice showed similar levels of vRNA, and viral titer to those mice treated with vehicle (Figure 11A and 11B). Further, quercetin pretreated mice infected with RV1B also showed airways hyperresponsiveness to methacholine challenge similar to mice pretreated with vehicle (Figure 11D). These results indicate that quercetin treatment during or after viral infection rather than pretreatment is more effective *in vivo*. These findings are not surprising, because half-life of quercetin ranges between 11 to 28h (Boots et al., 2008) and therefore one would expect the plasma quercetin levels return to basal levels by 48 h.

## Discussion

In the present study, we demonstrate that quercetin decreases RV infection at multiple steps in the viral life cycle. Quercetin decreases viral endocytosis and inhibits RV-induced IL-8 responses in cultured airway epithelial cells. When quercetin was added after viral endocytosis, it also decreased RV-stimulated IL-8 and IFN responses, consistent with downstream effects. Quercetin inhibited replication of RV by suppressing transcription of RV genome. Further, quercetin decreased RV-induced cleavage of eIF4GI and increased phosphorylation of eIF2 $\alpha$ . Finally, we showed that quercetin treatment of RV-infected mice decreased viral replication, expression of pro-inflammatory cytokines and chemokines, and airways hyperresponsiveness to methacholine challenge. Collectively, these results suggest that quercetin may be beneficial in limiting viral replication and reducing symptoms associated with RV infection.

Quercetin is a potent antioxidant and possesses anti-inflammatory properties. It inhibits several lipid, protein tyrosine and serine/threonine kinases by competing for the ATP binding site (reviewed in (Williams et al., 2004)). In fact, a potent inhibitor of PI-3 kinase, LY 294002 was designed based on the structure of quercetin (Vlahos et al., 1994). We and others have shown that PI-3 kinase activity is required for RV endocytosis and also for viral replication-independent IL-8 responses (Bentley et al., 2007; Lau et al., 2008; Newcomb et al., 2005). Therefore, treatment of cells with quercetin prior to or during RV infection could inhibit viral infection by abolishing RV endocytosis, an essential initial step of RV infection. Consistent with this idea, pretreatment of cells with quercetin decreased Akt phosphorylation, endocytosis of RV and attenuated the associated chemokine responses in human airway epithelial cells.

In addition to inhibiting viral entry, quercetin may also inhibit subsequent steps in the picornavirus life cycle. For instance, quercetin and its derivative 3-methylquercetin were shown to inhibit synthesis of poliovirus RNA (Castrillo and Carrasco, 1987; Neznanov et al., 2008) and this was attributed to inhibition of viral RNA polymerase 3D<sup>pol</sup> (Castrillo and Carrasco, 1987). Viral RNA polymerase 3D<sup>pol</sup> is required for conversion of viral genome to create negative-strand RNA, which serves as a template for generating positive strand viral genome (Castrillo and Carrasco, 1987; Hung et al., 2002). Recently, polyphenols including quercetin were shown to increase zinc uptake in Caco2 cells and zinc ions have been shown to inhibit 3D<sup>pol</sup> *in vitro* (Hung et al., 2002; Krenn et al., 2005; Sreenivasulu et al., 2010). Further, quercetin may also interfere with the processing of RV polyprotein by RV proteases that is required for activation of RNA polymerase (Hellen et al., 1989). In addition to processing viral 3D<sup>pol</sup>, viral protease 2A also cleaves eIF4G to inhibit cap-dependent host protein synthesis and promote viral genome translation (Glaser and Skern, 2000; Gradi et al., 2003). Our results demonstrate that quercetin significantly reduced both negative and positive strand viral RNA and this was associated with reduced cleavage of eIF4GII and reduction in viral capsid protein, VP2, suggesting that quercetin may inhibit initial

polypeptide processing that is required for both viral RNA polymerase processing and for cleavage of eIFG4II thereby blocking all the downstream steps in RV replication. Another possibility is that quercetin may directly inhibit viral RNA polymerase thereby blocking genome translation and synthesis of new progeny virus.

One of the mechanisms by which host limits viral replication is through shutting down the global protein synthesis by increasing phosphorylation of eIF2 $\alpha$  (Jacobs and Langland, 1996), (Qian et al., 2004). We observed that quercetin increases phosphorylation of eIF2 $\alpha$ , beyond that stimulated by RV infection itself and therefore it is likely that quercetin also limits viral replication by enhancing host innate immune responses. Recently, replication of enterovirus, a member of picornaviridae family was shown to occur in specialized organelles enriched in phosphoinositide-4-phosphate (PI4P) lipids generated by PI-4-kinase III $\beta$  (Hsu et al., 2010). If RV replication requires PI4P-enriched organelles, it is possible that quercetin which inhibits PI-4 kinases (Prajda et al., 1995) may interfere with formation of these organelles thus blocking viral replication. Together, these observations suggest that quercetin may interfere with viral replication by more than one mechanism.

In our previous studies, we demonstrated that RV1B infects normal mice, transiently replicates, increases the levels of pro-inflammatory cytokines and IFNs in the lung, and increases airways responsiveness to methacholine challenge (Newcomb et al., 2008; Sajjan et al., 2009; Wang et al., 2011). Using this model system, we show for the first time that quercetin not only inhibits RV replication *in vitro*, but also inhibits RV replication, RV-induced pro-inflammatory cytokine and chemokine expression, and RV-induced airway hyperresponsiveness *in vivo*. Mice treated with quercetin showed a significantly decreased viral load, diminished levels of negative-strand RNA, and decreased replication-dependent expression of type I and type III IFNs (Wang et al., 2009, 2011), providing direct evidence that quercetin effectively inhibits viral replication *in vivo*. In addition, quercetin treatment decreased RV-induced lung inflammation, as evidenced by reduced expression of CXCL-1, CXCL-2, CCL2 and TNF- $\alpha$ . We have previously shown that, in addition to their role in lung inflammation, CXCR2 ligands and TNF- $\alpha$  contribute to RV-induced airways hyperresponsiveness (Nagarkar et al., 2009). Neutrophils, attracted to the airways by CXCR2 ligands, induce a state of hyperresponsiveness by elaboration of TNF- $\alpha$  (Nagarkar et al., 2009). Therefore, decreased airways responsiveness observed in RV-infected mice treated with quercetin may be due to decreased lung CXCL1, CXCL2 and TNF- $\alpha$  levels.

Quercetin dihydrate is an aglycon and is passively absorbed by small intestine. Only 24% of the ingested quercetin aglycon is absorbed as oppose to 52% of naturally occurring quercetin glycosides (Hollman et al., 1995). Plasma quercetin levels are in the low nanomolar range (<100 nM) in humans consuming normal diet which is estimated to provide 5 to 40 mg/day quercetin (Boots et al., 2007; Harwood et al., 2007). Upon supplementation with 1g of quercetin aglycone/day increases plasma levels to 1.5  $\mu$ M (Boots et al., 2007; Conquer et al., 1998). Previously we have shown that treatment of mice with 0.2 mg of quercetin yield total plasma quercetin (including both quercetin and its primary conjugates) levels on the order of 150 nM (Ganesan et al., 2010; Nanua et al., 2006). This dose of quercetin was sufficient to reduce lung inflammation in a mouse model of chronic obstructive pulmonary disease, and also an allergic mouse model of asthma (Ganesan et al., 2010; Nanua et al., 2006; Rogerio et al., 2007). Based on the present study, it appears that a nanomolar range of plasma quercetin is also sufficient to inhibit viral replication *in vivo*, even though a higher concentration (10  $\mu$ M) of quercetin is required *in vitro*. Although, the precise explanation for this is unclear, it may depend on the differences in the stability and bioavailability of quercetin. One possibility is that quercetin may be absorbed better *in vivo* via small intestine compared to respiratory epithelial cells *in vitro*. Another possibility is that quercetin metabolites generated *in vivo* may be more potent as an antiviral agent than the parent

compound. Quercetin dihydrate is metabolized in different organs including small intestine and liver similar to quercetin glycosides (Harwood et al., 2007). Quercetin glycosides are in fact precursors of quercetin aglycon *in vivo*. Quercetin glycosides are initially hydrolyzed to form quercetin aglycon in the epithelial cells of small intestine (Murota and Terao, 2003) and the aglycon generated is metabolized by phase II enzymes present in the small intestine, liver and other organs to methylated, sulfated and glycosylated forms. One hour after intragastric administration of quercetin to rats, the majority of the absorbed quercetin was found in the form of quercetin glucuronides, sulfoglucuronides and sulfates as well as isorhamnetic conjugates (Justino et al., 2004). Quercetin glucuronide was proposed to be a more potent modulator of reactive oxygen-generating enzymes than quercetin (Terao et al., 2011). Further studies are required to assess the antiviral effects of known quercetin metabolites.

## Conclusion

We showed that quercetin, a potent antioxidant and anti-inflammatory agent, also possesses anti-rhinoviral effects. Quercetin inhibits viral infection at multiple stages, including endocytosis, transcription of the viral genome and viral protein synthesis. Further studies are needed to determine whether quercetin may be beneficial in preventing or treating rhinovirus infections, or reducing symptoms related to viral infection.

## Acknowledgments

This work was supported by NIH grants AT4793 and HL897720 (to U.S.) and HL81420 (to M.B.H.). S.G. conducted the study, analyzed the data and prepared the first draft of the manuscript; A.N.F., A.T.C., Q.W. and N.A. provided assistance with the experiments; M.B.H. performed critical review of the manuscript; M.B.H. and U.S. obtained the funding and conceived the study, and U.S. supervised and finalized the manuscript.

## References

- Agullo G, Gamet-Payraastre L, Manenti S, Viala C, Remesy C, Chap H, Payraastre B. Relationship between flavonoid structure and inhibition of phosphatidylinositol 3-kinase: a comparison with tyrosine kinase and protein kinase C inhibition. *Biochem Pharmacol.* 1997; 53:1649–1657. [PubMed: 9264317]
- Bentley JK, Newcomb DC, Goldsmith AM, Jia Y, Sajjan US, Hershenson MB. Rhinovirus activates interleukin-8 expression via a Src/p110beta phosphatidylinositol 3-kinase/Akt pathway in human airway epithelial cells. *J Virol.* 2007; 81:1186–1194. [PubMed: 17121804]
- Boots AW, Haenen GR, Bast A. Health effects of quercetin: from antioxidant to nutraceutical. *Eur J Pharmacol.* 2008; 585:325–337. [PubMed: 18417116]
- Boots AW, Li H, Schins RP, Duffin R, Heemskerk JW, Bast A, Haenen GR. The quercetin paradox. *Toxicol Appl Pharmacol.* 2007; 222:89–96. [PubMed: 17537471]
- Castrillo JL, Carrasco L. Action of 3-methylquercetin on poliovirus RNA replication. *J Virol.* 1987; 61:3319–3321. [PubMed: 2442414]
- Chattoraj SS, Ganesan S, Faris A, Comstock A, Lee WM, Sajjan US. *Pseudomonas aeruginosa* suppresses interferon response to rhinovirus infection in Cystic fibrosis, but not in normal bronchial epithelial cells. *Infect Immun.* 2011; 79:4131–4145. [PubMed: 21825067]
- Chen L, Li J, Luo C, Liu H, Xu W, Chen G, Liew OW, Zhu W, Puah CM, Shen X, Jiang H. Binding interaction of quercetin-3-beta-galactoside and its synthetic derivatives with SARS-CoV 3CL(pro): structure-activity relationship studies reveal salient pharmacophore features. *Bioorg Med Chem.* 2006; 14:8295–8306. [PubMed: 17046271]
- Chiang LC, Chiang W, Liu MC, Lin CC. In vitro antiviral activities of *Caesalpinia pulcherrima* and its related flavonoids. *J Antimicrob Chemother.* 2003; 52:194–198. [PubMed: 12837746]

- Conquer JA, Maiani G, Azzini E, Raguzzini A, Holub BJ. Supplementation with quercetin markedly increases plasma quercetin concentration without effect on selected risk factors for heart disease in healthy subjects. *J Nutr.* 1998; 128:593–597. [PubMed: 9482769]
- Davies SP, Reddy H, Caivano M, Cohen P. Specificity and mechanism of action of some commonly used protein kinase inhibitors. *Biochem J.* 2000; 351:95–105. [PubMed: 10998351]
- Davis JM, Murphy EA, McClellan JL, Carmichael MD, Gangemi JD. Quercetin reduces susceptibility to influenza infection following stressful exercise. *Am J Physiol Regul Integr Comp Physiol.* 2008; 295:R505–509. [PubMed: 18579649]
- De Meyer N, Haemers A, Mishra L, Pandey HK, Pieters LA, Vanden Berghe DA, Vlietinck AJ. 4'-Hydroxy-3-methoxyflavones with potent antipicornavirus activity. *J Med Chem.* 1991; 34:736–746. [PubMed: 1847431]
- Foeger N, Schmid EM, Skern T. Human rhinovirus 2 A pro recognition of eukaryotic initiation factor 4GI. Involvement of an exosite. *J Biol Chem.* 2003; 278:33200–33207. [PubMed: 12791690]
- Gale MJ Jr, Korth MJ, Tang NM, Tan SL, Hopkins DA, Dever TE, Polyak SJ, Gretch DR, Katze MG. Evidence that hepatitis C virus resistance to interferon is mediated through repression of the PKR protein kinase by the nonstructural 5A protein. *Virology.* 1997; 230:217–227. [PubMed: 9143277]
- Ganesan S, Faris AN, Comstock AT, Chatteraj SS, Chatteraj A, Burgess JR, Curtis JL, Martinez FJ, Zick S, Hershenson MB, Sajjan US. Quercetin prevents progression of disease in elastase/LPS-exposed mice by negatively regulating MMP expression. *Respir Res.* 2010; 11:131. [PubMed: 20920189]
- Glaser W, Skern T. Extremely efficient cleavage of eIF4G by picornaviral proteinases L and 2A in vitro. *FEBS Lett.* 2000; 480:151–155. [PubMed: 11034318]
- Gonzalez O, Fontanes V, Raychaudhuri S, Loo R, Loo J, Arumugaswami V, Sun R, Dasgupta A, French SW. The heat shock protein inhibitor Quercetin attenuates hepatitis C virus production. *Hepatology.* 2009; 50:1756–1764. [PubMed: 19839005]
- Gradi A, Svitkin YV, Sommergruber W, Imataka H, Morino S, Skern T, Sonenberg N. Human rhinovirus 2A proteinase cleavage sites in eukaryotic initiation factors (eIF) 4GI and eIF4GII are different. *J Virol.* 2003; 77:5026–5029. [PubMed: 12663812]
- Groskreutz DJ, Babor EC, Monick MM, Varga SM, Hunninghake GW. Respiratory syncytial virus limits alpha subunit of eukaryotic translation initiation factor 2 (eIF2alpha) phosphorylation to maintain translation and viral replication. *J Biol Chem.* 2010; 285:24023–24031. [PubMed: 20519500]
- Harwood M, Danielewska-Nikiel B, Borzelleca JF, Flamm GW, Williams GM, Lines TC. A critical review of the data related to the safety of quercetin and lack of evidence of in vivo toxicity, including lack of genotoxic/carcinogenic properties. *Food Chem Toxicol.* 2007; 45:2179–2205. [PubMed: 17698276]
- Hellen CU, Krausslich HG, Wimmer E. Proteolytic processing of polyproteins in the replication of RNA viruses. *Biochemistry.* 1989; 28:9881–9890. [PubMed: 2695162]
- Hollman PC, de Vries JH, van Leeuwen SD, Mengelers MJ, Katan MB. Absorption of dietary quercetin glycosides and quercetin in healthy ileostomy volunteers. *Am J Clin Nutr.* 1995; 62:1276–1282. [PubMed: 7491892]
- Hsu NY, Ilnytska O, Belov G, Santiana M, Chen YH, Takvorian PM, Pau C, van der Schaar H, Kaushik-Basu N, Balla T, Cameron CE, Ehrenfeld E, van Kuppeveld FJ, Altan-Bonnet N. Viral reorganization of the secretory pathway generates distinct organelles for RNA replication. *Cell.* 2010; 141:799–811. [PubMed: 20510927]
- Huang YT, Hwang JJ, Lee PP, Ke FC, Huang JH, Huang CJ, Kandaswami C, Middleton E Jr, Lee MT. Effects of luteolin and quercetin, inhibitors of tyrosine kinase, on cell growth and metastasis-associated properties in A431 cells overexpressing epidermal growth factor receptor. *Br J Pharmacol.* 1999; 128:999–1010. [PubMed: 10556937]
- Hung M, Gibbs CS, Tsiang M. Biochemical characterization of rhinovirus RNA-dependent RNA polymerase. *Antiviral Res.* 2002; 56:99–114. [PubMed: 12367717]
- Jacobs BL, Langland JO. When two strands are better than one: the mediators and modulators of the cellular responses to double-stranded RNA. *Virology.* 1996; 219:339–349. [PubMed: 8638399]


- Jiang JS, Shih CM, Wang SH, Chen TT, Lin CN, Ko WC. Mechanisms of suppression of nitric oxide production by 3-O-methylquercetin in RAW 264.7 cells. *J Ethnopharmacol.* 2006; 103:281–287. [PubMed: 16213685]
- Johnston SL, Pattemore PK, Sanderson G, Smith S, Lampe F, Josephs L, Symington P, O'Toole S, Myint SH, Tyrrell DA, et al. Community study of role of viral infections in exacerbations of asthma in 9–11 year old children. *Brit Med J.* 1995; 310:1225–1229. [PubMed: 7767192]
- Johnston, SL.; Tyrrell, DAJ. Rhinoviruses. In: Lennette, EH.; Schmidt, NJ., editors. *Diagnostic Procedures for Viral, Rickettsial, and Chlamydial Infections.* American Public Health Association; Washington D.C: 1997. p. 533-563.
- Justino GC, Santos MR, Canario S, Borges C, Florencio MH, Mira L. Plasma quercetin metabolites: structure-antioxidant activity relationships. *Arch Biochem Biophys.* 2004; 432:109–121. [PubMed: 15519302]
- Kaul TN, Middleton E Jr, Ogra PL. Antiviral effect of flavonoids on human viruses. *J Med Virol.* 1985; 15:71–79. [PubMed: 2981979]
- Kim MK, Park KS, Yeo WS, Choo H, Chong Y. In vitro solubility, stability and permeability of novel quercetin-amino acid conjugates. *Bioorg Med Chem.* 2009; 17:1164–1171. [PubMed: 19128975]
- Kim MK, Park KS, Lee C, Park HR, Choo H, Chong Y. Enhanced stability and intracellular accumulation of quercetin by protection of the chemically or metabolically susceptible hydroxyl groups with a pivaloxymethyl (POM) promoiety. *J Med Chem.* 2010a; 53:8597–8607. [PubMed: 21090565]
- Kim Y, Narayanan S, Chang KO. Inhibition of influenza virus replication by plant-derived isoquercetin. *Antiviral Res.* 2010b; 88:227–235. [PubMed: 20826184]
- Krenn BM, Holzer B, Gaudernak E, Triendl A, van Kuppeveld FJ, Seipelt J. Inhibition of polyprotein processing and RNA replication of human rhinovirus by pyrrolidine dithiocarbamate involves metal ions. *J Virol.* 2005; 79:13892–13899. [PubMed: 16254325]
- Lau C, Wang X, Song L, North M, Wiehler S, Proud D, Chow CW. Syk associates with clathrin and mediates phosphatidylinositol 3-kinase activation during human rhinovirus internalization. *J Immunol.* 2008; 180:870–880. [PubMed: 18178826]
- Li S, Min JY, Krug RM, Sen GC. Binding of the influenza A virus NS1 protein to PKR mediates the inhibition of its activation by either PACT or double-stranded RNA. *Virology.* 2006; 349:13–21. [PubMed: 16466763]
- Makela MJ, Puhakka T, Ruuskanen O, Leinonen M, Saikku P, Kimpimaki M, Blomqvist S, Hyypia T, Arstila P. Viruses and bacteria in the etiology of the common cold. *J Clin Microbiol.* 1998; 36:539–542. [PubMed: 9466772]
- Mallia P, Message SD, Gielen V, Contoli M, Gray K, Keadze T, Aniscenko J, Laza-Stanca V, Edwards MR, Slater L, Papi A, Stanciu LA, Kon OM, Johnson M, Johnston SL. Experimental rhinovirus infection as a human model of chronic obstructive pulmonary disease exacerbation. *Am J Respir Crit Care Med.* 2011; 183:734–742. [PubMed: 20889904]
- Mallia P, Message SD, Keadze T, Parker HL, Kon OM, Johnston SL. An experimental model of rhinovirus induced chronic obstructive pulmonary disease exacerbations: a pilot study. *Respir Res.* 2006; 7:116. [PubMed: 16956406]
- Meredith JM, Rohll JB, Almond JW, Evans DJ. Similar interactions of the poliovirus and rhinovirus 3D polymerases with the 3' untranslated region of rhinovirus 14. *J Virol.* 1999; 73:9952–9958. [PubMed: 10559308]
- Message SD, Laza-Stanca V, Mallia P, Parker HL, Zhu J, Keadze T, Contoli M, Sanderson G, Kon OM, Papi A, Jeffery PK, Stanciu LA, Johnston SL. Rhinovirus-induced lower respiratory illness is increased in asthma and related to virus load and Th1/2 cytokine and IL-10 production. *Proc Natl Acad Sci USA.* 2008; 105:13562–13567. [PubMed: 18768794]
- Mosser AG, Brockman-Schneider R, Amineva S, Burchell L, Sedgwick JB, Busse WW, Gern JE. Similar frequency of rhinovirus-infectible cells in upper and lower airway epithelium. *J Infect Dis.* 2002; 185:734–743. [PubMed: 11920291]
- Murota K, Terao J. Antioxidative flavonoid quercetin: implication of its intestinal absorption and metabolism. *Arch Biochem Biophys.* 2003; 417:12–17. [PubMed: 12921774]

- Nagarkar DR, Wang Q, Shim J, Zhao Y, Tsai WC, Lukacs NW, Sajjan U, Hershenson MB. CXCR2 is required for neutrophilic airway inflammation and hyperresponsiveness in a mouse model of human rhinovirus infection. *J Immunol.* 2009; 183:6698–6707. [PubMed: 19864593]
- Nanua S, Zick SM, Andrade JE, Sajjan US, Burgess JR, Lukacs NW, Hershenson MB. Quercetin blocks airway epithelial cell chemokine expression. *Am J Respir Cell Mol Biol.* 2006; 35:602–610. [PubMed: 16794257]
- Newcomb DC, Sajjan U, Nanua S, Jia Y, Goldsmith AM, Bentley JK, Hershenson MB. Phosphatidylinositol 3-kinase is required for rhinovirus-induced airway epithelial cell interleukin-8 expression. *J Biol Chem.* 2005; 280:36952–36961. [PubMed: 16120607]
- Newcomb DC, Sajjan US, Nagarkar DR, Wang Q, Nanua S, Zhou Y, McHenry CL, Hennrick KT, Tsai WC, Bentley JK, Lukacs NW, Johnston SL, Hershenson MB. Human rhinovirus 1B exposure induces phosphatidylinositol 3-kinase-dependent airway inflammation in mice. *Am J Respir Crit Care Med.* 2008; 177:1111–1121. [PubMed: 18276942]
- Neznanov N, Kondratova A, Chumakov KM, Neznanova L, Kondratov R, Banerjee AK, Gudkov AV. Quercetinase pirin makes poliovirus replication resistant to flavonoid quercetin. *DNA Cell Biol.* 2008; 27:191–198. [PubMed: 18067462]
- Nieman DC, Henson DA, Gross SJ, Jenkins DP, Davis JM, Murphy EA, Carmichael MD, Dumke CL, Utter AC, McAnulty SR, McAnulty LS, Mayer EP. Quercetin reduces illness but not immune perturbations after intensive exercise. *Med Sci Sports Exerc.* 2007; 39:1561–1569. [PubMed: 17805089]
- Nieman DC, Williams AS, Shanely RA, Jin F, McAnulty SR, Triplett NT, Austin MD, Henson DA. Quercetin's influence on exercise performance and muscle mitochondrial biogenesis. *Med Sci Sports Exerc.* 2010; 42:338–345. [PubMed: 19927026]
- Peet GW, Li J. IkappaB kinases alpha and beta show a random sequential kinetic mechanism and are inhibited by staurosporine and quercetin. *J Biol Chem.* 1999; 274:32655–32661. [PubMed: 10551820]
- Prajda N, Singhal RL, Yeh YA, Olah E, Look KY, Weber G. Linkage of reduction in 1-phosphatidylinositol 4-kinase activity and inositol 1,4,5-trisphosphate concentration in human ovarian carcinoma cells treated with quercetin. *Life Sci.* 1995; 56:1587–1593. [PubMed: 7723587]
- Prchla E, Kuechler E, Blaas D, Fuchs R. Uncoating of human rhinovirus serotype 2 from late endosomes. *J Virol.* 1994; 68:3713–3723. [PubMed: 8189509]
- Proud CG. eIF2 and the control of cell physiology. *Semin Cell Dev Biol.* 2005; 16:3–12. [PubMed: 15659334]
- Qian X, Karpova T, Sheppard AM, McNally J, Lowy DR. E-cadherin-mediated adhesion inhibits ligand-dependent activation of diverse receptor tyrosine kinases. *EMBO J.* 2004; 23:1739–1748. [PubMed: 15057284]
- Rogério AP, Kanashiro A, Fontanari C, da Silva EV, Lucisano-Valim YM, Soares EG, Faccioli LH. Anti-inflammatory activity of quercetin and isoquercitrin in experimental murine allergic asthma. *Inflamm Res.* 2007; 56:402–408. [PubMed: 18026696]
- Sajjan U, Ganesan S, Comstock AT, Shim J, Wang Q, Nagarkar DR, Zhao Y, Goldsmith AM, Sonstein J, Linn MJ, Curtis JL, Hershenson MB. Elastase- and LPS-exposed mice display altered responses to rhinovirus infection. *Am J Physiol.* 2009; 297:L931–944.
- Seemungal T, Harper-Owen R, Bhowmik A, Moric I, Sanderson G, Message S, Maccallum P, Meade TW, Jeffries DJ, Johnston SL, Wedzicha JA. Respiratory viruses, symptoms, and inflammatory markers in acute exacerbations and stable chronic obstructive pulmonary disease. *Am J Respir Crit Care Med.* 2001; 164:1618–1623. [PubMed: 11719299]
- Seemungal TA, Harper-Owen R, Bhowmik A, Jeffries DJ, Wedzicha JA. Detection of rhinovirus in induced sputum at exacerbation of chronic obstructive pulmonary disease. *Eur Respir J.* 2000; 16:677–683. [PubMed: 11106212]
- Sreenivasulu K, Raghu P, Nair KM. Polyphenol-rich beverages enhance zinc uptake and metallothionein expression in Caco-2 cells. *J Food Sci.* 2010; 75:H123–128. [PubMed: 20546406]
- Tal MC, Iwasaki A. Mitoxosome: a mitochondrial platform for cross-talk between cellular stress and antiviral signaling. *Immunol Rev.* 2011; 243:215–234. [PubMed: 21884179]

- Tan SL, Katze MG. How hepatitis C virus counteracts the interferon response: the jury is still out on NS5A. *Virology*. 2001; 284:1–12. [PubMed: 11352662]
- Terao J, Murota K, Kawai Y. Conjugated quercetin glucuronides as bioactive metabolites and precursors of aglycone in vivo. *Food Funct*. 2011; 2:11–17. [PubMed: 21773581]
- Vlahos CJ, Matter WF, Hui KY, Brown RF. A specific inhibitor of phosphatidylinositol 3-kinase, 2-(4-morpholinyl)-8-phenyl-4H-1-benzopyran-4-one (LY294002). *J Biol Chem*. 1994; 269:5241–5248. [PubMed: 8106507]
- Wang Q, Miller DJ, Bowman ER, Nagarkar DR, Schneider D, Zhao Y, Linn MJ, Goldsmith AM, Bentley JK, Sajjan US, Hershenson MB. MDA5 and TLR3 Initiate Pro-Inflammatory Signaling Pathways Leading to Rhinovirus-Induced Airways Inflammation and Hyperresponsiveness. *PLoS Pathog*. 2011; 7:e1002070. [PubMed: 21637773]
- Wang Q, Nagarkar DR, Bowman ER, Schneider D, Gosangi B, Lei J, Zhao Y, McHenry CL, Burgens RV, Miller DJ, Sajjan U, Hershenson MB. Role of double-stranded RNA pattern recognition receptors in rhinovirus-induced airway epithelial cell responses. *J Immunol*. 2009; 183:6989–6997. [PubMed: 19890046]
- Williams RJ, Spencer JP, Rice-Evans C. Flavonoids: antioxidants or signalling molecules? *Free Radic Biol Med*. 2004; 36:838–849. [PubMed: 15019969]
- Winther B. Rhinovirus infections in the upper airway. *Proc Am Thor Soc*. 2011; 8:79–89.


- Rhinovirus is responsible for majority of common colds
- There are no FDA approved drugs are available to treat rhinovirus infection.
- We show that quercetin, a plant flavonoid blocks viral replication in vitro.
- In addition quercetin inhibits virus-stimulated cytokine expression.
- Quercetin also inhibits viral replication and decreasing lung inflammation in vivo.






**Figure 1.**

Cells pretreated with quercetin show attenuated RV-stimulated IL-8 responses. BEAS-2B cells were pretreated with DMSO or quercetin for 1 h. Cells were infected with sham, UV-RV1B, RV1B, UV-RV39 or RV39 in the presence of DMSO or quercetin and incubated for 24 h. IL-8 in the media was measured by ELISA. Data represent average and SD calculated from 3 independent experiments (\*p 0.05, different from sham; †p 0.05, different from media or DMSO treated cells).

**Figure 2.**


Quercetin pretreatment decreases RV endocytosis and RV-induced Akt phosphorylation. BEAS-2B cells were pretreated with 10  $\mu$ M quercetin or DMSO and incubated with sham, UV-RV1B, RV1B, UV-RV39 or RV39 for 30 min at 4°C, washed to remove unbound virus and cells were further incubated for 30 min at 37°C. Endocytosed virus was detected using antibody to RV followed by flow cytometry (A and B). Data represent average and SD calculated from 3 independent experiments (\*p 0.05, different from sham; †p 0.05, different from media or DMSO treated cells). C. Cells were pretreated with DMSO or quercetin and incubated with RV1B for 30 minutes. Cells were washed and incubated with antibody to RV capsid protein (green) and EEA1 antibody (red). Nuclei were counter-stained with DAPI (blue) and cells were visualized by confocal microscopy. Arrows represent co-localization of RV1B with EEA1. Cells pretreated with quercetin were infected

with sham or RV1B and incubated for 30 min. Total cell lysates were subjected to Western blot analysis with antibodies to p-Akt or total Akt (D). Images and blot are representative of three independent experiments.




**Figure 3.**

Addition of quercetin 6 h post RV infection attenuates virus-stimulated IFN responses in airway epithelial cells. BEAS-2B cells were infected with sham, UV-RV1B, RV1B, UV-RV39 or RV1B, and incubated for 90 min. Infection medium was replaced with fresh medium and incubated for another 4.5 h. DMSO or 10  $\mu$ M quercetin was added to the cells and incubation continued for another 18 h. Levels of IFN mRNA (panels A, B and C) and IL-8 protein (D) were determined by qPCR and ELISA respectively. Expression of IFN genes were normalized to G3PDH and then expressed as fold change over sham-infected cells. Data represent average and SD calculated from 3 independent experiments (\*p 0.05, different from UV-RV in A, B and C; and different from sham in D; †p 0.05, different from media or DMSO treated cells).




**Figure 4.**

Quercetin added 6h after RV infection decreases viral load and replication. Cells were infected and treated with quercetin as described in Figure 3. Cells along with media was collected at 6 (just before adding quercetin) or 24 h (18 h after addition of quercetin or DMSO) after RV infection and infectious viral load was determined (A). Total RNA extracted from cells incubated for a total of 24 h post RV infection was used for determination of vRNA (B). The amplified qPCR product of negative strand RNA was subjected to agarose gel electrophoresis to show the amplified negative-strand vRNA (C). This result was representative of 3 experiments. (\*p 0.05, different from UV-RV in A and B; and different from sham in D; †p 0.05, different from media or DMSO treated cells).




**Figure 5.** Quercetin inhibits cleavage of eIF4GI, increases phosphorylation of eIF2 $\alpha$  and decreases levels of VP2. BEAS-2B cells were infected with RV and incubated for 90 min. Infection medium was replaced with fresh media, incubated for 4.5 h and then quercetin (10  $\mu$ M) (or DMSO) were added to the cells. After a total of 24 h, total protein was isolated and subjected to Western blot analysis using antibody to eIF4GI (A), eIF2 $\alpha$  (B) or VP2 (C). Blots are representative of 3 independent experiments.




**Figure 6.**

Withdrawal of quercetin several hours before RV infection does not limit rhinovirus infectivity *in vitro*. BEAS-2B cells were treated with DMSO or 10 μM quercetin overnight. Cells were shifted to fresh media without quercetin or DMSO and incubated for 8 h. Cells were then infected in the absence of quercetin, and incubated for 18h. Infectious viral load was determined as described in Figure 4(A). vRNA and IFN responses (B to E) were determined by qPCR. IL-8 protein levels were determined by ELISA (F). Data represent average and SD values calculated from 3 independent experiments (\*p < 0.05, different from sham infected cells).




**Figure 7.** Quercetin decreases viral load *in vivo*. C57BL/6 mice were infected with RV1B or sham by the intranasal route. Two hours after RV infection, mice were orally gavaged with quercetin (0.2 mg/mouse) or 50% propylene glycol (PG) and then once a day up to 4 days. Mice were sacrificed and infectious viral load (A) was determined by measuring CCID<sub>50</sub>/ml. Positive strand vRNA (B) was determined by qPCR and negative strand vRNA (C) was detected by qPCR followed by agarose gel electrophoresis. A and B, data represent geomean and range of data from 2 independent experiments carried out in triplicate ( $\dagger p < 0.05$ , different from mice infected with RV1B infected and treated with PG). C, PCR product from two representative animals from PG and quercetin group showing negative-strand vRNA and 18S RNA.






**Figure 8.** Quercetin decreases RV-stimulated chemokine and cytokine responses *in vivo*. Mice were infected with RV1B and treated with quercetin or vehicle as described in the Figure 5 legend. After sacrifice, mouse lungs were excised, homogenized in PBS, centrifuged and supernatant examined by ELISA (A to D) or total RNA was extracted from the lungs and mRNA expression of IFN- $\alpha$  and IFN- $\lambda_1$  was measured by qPCR (E and F). Data represent mean  $\pm$  SD calculated from triplicate experiments (\*p 0.05, different from sham; †p 0.05, different from mice infected with RV and treated with vehicle).




**Figure 9.**

Compared to vehicle-treated animals, RV-infected mice treated with quercetin show decreased airways responsiveness to methacholine. Mice were infected with RV1B and then treated with DMSO or quercetin as described above. After 1 day, mice were anesthetized and airways responsiveness to methacholine challenge was measured. (n= 3; \*p 0.05, different from sham infected mice; †p 0.05, different from mice infected with RV1B and treated with vehicle, two- way ANOVA).



**Figure 10.**

Treatment of quercetin during RV infection is necessary to limit RV-stimulated cytokine responses. Mice were treated with quercetin or vehicle as described in Figure 7 for 10 days. Forty hours after the last quercetin or vehicle treatment, mice were infected with RV1B or sham and sacrificed 24 h post-infection and lungs harvested for determination of cytokine proteins by ELISA (A to D) or IFN mRNA expression by qPCR (E to G). Data represent mean  $\pm$  SD calculated from triplicate experiments (n=5, \*p < 0.05, different from sham, ANOVA).



**Figure 11.** Treatment of quercetin during RV infection is necessary to reduce viral load and improve lung function. Mice were treated with quercetin or vehicle as described in Figure 7 for 10 days. Forty hours after the last quercetin or vehicle treatment, mice were infected with RV1B or sham. After 1 day, some mice sacrificed, lungs harvested for determination of vRNA (A) by qPCR and infectious viral load (B). Rest of the mice were anesthetized and airways responsiveness to methacholine challenge was measured (C) (n= 3–5, data in A and B represents range and median; \*p 0.05, different from sham infected mice, two- way ANOVA).