



Published in final edited form as:

J Med Chem. 2014 November 13; 57(21): 9130–9138. doi:10.1021/jm501245b.

Radiosynthesis and Evaluation of an ^{18}F -Labeled Positron Emission Tomography (PET) Radioligand for Metabotropic Glutamate Receptor Subtype 4 (mGlu₄)

Kun-Eek Kil, Pekka Poutiainen, Zhaoda Zhang, Aijun Zhu, Ji-Kyung Choi, Kimmo Jokivarsi[†], and Anna-Liisa Brownell^{*}

Athinoula A. Martinos Center for Biomedical Imaging, Department of Radiology, Massachusetts General Hospital, Charlestown, MA 02129

Abstract

Four 4-phthalimide derivatives of *N*-(3-chlorophenyl)-2-picolinamide were synthesized as potential ligands for the PET imaging of mGlu₄ in the brain. Of these compounds, *N*-(3-chloro-4-(4-fluoro-1,3-dioxo-isoindolin-2-yl)phenyl)-2-picolinamide (**3**, KALB001) exhibited improved binding affinity (IC₅₀ = 5.1 nM) compared with ML128 (**1**) and was subsequently labeled with ^{18}F . When finally formulated in 0.1 M citrate buffer (pH4) with 10% ethanol, the specific activity of [^{18}F]**3** at the end of synthesis (EOS) was 233.5±177.8 GBq/μmol (n=4). The radiochemical yield of [^{18}F]**3** was 16.4±4.8% (n=4) and the purity was over 98%. *In vivo* imaging studies in a monkey showed that the radiotracer quickly penetrated the brain with the highest accumulation in the brain areas known to express mGlu₄. Despite some unfavorable radiotracer properties like fast washout in rodent studies, [^{18}F]**3** is the first ^{18}F -labeled mGlu₄ radioligand, which can be further modified to improve pharmacokinetics and brain penetrability for future human studies.

INTRODUCTION

L-Glutamate is the most abundant excitatory neurotransmitter in the CNS of vertebrates.^{1,2} mGluRs and iGluRs are two major classes involved in glutamate signal transfer.² While iGluRs are dedicated to quick synaptic responses to open ion channels, mGluRs create slow synaptic responses and their subsequent biochemical effects.² The mGluRs belong to class C of the GPCR superfamily. mGluRs are found preferably as dimers and have a distinct large extracellular N-terminus. This extracellular N-terminal domain contains an orthosteric binding site for the endogenous ligand, formed by two hinged globular domains, referred to

^{*}Corresponding Author Information Anna-Liisa Brownell, Athinoula A. Martinos Center for Biomedical Imaging 149, 13th Street Suite 2301 Charlestown, MA 02129 United States. abrownell@partners.org. Phone: 617-726-3807.

[†]Present/Current Author Addresses

Department of Neurobiology, A.I. Virtanen Institute, University of Eastern Finland, Kuopio, Finland

Supporting Information

Chromatograms for the separation and analyses of [^3H]**1** and [^{18}F]**3** are presented in Supporting Information and the stability study and *in vitro* selectivity studies of **3** are also provided. These materials are available free of charge via the Internet at <http://pubs.acs.org/>.

The authors declare no competing financial interest.

as the Venus Flytrap Domain (VFD).³ The mGluRs can be further divided into three subgroups including eight known receptor subtypes (group I: mGlu₁ and mGlu₅, group II: mGlu₂ and mGlu₃, and group III: mGlu₄, mGlu₆, mGlu₇ and mGlu₈) based on their structural similarity, ligand specificity, and preferred coupling mechanisms.⁴ The mGluRs are involved in glutamate signal transfer in almost every excitatory synapse in CNS, and they have distinctive biodistribution in CNS depending on subtypes and subgroups.⁵ In recent years, mGlu₄ has received a lot of attention because of the potential benefits of mGlu₄ activation in several diseases such as Parkinson's Disease (PD).^{6,7} PD is caused by the degeneration of dopaminergic neurons in the basal ganglia and results in motor symptoms such as tremors and bradykinesia.⁸ As a group III mGluR, mGlu₄ interacts with the G_{ai/o} subunit of the G-protein which negatively couples with adenylate cyclase to inhibit cAMP dependent signal pathways.^{9,10} The mGlu₄ is expressed at multiple synapses throughout the basal ganglia, mainly localized presynaptically and expressed in the striatum, hippocampus, thalamus, and cerebellum.^{4,11-15} Its activation reduces neurotransmitter release, a mechanism implicated in the pathophysiology of PD. Since most orthosteric ligands of mGluRs lack clear subtype selectivity and BBB penetration, much effort has been done in the development of allosteric modulators, which target the seven-transmembrane spanning domain. In particular, mGlu₄ has received much interest following the discovery that PHCCC (*N*-phenyl-7-(hydroxyimino)cyclopropa[*b*]chromen-1*a*-carboxamide), a partially selective mGlu₄ positive allosteric modulator (PAM), demonstrated activity in models of neuroprotection and PD. Subsequent results with other PAMs of mGlu₄ have further validated the antiparkinsonian activity in animal models of PD.^{14,15} This approach has opened a new avenue for developing nondopaminergic treatments for PD and for identifying novel disease modifying therapeutics. In 2009, Engers et al. (Vanderbilt University) reported a series of *N*-phenyl-2-picolinamide derivatives as mGlu₄ PAMs, which were potent and centrally penetrating.¹⁶ Compounds **1** (*N*-(chloro-3-methoxyphenyl)-2-picolinamide, ML128) and **2** (*N*-(3-(difluoromethoxy)phenyl)-2-picolinamide) were the most potent mGlu₄ PAMs in this series, in which **1** was the first mGlu₄ PAM to show efficacy in a preclinical PD model upon systemic dosing.⁶

To better understand the role of mGlu₄ under normal and disease conditions, we are interested in developing an mGlu₄ selective radioligand for *in vivo* study. As a noninvasive medical imaging technique and powerful tool in neurological research, PET offers the possibility to visualize and analyze target receptor expression under physiological and pathophysiological conditions. Moreover, PET tracers serve as invaluable biomarkers during the development of potential therapeutic drugs. Recently, we have reported a carbon-11 labeled mGlu₄ PAM [¹¹C]**1** (*N*-(chloro-3-[¹¹C]methoxyphenyl)-2-picolinamide) as a PET radioligand.¹⁷ This compound demonstrated some promising features as a PET radioligand such as fast uptake into the brain and the specific accumulation of [¹¹C]**1** in mGlu₄-rich regions of the brain. However, in comparison to one of the best mGlu₅ PET tracers, [¹⁸F]FPEB (3-[¹⁸F]fluoro-5-(2-pyridinylethynyl)benzotrile),^{18,19} [¹¹C]**1** showed decreased retention time in the brain, which may affect the quality of the imaging. As a result of our efforts to further develop mGlu₄ PET ligands, we report here the synthesis and evaluation of [¹⁸F]**3**, the first fluorine-18 labeled PET radioligand for mGlu₄ imaging.

RESULTS AND DISCUSSION

Chemistry

Aimed at improving the potency of the previous PET ligand, we have looked for a new type of mGlu₄ PAM with improved properties for developing a new PET tracer. In recent years, a series of phthalimide derivatives of *N*-phenyl-2-picolinamides have been developed as mGlu₄ PAMs by the Merck Research Laboratory and Vanderbilt University.^{20,21} Four representative compounds (**3-6**) from this series are shown in Fig 1. ML182 (**6**) was a saturated version of the phthalimide, which proved to be orally active in the haloperidol induced catalepsy model, a well-established antiparkinsonian model. It was also shown that compounds **3** (hEC₅₀: 50.4 nM; IC₅₀: 36 nM), **4** (hEC₅₀: 43.6 nM; IC₅₀: 11 nM), and **5** (hEC₅₀: 101 nM; IC₅₀: 12nM) had enhanced activity compared with **1** (hEC₅₀: 240 nM) and gave excellent binding affinity (depicted as the IC₅₀ value) to mGlu₄. Compound **4** was radiolabeled with tritium as a tool compound for *in vitro* studies.²⁰ By checking the structures of **3-5**, it was determined that the picolinamide (left-side) was critical for the mGlu₄ potency,¹⁶ and the phthalimide group at the 4-position on the phenyl ring enhanced potency compared to **1**. In addition, although both the chloro- and bromo-substitutions at the 3-position of the phenyl ring (**4** and **5**) gave good binding affinities, the chloro-substitution was more favorable since the bromo-substitution increased molecular weight as well as clogP value, but not the potency at mGlu₄. Because of their promising affinity to mGlu₄, we have synthesized four phthalimide derivatives including **3** and **4** as well as their analogs **7** and **8** for further studies. **3**, **4**, **7**, and **8** only differ in the halide and methyl groups on the phenyl ring of the phthalimide, which may offer an appropriate site for labeling the corresponding positron emitting radionuclides such as fluorine-18, carbon-11, bromine-76, and iodine-124, respectively.

As shown in Scheme 1, compounds **3**, **4**, **7**, and the nitro-precursor **17** that is used for radiolabeling were prepared in three steps from commercially available chemicals. Iodine derivative **8** was synthesized in five steps. 2-Picolinic acid (**9**) was converted into the corresponding acid chloride and then coupled with aniline **10** to give picolinamide derivative **11** in 41% yield. The nitro group in **11** was subsequently reduced by hydrogenation to afford aniline **12** in 91% yield. Aniline **12** was coupled with the phthalic anhydride derivatives **13** through **16** in acetic acid to give the corresponding phthalimides derivatives, **3**, **4**, **7**, and **17** in 78%, 66%, 92%, and 80% yields, respectively. To prepare iodine derivative **8**, bromo-derivative **7** was converted into trimethyltin derivative **18**, which subsequently reacted with iodine to obtain **8** in 7% yield.

Binding Affinity to mGlu₄

Since we had studied [¹¹C]**1** as a PET ligand and there was no report on its affinity to mGlu₄, we would like to directly compare the affinity of the phthalimide derivatives **3**, **4**, **7** and **8** with compound **1**. Thus, we performed radiolabeling of compound **1** with tritium for a competitive binding assay (Scheme 2). The chromatography of [³H]**1** radiosynthesis is shown in Supporting Information A.

Table 1 presents the IC₅₀ values of compounds **1-4**, **7** and **8** obtained from *in vitro* competitive binding assay using [³H]**1**. The results show that the lead compounds from of *N*-phenyl-2-picolinamide series **1** and **2** have IC₅₀ values of 12.8 nM and 9.9 nM, respectively. Compound **2** has been used for *in vivo* mGlu₄ blocking experiments for [¹¹C]**1**.¹⁷ The phthalimide derivatives, **3** (5.1 nM) and **4** (4.2 nM) gave improved affinity compared to **1** while compounds **7** (11.8 nM) and **8** (26.3 nM) showed similar or less affinity to mGlu₄ compared to **1**. This result shows that **3** and **4** have similar affinity while Merck's patent²⁰ presents that **4** (11 nM) has better affinity than **3** (36 nM), which may be caused from using cells expressing different mGlu₄ receptors. We used the rat mGlu₄ while Merck employed the human mGlu₄. The results verified that **3** and **4** are more potent compounds for developing an mGlu₄ PET ligand. The results also indicate that these phthalimide compounds interact with the same allosteric site in mGlu₄ where **1** binds because the quantitative specific binding curves of these compounds show complete replacement of the radioligand at a higher concentration (Fig. 2). In addition, the selectivities of **3** to mGlu₁, mGlu₅ and mGlu₈ were determined by the functional assays, which showed little activity against to these mGluRs (Supporting information B).^{22, 23}

Based on their affinity, both compounds **3** and **4** can be the ideal candidates for developing PET radioligands. Nonetheless, fluorine-18 is often the radionuclide of choice for both its physical and nuclear characteristics. Its half-life is long enough to carry out relatively extended imaging protocols when compared to what is possible with carbon-11, which facilitates kinetic studies and high-quality metabolic and plasma analyses. Therefore, we have selected compound **3** for radiolabeling and *in vivo* PET imaging.

Radiochemistry

As shown in Scheme 3, [¹⁸F]**3** was prepared from the reaction of **17** in DMSO with non-carrier-added (nca) [¹⁸F] fluoride in the presence of kryptofix222 (K222) and potassium carbonate at 120°C for 5 min. It was necessary to use an adequate amount of **17** (6 mg) for this reaction because our result showed that K222 activated carbonate anion and could cause the decomposition of substrate **17**. The radioactive product was purified by semi-preparative HPLC eluting with a 0.1% formic acid solution of water and acetonitrile (52:48) at a flow rate of 4 ml/min. [¹⁸F]**3** was eluting at 17.4-18.8 min and was well separated from starting material **17** (t_R:19.6-22.4 min) as well as other impurities (Supporting Information C). It was subsequently found that the radiochemical purity of [¹⁸F]**3** was only 79% when the purified [¹⁸F]**3** was first formulated in a neutral 10% ethanol-saline solution (Supporting Information D). This result indicates that the phthalimide derivative was not very stable and underwent hydrolysis in a neutral condition.²⁴ To confirm this issue, we studied the stability of **3** in both neutral and acidic conditions by using a LC-MS method. When **3** was dissolved in a neutral HPLC solution (0.1 M ammonium formate water-acetonitrile solution) and analyzed by using the same neutral eluants, two side products were detected in which one was **12** and the other was the hydrolysis product **20a** and/or **20b** (Supporting Information E). Conversely, the stability test in acidic media gave minimal hydrolysis products. Thus, [¹⁸F]**3** was purified and formulated in acidic media after radiosynthesis to minimize unwanted hydrolysis. When finally formulated in 0.1 M citrate buffer (pH4) with 10% ethanol, the specific activity of [¹⁸F]**3** at the end of synthesis (EOS) was 233.5±177.8 GBq/μmol (n=4).

The average decay corrected radiochemical yield (RCY) of [^{18}F]**3** was $16.4\pm 4.8\%$ ($n=4$). The radiochemical purity of [^{18}F]**3** was over 98%.

QC analysis using analytical HPLC and radio-TLC by co-injection with nonradioactive **3** has confirmed the isolated radioactive product as [^{18}F]**3** (supporting information F). Both methods validated that the radiochemical purity of [^{18}F]**3** was over 98%.

PET Imaging of [^{18}F]**3**

The characterization of [^{18}F]**3** was first conducted with rat (male Sprague Daley) studies. These studies demonstrated that [^{18}F]**3** crossed BBB and occupied brain areas known to express mGlu₄. TACs showed fast uptake and washout in different brain regions within 20 min (Fig. 3a). These results are similar to our previous PET studies with [^{11}C]**1**.¹⁷

Pre-treatment with 10 mg/kg of **2**·HCl as an mGlu₄ blocking agent^{16,17} and 10 mg/kg of 3-((2-methyl-1,3-thiazole-4-yl)ethynyl)pyridine (MTEP) as a selective mGlu₅ blocking agent^{25,26} resulted in modest blocking effects in rat brains of 10-20% and 0-5% reduced uptake, respectively. (Fig. 3b)

The low blocking effects induced by **2** can be attributed to insufficient dosage and fast pharmacokinetics behavior of **2** in brain.¹⁶ 10 mg/kg of **2**·HCl may not be enough to achieve noticeable blocking effects in the brain. The low blocking effects can also be due to fast pharmacokinetic properties of **2** in the brain.¹⁶ The time difference between the injection of the blocking agent and radiotracer was 5 min, and some of blocking agent **2** might have washed off before [^{18}F]**3** injection. Therefore, the current results might be improved to obtain better specificity and selectivity with further adjusted blocking.

In addition, the arterial plasma analysis (Fig. 3c) showed that after the intravenous bolus injection, about 90% of [^{18}F]**3** was quickly metabolized within 5 min. It is important to improve its metabolic stability.

After the characterization of [^{18}F]**3** in rats studies, combined PET-MRI studies showed similar results in the monkey brain with high accumulation of [^{18}F]**3** in the regions such as the striatum, thalamus, hippocampus, and cerebellum where previous literature has reported high expression of mGlu₄ (Fig. 4a).^{4,15} T1-weighted MRI study (Fig. 4c) and fused PET-MRI studies (Fig. 4b) illustrated anatomical details and the regional distribution of [^{18}F]**3**.

Time activity curves (TAC)s derived from the different monkey brain areas showed fast accumulation of [^{18}F]**3** with the highest accumulation about 3 min after the administration of the radioligand. About 50% of the activity stayed for 20 min followed by slower washout until 40 min, allowing suitable retention time for binding. The binding potential determined by the metabolite corrected plasma input function and volume distribution model was highest in the thalamus (2.35) followed by the putamen (2.13), the caudate (1.91), the prefrontal cortex (1.47), and the cerebellum (1.19). These studies confirm that the cerebellum cannot be used as a reference tissue in quantitative modeling.

The fast pharmacokinetics of [^{18}F]**3** in the brain can be also related to metabolic instability of the radiotracer. The percent of parent [^{18}F]**3** in plasma was quickly dropped within 5 min

in both rats and the monkey (Fig. 3c and Fig. 5). However, [¹⁸F]**3** showed better permeability in the monkey brain than in the rat brain and was retained in the brain until 40 min (Fig. 3a and Fig. 5).

CONCLUSION

Four 4-phthalimide derivatives of *N*-(3-chlorophenyl)-2-picolinamide **3**, **4**, **7**, and **8**, and precursor **17** were synthesized by three or five steps of syntheses from commercially available chemicals. Compounds **3**, **4**, **7**, and **8** were studied for their binding affinity using rat mGlu₄ transfected Chinese hamster ovary (CHO) cells. Among these compounds, **3** and **4** exhibited the highest binding affinity, which demonstrated improved affinity compared to the previously reported mGlu₄ PET ligand, [¹¹C]**1**. **3** was selected and labeled with fluorine-18 since fluorine-18 has been the choice of radionuclide for PET.

One step labeling reaction with fluorine-18 was successfully performed after careful optimization of reaction conditions with 16.4±4.8% of RCY (n = 4) and 233.5±177.8 GBq/μmol of specific activity (n = 4). [¹⁸F]**3** was formulated with 10% ethanol in pH4 citrate buffer to achieve over 98% radiochemical purity. *In vivo* imaging studies in the monkey showed that the radiotracer quickly penetrated the brain with the highest accumulation in the brain areas known to express mGlu₄. In the rat studies, the fast accumulation was followed with fast washout from the brain within 20 min while it was 40 min in the monkey study. Even though [¹⁸F]**3** did not improve the problems of fast washout in rodent studies which we reported with [¹¹C]**1**, it might be a more promising radiotracer to further development for use in monkey and possibly human studies since it displayed higher peak uptake in the brain and it was retained in the brain for a longer period. *In vitro* tests also indicated that [¹⁸F]**3** would give better results with mGlu₄ human cell lines than [¹¹C]**1**.^{6,16,20} Despite of some unfavorable properties as a radiotracer such as fast washout in rodent studies, [¹⁸F]**3** is the first ¹⁸F-labeled radioligand and can be further modified to improve pharmacokinetic properties and brain penetrability as a candidate for future human studies.

EXPERIMENTAL SECTION

Animal Procedures

The animal studies were approved and done under the strict supervision of the Subcommittee on Research Animals of the Massachusetts General Hospital and Harvard Medical School and performed in accordance with the Guide of NIH for the Care and Use of Laboratory Animals.

Materials and General Methods

Unless otherwise specified, all chemicals were purchased from Sigma-Aldrich (St. Louis, MO). The reactions were monitored by TLC using aUV lamp monitored at 254 nm. If necessary, the reactions were also checked by LC-MS using the Agilent 1200 Series HPLC system coupled with a multi-wavelength UV detector and a6310 Ion Trap Mass Spectrometer (Santa Clara, CA), equipped with an Agilent Eclipse C₈ analytical column (150 mm × 4.6 mm, 5 μm), eluted with a 0.1% formic acid solution of water (A) and acetonitrile (B). Flash chromatography was performed with a CombiFlash R_f Purification

System (Teledyne Isco) using a Silica ReadySep R_f Column. Hydrogenation reaction was performed using a Parr Shaker Hydrogenation Apparatus (Moline, IL). The products were identified by LC-MS and ¹H-NMR and ¹³C-NMR using a Varian 500 MHz Spectrometer. ¹⁹F-NMR was used to identify the fluorine compound. All NMR samples were dissolved in chloroform-*d* (CDCl₃) containing tetramethylsilane as a reference standard. Chemical shifts were expressed as ppm and calculated downfield or up field from the NMR signal of reference standard. *J* was expressed as Hz, and its splitting patterns were reported as s, d, t, q, or m. HRMS was obtained from the High Resolution Mass Spectrometry Facility at the University California, Riverside using electrospray ionization (ESI)/atmospheric pressure chemical ionization (APCI) technique (Agilent Time of Flight (TOF) LC-MS). mp was measured using a Mettler MP50 Melting Point System. Unless otherwise specified, the purities of all new compounds were over 95% determined by HPLC.

The mGlu₄ blocking agent **2**, and its hydrochloride salt were prepared by a published method.¹⁷ [³H]Iodomethane (1 Ci/ml in DMF) was purchased from American Radiolabeled Chemicals Inc. and used without further purification.

The National Institute of Mental Health's Psychoactive Drug Screening Program, Contract # HHSN-271-2008-00025-C (NIMH PDSP), generously performed selectivity tests for mGlu₁ and mGlu₅ using intracellular inositol phosphates (IPs) accumulation assay with CHO cells that stably expressed mGlu_{1a} and mGlu_{5a}. Detailed procedures of selectivity tests were described in previous literature.^{17,22,23} For the selectivity test for mGlu₈, split luciferase biosensor cAMP assay was performed using mGlu₈ transfected CHO cells.^{22,23}

Syntheses of Phthalimide Derivatives **3**, **4**, **7**, and **17**

A solution of **12** and phthalic anhydride derivatives (**13**, **14**, **15**, and **16**) in acetic acid was heated to 100°C for 5 h. After acetic acid was removed at reduced pressure, residue was recrystallized in a solution of ethyl acetate and hexane to give the corresponding phthalimide derivatives **3**, **4**, **7**, and **17** as brown powders, respectively.

***N*-(3-Chloro-4-(4-fluoro-1,3-dioxo-isoindolin-2-yl)phenyl)-2-picolinamide (3)**

12 (0.20 mmol, 50 mg), 3-fluorophthalic anhydride (**13**, 0.24 mmol, 40 mg) and acetic acid (4.0 ml) were used to afford **3** (62 mg, 78% yield); mp: 218-220°C. ¹H NMR (500 MHz, CDCl₃): δ 10.21 (s, 1H), 8.64 (d, *J* = 4.5 Hz, 1H), 8.31 (d, *J* = 8.0 Hz, 1H), 8.18 (d, *J* = 2.0 Hz, 1H), 7.95 (t, *J* = 8.0 Hz, 1H), 7.80-7.83 (m, 3H), 7.53 (dd, *J* = 7.5 Hz, 5.0 Hz, 1H), 7.48 (t, *J* = 8.0 Hz, 1H), 7.35 (d, *J* = 8.5 Hz, 1H). ¹³C NMR (125 MHz, CDCl₃): δ 165.97 (d, *J* = 2.6 Hz), 163.71, 162.57, 158.30 (d, *J* = 265.6 Hz), 149.46, 148.43, 140.23, 138.23, 137.38 (d, *J* = 6.8 Hz), 134.39, 134.17, 131.28, 127.27, 124.92, 123.18 (d, *J* = 20.5 Hz), 122.99, 121.31, 120.52 (d, *J* = 3.9 Hz), 118.88, 118.16 (d, *J* = 1.3 Hz). ¹⁹F NMR (470 MHz, CDCl₃) δ -111.6 (t, *J* = 5.9 Hz). HRMS *m/z* calcd for C₂₀H₁₂ClFN₃O₃ (M+H)⁺, 396.0546; found 396.0554.

***N*-(3-Chloro-4-(4-methyl-1,3-dioxo-isoindolin-2-yl)phenyl)-2-picolinamide (4)**

12 (0.40 mmol, 100 mg), 3-methylphthalic anhydride (**14**, 0.42 mmol, 39 mg) and acetic acid (8.0 ml) were used to afford **4** (66 mg, 66% yield); mp: 184-185 °C. ¹H NMR (500

MHz, CDCl₃): δ 10.19 (s, 1H), 8.63 (d, J = 5.0 Hz, 1H), 8.31 (d, J = 8.0 Hz, 1H), 8.17 (d, J = 2.0 Hz, 1H), 7.94 (td, J = 8.0 Hz, 1.0 Hz, 1H), 7.81 (m, 2H), 7.66 (t, J = 8.0 Hz, 1H), 7.52-7.57 (m, 2H), 7.35 (d, J = 8.5 Hz, 1H), 2.76 (s, 3H). ¹³C NMR (125 MHz, CDCl₃): δ 167.91, 167.19, 162.52, 149.51, 148.41, 139.95, 139.07, 138.19, 137.15, 134.29, 134.23, 132.71, 131.35, 128.99, 127.21, 125.54, 122.93, 121.93, 121.24, 118.78, 18.13. HRMS m/z calcd for C₂₁H₁₅ClN₃O₃(M+H)⁺, 392.0796; found 392.0805.

***N*-(3-Chloro-4-(4-bromo-1,3-dioxo-isoindolin-2-yl)phenyl)-2-picolinamide (7)**

12 (2.02 mmol, 0.500 g), 3-bromophthalic anhydride (**15**, 2.22 mmol, 0.504 g) and acetic acid (10 ml) were used to afford **7** (844 mg, 92%); mp: 187-189 °C. ¹H NMR (500 MHz, CDCl₃): δ 10.21 (s, 1H), 8.62 (d, J = 4.5 Hz, 1H), 8.30 (d, J = 7.5 Hz, 1H), 8.17 (d, J = 1.0 Hz, 1H), 7.92-7.95 (m, 3H), 7.79 (dd, J = 8.5 Hz, 1.5 Hz, 1H), 7.65 (t, J = 7.5 Hz, 1H), 7.51-7.54 (m, 1H), 7.35 (d, J = 8.5 Hz, 1H). ¹³C NMR (125 MHz, CDCl₃): δ 165.47, 165.08, 162.55, 149.42, 148.41, 140.17, 139.71, 138.21, 135.63, 134.43, 134.10, 131.21, 129.81, 127.25, 125.02, 123.33, 122.96, 121.24, 119.55, 118.82. HRMS m/z calcd for C₂₀H₁₂ClBrN₃O₃ (M+H)⁺, 455.9745; found 455.9742.

***N*-(3-Chloro-4-(4-nitro-1,3-dioxo-isoindolin-2-yl)phenyl)-2-picolinamide (17)**

12 (0.606 mmol, 0.150 g), 3-nitrophthalic anhydride (**16**, 0.666 mmol, 0.129 g) and acetic acid (6 ml) were used to afford **17** (0.204 g, 80% yield); mp: 199-201 °C. ¹H NMR (500 MHz, CDCl₃): δ 10.23 (s, 1H), 8.65 (d, J = 5.0 Hz, 1H), 8.32 (d, J = 8.0 Hz, 1H), 8.25 (d, J = 7.0 Hz, 1H), 8.20-8.22 (m, 2H), 8.01 (t, J = 7.8 Hz, 1H), 7.96 (td, J = 8.0 Hz, 1.5 Hz, 1H), 7.82 (dd, J = 8.8 Hz, 2.3 Hz, 1H), 7.54 (dd, J = 7.0 Hz, 5.0 Hz, 1H), 7.37 (d, J = 8.5 Hz, 1H). ¹³C NMR (125 MHz, CDCl₃): δ 164.62, 162.61, 161.68, 149.41, 148.45, 145.87, 140.50, 138.26, 136.14, 134.25, 134.05, 131.17, 129.41, 128.03, 127.31, 124.50, 124.00, 123.02, 121.31, 118.91. HRMS m/z calcd for C₂₀H₁₂ClN₄O₅(M+H)⁺, 423.0491; found 423.0487.

***N*-(3-Chloro-4-(4-iodo-1,3-dioxo-isoindolin-2-yl)phenyl)-2-picolinamide (8)**

Iodine (0.116 mmol, 29.4 mg) was added to a solution of **18** (72.5 μ mol, 39.2 mg) in DCM (1.4 ml), and stirred at room temperature for 2 h. The solution was quenched with 5% sodium metabisulfite solution until its color disappeared. The solution was then extracted with ethyl acetate three times. Organic layers were collected, dried over anhydrous sodium sulfate (Na₂SO₄), filtered, and evaporated. Residue was separated with 25% ethyl acetate in hexane by flash column chromatography to give **8** as a yellowish solid (2.5 mg, 7% yield); mp: 140-142 °C. ¹H NMR (500 MHz, CD₃CN): δ 10.37 (s, 1H), 8.72 (d, J = 4.5 Hz, 1H), 8.30 (d, J = 8.0 Hz, 1H), 8.26-8.27 (m, 2H), 8.05 (t, J = 7.5 Hz, 1H), 7.98 (d, J = 7.5 Hz, 1H), 7.92 (dd, J = 8.5 Hz, 2.0 Hz, 1H), 7.64 (dd, J = 8.0 Hz, 4.0 Hz, 1H), 7.59 (t, J = 7.8 Hz, 1H), 7.48 (d, J = 9.0 Hz, 1H). HRMS m/z calcd for C₂₀H₁₂ClN₃O₃I (M+H)⁺, 503.9606; found 503.9417.

***N*-(3-Chloro-4-nitrophenyl)-2-picolinamide (11)**

Thionyl chloride (163 mmol, 11.8 ml) was added to 2-picolinic acid (**9**, 40.6 mmol, 4.00 g) in benzene (60 ml), and the mixture was refluxed for 2 h and then the solvent and extra

thionyl chloride were removed by vacuum distillation. The resulting 2-picolinic acid chloride was isolated as a grey powder. A solution of 3-chloro-4-nitroaniline (**6**, 10.2 mmol, 1.75 g) (Alfa Aesar) and triethylamine (122 mmol, 17.0 ml) in 30 ml of THF was then added to the grey salt powder in 30 ml of THF. The mixture was refluxed for 2 h and was then quenched with saturated ammonium chloride solution. The reaction mixture solution was extracted with ethyl acetate three times and dried over anhydrous Na₂SO₄, filtered, and evaporated. The residue was separated by flash chromatography with eluants of ethyl acetate (20%) and hexane (80%) to give the crude product as a solid, which was recrystallized in a solution of ethyl acetate and hexane to give **11** as a white crystal (1.153 g, 41% yield); mp: 179-180°C. ¹H NMR (500 MHz, CDCl₃): δ 10.35 (s, 1H), 8.65 (d, *J* = 5.0 Hz, 1H), 8.31 (d, *J* = 8.0 Hz, 1H), 8.15 (d, *J* = 2.0 Hz, 1H), 8.05 (d, *J* = 9.0 Hz, 1H), 7.97 (td, *J* = 8.0 Hz, 1.0 Hz, 1H), 7.57 (dd, *J* = 7.0 Hz, 5.0 Hz, 1H). ¹³C NMR (125 MHz, CDCl₃): δ 162.77, 148.90, 148.53, 143.16, 142.61, 138.39, 129.45, 127.69, 127.66, 123.12, 122.04, 117.87. HRMS *m/z* calcd for C₁₂H₉ClN₃O₃ (M+H)⁺, 278.0327; found 278.0329.

***N*-(3-Chloro-4-aminophenyl)-2-picolinamide (12)**

Compound **11** (3.96 mmol, 1.10 g) and 10% palladium/charcoal (0.220 g) were added to a solution of methanol (70 ml) and ethyl acetate (70 ml). The mixture was shaken under 3 atm of hydrogen for 20 h at room temperature. The solution was filtered and evaporated. The residue was separated by flash chromatography with eluants of 30% ethyl acetate and 70% hexane to give a crude product, which was recrystallized in a solution of ethyl acetate and hexane to afford **12** as a yellowish solid (0.889 g, 91% yield); mp: 144-145°C. ¹H NMR (500 MHz, CDCl₃): δ 9.85 (s, 1H), 8.60 (d, *J* = 5.0 Hz, 1H), 8.29 (d, *J* = 8.0 Hz, 1H), 7.91 (td, *J* = 8.0 Hz, 1.0 Hz, 1H), 7.84 (d, *J* = 2.0 Hz, 1H), 7.48 (dd, *J* = 7.5 Hz, 5.0 Hz, 1H), 7.44 (dd, *J* = 9.0 Hz, 2.0 Hz, 1H), 6.80 (d, *J* = 8.5 Hz, 1H), 4.00 (s, 2H). ¹³C NMR (125 MHz, CDCl₃): δ 161.98, 150.12, 148.27, 140.13, 137.98, 129.87, 126.67, 122.60, 121.56, 120.11, 119.69, 116.28.

***N*-(3-Chloro-4-(4-trimethylstannyl-1,3-dioxo-isoindolin-2-yl)phenyl)-2-picolinamide (18)**

Bis(trimethyltin) (0.40 mmol, 84 μl) was added to a mixed solution of **7** (0.18 mmol, 80 mg) and tetrakis(triphenylphosphine)palladium(0) (1.8 μmol, 2.0 mg) in degassed toluene (2 ml) at room temperature, and then was refluxed for 8 h. The resulting mixture was diluted with saturated ammonium chloride solution and extracted three times with ethyl acetate. The organic layers were combined, dried over anhydrous Na₂SO₄, filtered, and evaporated. The residue was separated by flash chromatography with 10% ethyl acetate in hexane to give **18** as a white solid (44 mg, 46% yield); mp: 213-214 °C. ¹H NMR (500 MHz, CDCl₃): δ 10.19 (s, 1H), 8.63 (d, *J* = 4.0 Hz, 1H), 8.30 (d, *J* = 8.0 Hz, 1H), 8.18 (s, 1H), 7.90-7.96 (m, 3H), 7.80 (d, *J* = 8.0 Hz), 7.72 (t, *J* = 5.5 Hz, 1H), 7.38 (d, *J* = 8.5 Hz, 1H), 0.41 (s, 9H). ¹³C NMR (125 MHz, CDCl₃): δ 169.16, 167.78, 162.48, 149.46, 148.39, 142.39, 142.09, 139.92, 138.17, 134.08, 133.28, 131.75, 131.23, 127.20, 125.49, 124.02, 122.92, 121.27, 118.78, -8.37. HRMS *m/z* calcd for C₂₃H₂₁ClN₃O₃¹²⁰ Sn (M+H)⁺, 542.0288; found 542.0294.

Radiosynthesis of [³H]1

[³H]Iodomethane (1 Ci/ml in DMF) was added to a DMF solution of *N*-(4-chloro-3-hydroxyphenyl)picolinamide (**19**, 1.5 mg, 6.0 μmol) and 5 M potassium hydroxide solution (3.0 μl, 15 μmol). The resulting mixture was heated at 90°C for 10 min and diluted with 1.0 ml of HPLC solvents. Then, the aliquot was injected into HPLC equipped with a Gemini-NX C₁₈ Semi-Preparative Column (250 mm × 10 mm, 5 μm, Phenomenex Inc.), a flow scintillation detector, and an internal UV detector eluting with a solution of 55% acetonitrile and 45% 0.1 M ammonium formate at a flow rate of 4 ml/min. An eluant containing [³H]1 was collected between 11–13 min. The radioactive product was diluted with 30 ml of water and passed through a C₁₈ Sap-Pak Plus followed by additional wash with 5 ml of sterile water. [³H]1 was then eluted from cartridge with ethanol. The final product was diluted with 9 volume of sterile saline to make a 10% ethanol solution in saline for cell studies.

Preparation of mGlu₄ Transfected CHO Cells

The vector of mGlu₄ from rat was obtained as a gift from Drs. Tanabe and Nakanishi's laboratory (Osaka Bioscience Institute, Osaka, Japan), and its structure was described in their previous literature.²⁷ To extract an mGlu₄ cDNA insert out of its backbone, pBluescript II KS(+), EcoR1 (New England Biolabs) was treated and incubated for 4 h at 37 °C. Then electrophoresis was performed and the mGlu₄ insert (3704 bp) was isolated from its backbone (2954 bp) using a gel purification kit (Qiagen Inc.) following the procedure provided by the kit. The target backbone, pcDNA, was also digested by the same procedure. After their concentrations were measured using a NanoDrop (Thermo Scientific), pcDNA (30 fmol) and the mGlu₄ insert (90 fmol) were mixed and incubated at room temperature for 2 h mediated by T4 Ligase (Invitrogen® Life Technology). After the ligase was deactivated by heating, the combined vectors were incorporated into DH5α-E competent cells (Invitrogen® Life Technology) by electroporation. The aliquot was transferred to a LB agar dish containing ampicillin and grown at 37 °C overnight. Some clones were selected and amplified in LB media containing 0.1% ampicillin. Successful clones were screened by electrophoresis and DNA sequencing. They continued to culture in LB broth (1 L) containing 0.1% ampicillin stock solution. The resultant vectors containing mGlu₄ cDNA was obtained using a Maxi-Prep Column Kit (Qiagen Inc.) and its standard procedure.

The mGlu₄ DNA and Plus™ Reagent (Invitrogen® Life Technology) were mixed in reduced serum media (Opti-MEM® Life Technology) and added to the reduced serum media containing Lipofectamine® (Invitrogen® Life Technology). The resulting aliquot was incubated for 15 min at room temperature. Meanwhile, CHO cells were loaded to a six-wall plate and cultured overnight until confluence at 37 °C in the culture media. After the culture media in the six-wall plate was replaced with reduced serum media, the mGlu₄ aliquot containing Lipofectamine and Plus reagent was loaded to the six-wall plate. Transfection was performed at 37 °C for 3 h. The media was replaced with culture media containing G418 to select and grow only transfected cells. The cells in each wall continued to grow until non-transfected cells died. The selection media was replaced every three days during culturing. The successful colonies were cropped and moved to bigger culture bottles. These cells continued to grow until confluence by changing media containing G418 every three days. Each colony was further selected to identify successful colonies by western blot.

In Vitro mGlu₄ Binding Assay

CHO cells expressing mGlu₄ were used for all binding assay protocols. For competitive binding, 10 nM of [³H]1 was used together with increased concentrations of test compounds ranging from 0.01 nM to 10 μM. Each test tube contained 50,000 freshly harvested CHO cells expressing mGlu₄ cultured in HAMs cell culture medium, penicillin streptomycin (100 units), and 1 mM G418. [³H]1 (10 nM) was added to the cell extract with and without test compounds on ice and then incubated for 30 min at room temperature. Samples were centrifuged at 1200 rpm for 10 min at 4°C and washed three times with a cold cell culture medium as a washing buffer. Samples were lysed by adding 100 μl of 0.5% NaOH and heated using a heating block (56°C, 30 min). Samples were cooled with an ice bath and transferred to Solvent-Saver™ scintillation vials (VWR International LLC.). To obtain binding parameters, the scintillation liquid (Perkin Elmer, Optima Gold) was added prior to counting with a scintillation counter (Packard TriCarb Model, 1 min/vial). Nonspecific binding was determined using 10 μM nonradioactive **1** and specific binding was determined by extracting the nonspecific binding from total binding. All measurements were done in triplicate and analyzed with GraphPad Prism software (GraphPad Software Inc.).

Radiosynthesis of [¹⁸F]3

[¹⁸F]Fluoride was generated by a Siemens Eclipse HP 11 MeV Cyclotron (Malvern, PA) using [¹⁸O]-enriched water (Isoflex Isotope, San Francisco, CA) with proton bombardment. Fluorine-18 labeling chemistry was carried out in a Siemens Explora® GN synthetic module. A [¹⁸F]fluoride reagent was passed through a QMA Sep-Pak Cartridge (Waters, Milford, MA) to trap [¹⁸F]fluoride ions. A solution of potassium carbonate (1 mg) and K222 (3 mg) in acetonitrile and water (1:1, 0.4 ml) was passed through the cartridge to elute [¹⁸F]fluoride ions. The solvents were evaporated at 115°C in a stream of nitrogen. To remove water completely, 1 ml of acetonitrile was added and evaporated twice. **17** (6 mg) in 0.6 ml of DMSO was added to the residue and the mixture was heated to 120°C for 5 min. The resulting mixture was cooled to 100°C and 1.5 ml of the HPLC solvent was added. The mixture was then purified by HPLC (Agilent 1100 series equipped with UV detector and radioactivity detector) using an Eclipse XDB-Phenyl Semi-Preparative Column (250 mm × 10 mm, 5 μm) and eluting with a 0.1% formic acid solution of water and acetonitrile (52:48) at a flow rate of 4 ml/min. The fraction containing [¹⁸F]**3** was diluted with a 0.1 M citrate buffer solution (pH4) to 20 ml and loaded on a C₁₈ Sep-Pak column. The final product was obtained after eluting 1 ml of ethanol through the C₁₈ Sep-Pak column and the collected product was diluted with a sterile 0.1 M citrate buffer solution (pH4) to make a 0.1 M citrate buffer solution with 10% ethanol.

QC Analysis of [¹⁸F]3

A purified radioactive aliquot (10 μl) and nonradioactive standard (**3**) were injected into the HPLC (Agilent 1100 series equipped with a UV detector monitored at 254 nm and a Ramsey Model 105S Single Channel Radiation Detector (Berkeley, CA)) using an Eclipse XDB-Phenyl Analytical Column (150 mm × 4.6 mm, 5 μm, Agilent) and eluting with a solution of acetonitrile and 0.1% formic acid (55:45) at a flow rate of 1 ml/min.

An AR-2000 radio-TLC Imaging Scanner (Bioscan Inc.) was also used to check radio-TLC for further confirmation of radioactive product. The radioactive aliquot was co-spotted with the nonradioactive **3** (in acetonitrile) on TLC and developed with a solution of ethyl acetate and hexane (1:1). The same R_f values of the radioactive spot and UV active spot were confirmed by the radio-TLC scanner and UV lamp at 254 nm, respectively.

Specific Activity of [^{18}F]**3**

Specific activity was measured by the published method.¹⁷ The radioactivity of the aliquot was measured and corrected to the time point of the EOS. The corresponding mass was measured by a calibration curve which showed the linear relationship between the molar amount of **3** and UV integration. The calibration curve was previously established with five different standard concentrations of **3**. The specific activity was calculated and expressed as GBq/ μmol .

PET Imaging of [^{18}F]**3** in Rats

PET imaging studies were conducted using seven normal Sprague Dawley male rats. The rats were anesthetized with isoflurane/nitrous oxide (1.0-1.5% isoflurane, with oxygen flow of 1-1.5 L/min), and a catheter was installed in their tail veins for the administration of [^{18}F]**3**. Dynamic volumetric PET data was acquired with a PET-CT scanner for 60 min (Triumph-II Preclinical Imaging System, Tri-Foil imaging, Northridge, CA). Vital signals such as heart and respiration rates were monitored during the scanning period. PET data acquisition started immediately after the administration of [^{18}F]**3** (37 MBq) followed by CT imaging to obtain anatomical information and data for attenuation correction of PET data. PET data was processed using a Maximum-Likelihood Expectation-Maximization (MLEM) algorithm with 30 iterations for dynamic volumetric images and were corrected for uniformity, scatter, and attenuation. The CT data was processed by a modified Feldkamp algorithm using matrix volumes of $512 \times 512 \times 512$ with a pixel size of 170 μm . Co-registration of CT and PET images and analysis of PET images were performed using PMOD3.2 software (PMOD Technology, Zurich, Switzerland).

For blocking studies, the same imaging protocol was used for baseline studies. For mGlu₄ blocking, 10 mg/kg of **2**·HCl salt dissolved in a saline solution with 10% ethanol and 10% Tween20 was injected 5 min before i.v. administration of [^{18}F]**3** in four normal male rats. For mGlu₅ blocking, 10 mg/kg of MTEP·HCl (abcam Inc.) in 0.9% saline was intravenously administered to three normal male rats 5 min before [^{18}F]**3** injection.

Blood Metabolite Studies

Blood metabolite studies were performed by the procedure reported in previous literature.¹⁷ Blood samples were taken from rats ($n = 2$) at 10, 30, and 60 min after [^{18}F]**3** injection. Each blood sample was centrifuged for 3 min to separate plasma and blood coagulants. The plasma (0.3 ml) was retrieved, and 0.3 ml of acetonitrile was added to the plasma. The suspended solution was vortexed, and centrifuged for 3 min to precipitate protein and to obtain protein-free plasma as supernatant. The resulting supernatant was divided 2 parts. One part was used for measuring the total radioactivity and another part (0.3 ml) was loaded to a Strata-X C₁₈ cartridge (500 mg, Phenomenex Inc.) that had been purged with water. A

series of elutions were performed using using 0.1% TFA/water and 0.1% TFA/acetonitrile in quantities: 95:5, 90:10, 85:15, 80:20, 70:30, 60:40, 30:70, 100% of 0.1% TFA/acetonitrile at a volume of 4 mL. The eluted fractions were collected in eight test tubes. A control experiment by injection of a small amount of [¹⁸F]3 onto a test cartridge elution described above indicated that the parent [¹⁸F]3 was collected in the seventh test tube exclusively. The radioactivities were measured using a 2480 Wizard² Automatic Gamma Counter (PerkinElmer Inc., Waltham, MA).

PET Imaging of [¹⁸F]3 in a Rhesus Monkey

A female rhesus monkey was restricted from food for 12 hrs before the PET-MR imaging. At first, 10 mg/kg of ketamine was administered intramuscularly. Anesthesia was maintained at 1-4% isoflurane (Forane) in a mixture of oxygen and nitrogen. The monkey was intubated and catheters were introduced for radiotracer injection and a radial arterial line for blood drawing needed for the input function and metabolite analysis. MR-PET images were obtained using an integrated 3T MR-BrainPET camera (Siemens, Munich, Germany) with a resolution of 2.5 mm and an axial field of view (FOV) of 19.25 cm and a transaxial FOV of 30 cm. A multi-echo Magnetization-Prepared Rapid Gradient-Echo (MEMPRAGE) sequence was initiated to acquire MRI data for attenuation correction and anatomical coregistration. After the injection of 189 MBq of [¹⁸F]3 (37 ng/kg) in a 10% ethanol solution in isotonic saline, dynamic PET data was acquired for 90 min. For quantitative analyses, blood samples were collected every 10 sec for the first 2 min after the administration of [¹⁸F]3 and subsequently obtained at the following time points during the scan: 3, 5, 10, 15, 20, 30, 45, 60, and 90 min.

The images were reconstructed using an ordinary Poisson ordered-subset expectation maximization (OP-OSEM) 3D algorithm from prompt and random coincidences, normalization, attenuation and scatter coincidences sinograms using 16 subsets and 6 iterations. The reconstructed volume consisted of 153 slices with 256×256 pixels (1.25×1.25×1.25 mm³). Regions of interest (ROI)s were drawn on all coronal levels, where structures were observed anatomically (fused data with MRI) and activity per unit volume, percent activity of injected dosage, and ligand concentration were calculated. Plasma data was corrected for counting efficiency, calibration factors and measured metabolites of injected ligand. Percent activity of injected dose and ligand concentration was calculated. Plasma data was used as an input function in kinetic modeling. The binding potential values of [¹⁸F]3 in the ROI were determined by PMOD3.2.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

The authors would like to thank the technical staff of the PET/MRI and radiochemistry facilities at the Athinoula A. Martinos Center for Biomedical Imaging at Massachusetts General Hospital for the operation of cyclotron and synthesis modules. We would like to thank Drs. Tanabe and Nakanishi's laboratory (Osaka Bioscience Institute, Osaka, Japan) for providing the vector of mGlu₄ from the rat as a gift. The authors would like to acknowledge the following supporting instrument grants: 1S10RR029495-01, 1S10RR026666-01, and 1S10RR023452-01. Financial support for PP from The Orion Farnos Research Foundation and Kuopio University Foundation is gratefully

acknowledged. Finally, We also appreciate the generous help from the NIMH PDSP program (Contract # HHSN-271-2008-00025-C) led by Dr. Bryan L. Roth MD, PhD at the University of North Carolina at Chapel Hill and Project Officer Jamie Driscoll at NIMH, Bethesda MD, USA.

Funding Sources

NIBIB-R01EB012864 and NIMH-R01MH91684 to A-L Brownell

References

1. Conn PJ. Physiological roles and therapeutic potential of metabotropic glutamate receptors. *Ann N Y Acad Sci.* 2003; 1003:12–21. [PubMed: 14684432]
2. Watkins JC, Jane DE. The glutamate story. *Br J Pharmacol.* 2006; 147(S1):S100–S108. [PubMed: 16402093]
3. Hampson, DR.; Rose, EM.; Antflick, JE. The Structure of Metabotropic Glutamate Receptors. In: Gereau, RW.; Swanson, GT., editors. *The Glutamate Receptors.* Human Press; Totowa: 2008. p. 363–386.
4. Conn PJ, Pin JP. Pharmacology and functions of metabotropic glutamate receptors. *Ann Rev Pharmacol Toxicol.* 1997; 37:205–237. [PubMed: 9131252]
5. Riedel G, Platt B, Micheau J. Glutamate receptor function in learning and memory. *Behav Brain Res.* 2003; 140:1–47. [PubMed: 12644276]
6. Robichaud AJ, Engers DW, Lindsley CW, Hopkins CR. Recent progress on the identification of metabotropic glutamate 4 receptor ligands and their potential utility as CNS therapeutics. *ACS Chem Neurosci.* 2011; 17:433–449. [PubMed: 22860170]
7. Amalric M, Lopez S, Goudet C, Fisone G, Battaglia G, Nicoletti F, Pin JP, Acher FC. Group III and subtype 4 metabotropic glutamate receptor agonists: Discovery and pathophysiological applications in Parkinson's disease. *Neuropharmacology.* 2013; 66:53–64. [PubMed: 22664304]
8. Schapira AHV, Bezard E, Brotchie J, Calon F, Collingridge GL, Ferger B, Hengerer B, Hirsch E, Jenner P, Le Novere N, Obeso JA, Schwarzschild MA, Spampinato U, Davidai G. Novel pharmacological targets for the treatment of Parkinson's disease. *Nat Rev Drug Discov.* 2006; 5:845–854. [PubMed: 17016425]
9. Fetzagutti F, Balani-Guerra B, Corsi M, Nakanishi S, Corti C. Activation of the extracellular signal regulated kinase 2 by metabotropic glutamate receptors. *Eur J Neurosci.* 1999; 11:2073–2082. [PubMed: 10336676]
10. Niswender CM, Conn PJ. Metabotropic glutamate receptors: physiology, pharmacology, and disease. *Ann Rev Pharmacol Toxicol.* 2010; 50:295–322. [PubMed: 20055706]
11. Corti C, Aldegheri L, Somogyi P, Ferraguti F. Distribution and synaptic localization of the metabotropic glutamate receptor 4 (mGluR4) in the rodent CNS. *Neurosci.* 2002; 110:403–420.
12. Kinoshita A, Ohishi H, Nomura S, Shigemoto R, Nakanishi S, Mizuno N. Presynaptic localization of a metabotropic glutamate receptor, mGluR4a, in the cerebellar cortex: a light and electron microscope study in the rat. *Neurosci Lett.* 1996; 207:199–202. [PubMed: 8728484]
13. Bradley SR, Standaert DG, Rhodes KJ, Rees HD, Testa CM, Levey AI, Conn PJ. Immunohistochemical localization of subtype 4a metabotropic glutamate receptors in the rat and mouse basal ganglia. *J Comp Neurol.* 1999; 407:33–46. [PubMed: 10213186]
14. Valenti O, Marino MJ, Wittmann M, Lis E, DiLella AG, Kinney GG, Conn PJ. Group III metabotropic glutamate receptor-mediated modulation of the striatopallidal synapse. *J Neurosci.* 2003; 23:7218–7226. [PubMed: 12904482]
15. Valenti O, Mannaioni G, Seabrook GR, Conn PJ, Marino MJ. Group III metabotropic glutamate-receptor-mediated modulation of excitatory transmission in rodent substantia nigra pars compacta dopamine neurons. *J Pharmacol Exp Ther.* 2005; 313:1296–1304.
16. Engers DW, Niswender CM, Weaver CD, Jadhav S, Menon UN, Zamorano R, Conn PJ, Lindsley CW, Hopkins CR. Synthesis and evaluation of a series of heterobiaryl amides that are centrally penetrant metabotropic glutamate receptor 4 (mGluR4) positive allosteric modulators (PAMs). *J Med Chem.* 2009; 52:4115–4118. [PubMed: 19469556]

17. Kil K-E, Zhang Z, Jokivarsi K, Gong C, Choi J-K, Kura S, Brownell A-L. Radiosynthesis of N-(4-chloro-3-[¹¹C]methoxyphenyl)-2-picolinamide ([¹¹C]ML128) as a PET radiotracer for metabotropic glutamate receptor subtype 4 (mGlu₄). *Bioorg Med Chem*. 2013; 21:5955–5962. [PubMed: 23978356]
18. Patel S, Hamill TG, Connolly B, Jagoda E, Li W, Gibson RE. Species differences in mGluR5 binding sites in mammalian central nervous system determined using in vitro binding with [¹⁸F]F-PEB. *Nucl Med Biol*. 2007; 34:1009–1017. [PubMed: 17998106]
19. Wang J, Tueckmantel W, Zhu A, Pellegrino D, Brownell A-L. Synthesis and preliminary biological evaluation of 3-[¹⁸F]fluoro-5-(2-pyridinylethynyl)benzotrile as a PET radiotracer for imaging metabotropic glutamate receptor subtype 5. *Synapse*. 2007; 61:951–961. [PubMed: 17787003]
20. McCauley JA, Hess JW, Liverton NJ, McIntyre CJ, Romano JJ, Rudd MT. Phthalimide derivative metabotropic glutamate R4 ligands. Merck Research Lab. 2010 WO2010033349A1.
21. Jones CK, Engers DW, Thompson AD, Field JR, Blobaum AL, Lindsley SR, Zhou Y, Gogliotti RD, Jadhav S, Zamorano R, Bogenpohl J, Smith Y, Morrison R, Daniels JS, Weaver CD, Conn PJ, Lindsley CW, Niswender CM, Hopkins CR. Discovery, synthesis, and structure-activity relationship development of a series of N-4-(2,5-dioxopyrrolidin-1-yl)phenylpicolinamides (VU0400195, ML182): Characterization of a novel positive allosteric modulator of the metabotropic glutamate receptor 4 (mGlu₄) with oral efficacy in an antiparkinsonian animal model. *J Med Chem*. 2011; 54:7639–7647. [PubMed: 21966889]
22. Shi Q, Savage JE, Hufeisen SJ, Rauser L, Grajkowska E, Ernsberger P, Wroblewski JT, Nadeau JH, Roth BL. L-Homocysteine sulfinic acid and other acidic homocysteine derivatives are potent and selective metabotropic glutamate receptor agonists. *J Pharmacol Exp Ther*. 2003; 305:131–142. [PubMed: 12649361]
23. Roth, BL. [October 20, 2014] National Institute of Mental Health Psychoactive Drug Screening Program Assay Protocol Book. <http://pdsp.med.unc.edu/PDSP%20Protocols%20II%202013-03-28.pdf>
24. Ernst ML, Schmir GL. Isoimides. A kinetic study of the reaction of nucleophiles with N-phenylphthalisoimides. *J Am Chem Soc*. 1966; 88:5001–5009.
25. Cosford NDP, Tehrani L, Roppe J, Schweiger E, Smith ND, Andersen J, Bristow L, Brodtkin J, Jiang X, McDonald I, Rao S, Washburn M, Varney MA. 3-[(2-Methyl-1,3-thiazol-4-yl)ethynyl]pyridine: A potent and highly selective metabotropic glutamate subtype 5 receptor antagonist with anxiolytic activity. *J Med Chem*. 2003; 46:204–206. [PubMed: 12519057]
26. Lea PM IV, Faden AI. Metabotropic glutamate receptor subtype 5 antagonists MPEP and MTEP. *CNS Drug Rev*. 2006; 12:149–166. [PubMed: 16958988]
27. Tanabe Y, Ishii T, Shigemoto R, Nakanishi S. A family of metabotropic glutamate receptors. *Neuron*. 1992; 8:169–179. [PubMed: 1309649]

ABBREVIATIONS USED

mGlu₄	metabotropic glutamate receptor subtype 4
CHO	Chinese hamster ovary
RCY	radiochemical yield
EOS	end of synthesis
K222	kryptofix222
TAC	time-activity curve
ROI	region of interest
MTEP	3-((2-methyl-1,3-thiazole-4-yl)ethynyl)pyridine

QC quality control

FOV field of view




Figure 1.
 The representative mGlu₄ PAMs from *N*-phenyl-2-picolinamides and the corresponding phthalimide derivatives




Figure 2.
The binding curves from competitive binding assay (mGlu₄)




Figure 3.

(a) TACs derived from the rat brain PET images $[^{18}\text{F}]\mathbf{3}$. (n = 7) (b) mGlu₄ blocking studies demonstrated reduced uptake in the brain regions by 10-20%, while mGlu₅ blocking agents displayed minimal blocking effect indicating good selectivity to mGlu₄. Tissue uptake was calculated as area under TAC expressed as % dose/cc × min. Str = striatum, Hipp = hippocampus, Thal = thalamus, Cort = cortex, Cereb = cerebellum, WB = whole brain. Baseline studies (n = 7), mGlu₄ (2) blocking studies (n = 4), mGlu₅ (MTEP) blocking studies (n = 3) (c) Percentage of $[^{18}\text{F}]\mathbf{3}$ in the plasma over time. (n = 2)




Figure 4.

(a) Color coded PET images show accumulation of [^{18}F]**3** in the selected 5 coronal slices of the monkey brain at the time point 5-15 min after the injection of radioactivity. These images show the highest accumulation of [^{18}F]**3** in the caudate, putamen, thalamus and several cortical areas. (b) Fused PET and MR images from (a) and (c). (c) T1-weighted coronal MR images show the anatomical structures of selected brain areas. (n = 1) 189 MBq of [^{18}F]**3** (37 ng/kg) was injected.







Figure 5. TACs derived from the different areas of the monkey brain show fast accumulation of $[^{18}\text{F}]\mathbf{3}$ and reversible binding.



Scheme 1.
Syntheses of the phthalimide derivatives



Scheme 2.
The radiosynthesis of [³H]**1**



Scheme 3.
Radiosynthesis of [^{18}F]3

Table 1The results from *in vitro* competitive binding assay

Compound	mGlu ₄ IC ₅₀ (nM)	log(IC ₅₀) ± SD	RBI % ± SD (10 μM)
1	12.8	-7.89 ± 0.09	100.1 ± 2.0
2	9.9	-8.00 ± 0.09	99.1 ± 0.5
3	5.1	-8.29 ± 0.08	99.4 ± 0.7
4	4.2	-8.38 ± 0.07	99.4 ± 0.9
7	11.8	-7.93 ± 0.09	96.6 ± 1.2
8	26.3	-7.53 ± 0.09	96.0 ± 3.0

-RBI = Relative binding inhibition

-SD = Standard deviation