

Published in final edited form as:

Chem Sci. 2015 February ; 6(2): 1115–1119. doi:10.1039/C4SC03106G.

Cobalt co-catalysis for cross-electrophile coupling: diarylmethanes from benzyl mesylates and aryl halides†

 Laura K. G. Ackerman^{#a}, Lukiana L. Anka-Lufford^{#a}, Marina Naodovic^{#a}, and Daniel J. Weix^a
^a Department of Chemistry, University of Rochester, Rochester, NY USA 14627-0216

[#] These authors contributed equally to this work.

Abstract

The nickel-catalyzed cross-coupling of aryl halides with alkyl radicals derived from alkyl halides has recently been extended to couplings with carbon radicals generated by a co-catalyst. In this study, a new co-catalyst, cobalt phthalocyanine (Co(Pc)), is introduced and demonstrated to be effective for coupling substrates not prone to homolysis. This is because Co(Pc) reacts with electrophiles by an S_N2 mechanism instead of by the electron-transfer or halogen abstraction mechanisms previously explored. Studies demonstrating the orthogonal reactivity of (bpy)Ni and Co(Pc), applying this selectivity to the coupling of benzyl mesylates with aryl halides, and the adaptation of these conditions to the less reactive benzyl phosphate ester and an enantioconvergent reaction are presented.

Introduction

Cross-coupling relies upon the selective, ordered activation of two different substrates. For the coupling of nucleophiles with electrophiles, a single catalyst reacts with the electrophile by oxidative addition and the nucleophile by transmetalation, resulting in high cross-selectivities. Cross-electrophile coupling,¹ the union of two different electrophiles, achieves selectivity by different mechanisms. Specifically, we have recently shown that in nickel-catalyzed reactions electrophiles can be differentiated by heterolysis and homolysis (Figure 1, entry 1).² Selectivity arises because (L)Ni⁰ reacts with aryl halides faster than alkyl halides, but (L)Ni^I forms alkyl radicals faster than aryl radicals.³

The key to successful cross-electrophile coupling is selective radical generation from R-Y (Figure 1).⁴ In order to expand the types of substrates that can be coupled with aryl halides by this electrophile + radical mechanism, alternative methods of generating radicals must be developed. A key advance was that radical generation and coupling can be accomplished by two different catalysts (Figure 1). Our group⁵ and the groups of Sanford,⁶ Molander,⁷ and

†Electronic Supplementary Information (ESI) available: [details of any supplementary information available should be included here]. See DOI: 10.1039/b000000x/

MacMillan and Doyle^{6b} have independently shown that a variety of co-catalysts can allow coupling of otherwise unreactive substrates under mild conditions (Figure 1, entries 2-4).

All of the methods reported to date convert the substrate into a radical by single-electron oxidation or reduction (Figure 1). As such, substrates must be easily oxidized or reduced. The development of co-catalysts that form radicals by different mechanisms would enable further expansion of substrate scope in this arylation strategy.

We report here that cobalt phthalocyanine (Co(Pc)) is an excellent, non-photochemical co-catalyst for radical generation that is compatible with nickel catalysis (Figure 1, entry 5).⁸ Co(Pc) differs from previously described co-catalysts because it generates radicals after 2-electron nucleophilic substitution⁹ rather than single-electron transfer. This gives Co(Pc) different selectivity than previous approaches. For example, alkylsulfonate and alkylphosphate esters are unreactive towards single-electron transfer due to the strength of the C-O bond, but react rapidly by nucleophilic substitution.

Results and Discussion

To demonstrate the potential of Co(Pc), we applied this co-catalyst to the synthesis of diarylmethanes from two abundant electrophiles, a benzyl alcohol derivative and an aryl halide. Although a variety of approaches to diarylmethanes have been developed,¹⁰ their prominence in medicinal chemistry¹¹ has justified continued attention. The majority of recent approaches involve the cross-coupling of benzylmetal reagents^{7, 12} or arylmetal reagents,¹³ but the need to pre-form each organometallic reagent can be limiting. A major advance was the development of methods where organozinc reagents were generated and coupled concurrently, but 2-4 equiv of the benzyl halide was still required for high yields and no examples with more abundant benzyl alcohol derivatives¹⁴ were reported.^{12d-f, 13e} Gosmini reported one example of the coupling of benzyl chloride with ethyl 4-bromobenzoate under conditions that might not involve an organozinc intermediate,¹⁵ but the scope of this method has not been explored. Finally, Reisman reported the coupling of secondary benzylic chlorides with vinyl bromides, but the use of aryl halides or benzyl alcohols was not reported.¹⁶ Compared to these known methods, our new approach avoids pre-formed nucleophiles, starts from benzyl alcohols instead of the less abundant benzyl halides, and does not use a large excess of one coupling partner.

The application of our reported conditions¹ to the coupling of aryl halides with benzyl bromide resulted primarily in the formation of bibenzyl (Scheme 1A). This is due to the fact that benzyl bromide reacts with (L)Ni faster than aryl halides (Scheme 1B). For example, benzyl bromide is converted to bibenzyl and toluene in only 60 min (Supporting Information Table S1).

In order to restore selectivity, we sought to take advantage of the low reactivity of bipyridine-ligated nickel catalysts with alkyl sulfonate esters in order to prevent formation of benzylnickel and favor formation of arylnickel (Scheme 1C). Benzyl mesylates can be conveniently generated *in situ* from abundant benzyl alcohols. Although this approach prevented bibenzyl formation (Table S1), it did not improve the yield of diarylmethane because nickel is slow to form a benzyl radical from a benzyl mesylate (Table 1, entry 1).

The addition of Co(Pc) as a co-catalyst could generate the missing benzyl radical (Scheme 1C), separating the oxidative addition and radical generation steps.¹⁷

Significantly, this optimized nickel and cobalt system allows for the coupling of benzyl mesylates with aryl bromides, which occurs in low yield in the absence of Co(Pc) (Table 1, entries 1-3).²⁰ Furthermore, it appears that both (dtbbpy)NiBr₂ and Co(Pc) are required for high selectivity as neither catalyst is effective without a ligand (entries 4 and 7). An alternative co-catalyst, sodium iodide,^{13j, 21} was not as effective as Co(Pc) (entry 5 and Table S1). The different products formed arise from different mechanisms of co-catalysis: Co(Pc) selectively forms benzyl radicals from BnOMs while NaI converts BnOMs to BnI which reacts similarly to benzyl bromide (Scheme 1B). For convenience, the aryl halide, ligand, Co(Pc), and zinc could be added with the alcohol, Ms₂O, and EtN(*i*-Pr)₂ or after mesylate formation was complete with little difference in yield. However, if nickel was added at the beginning, rapid reduction or dimerization of the starting materials resulted.²⁰

The nickel and cobalt co-catalytic strategy of synthesizing diarylmethanes was also successful when aryl iodides were employed as coupling partners. When benzyl mesylate was reacted with iodobenzene under the optimized reaction conditions, the diphenylmethane product was formed in good yield. Although product was formed in the absence of Co(Pc), the co-catalytic method improved yield and selectivity (Table 1, entries 9-12). Added NaI depressed the yield only slightly (entry 13 vs. entry 9), suggesting that PhI competes with *in situ* formed BnI more effectively than PhBr for oxidative addition to nickel (Scheme 1B).

Application of these conditions to a variety of different aryl bromides, aryl iodides, and benzyl alcohols demonstrated the generality of the method (Scheme 2). The reaction tolerates functional groups, such as aldehyde (**3e**) and methyl ketone (**3d**), which are susceptible to the nucleophilic and basic character of some organometallic reagents. Additionally, a boronic acid ester was coupled without chemoselectivity problems, demonstrating complementarity with existing cross-coupling methods (**3f**).¹³

The coupling of aryl bromides and aryl iodides with benzyl mesylate provided comparable yields. In examples where the selectivity over bibenzyl was high but the yield was low, hydrodehalogenation of the arene was responsible for diminished yields.²⁰ The aryl bromide conditions could be applied without further optimization to a vinyl bromide with reasonable success (**3j**). Both electron-rich and electron-poor benzyl alcohols couple effectively and steric hindrance on the benzyl alcohol did not result in lower yields (**3l**, **3c'**, **3m**, **3g'**, **3n**).²² Finally, we were able to apply this approach to the synthesis of beclobrate (**3o**), a diarylmethane compound that can be used to alter lipid levels in humans.²³

The reactivity and selectivity of the reaction can also be rationally optimized. For example, the coupling of benzyl diethyl phosphate ester²⁴ was sluggish under our standard conditions and starting materials were not consumed (Table 2). Presumably, this was due to the reduced leaving group ability of OP(O)(OEt)₂ compared to OMs.²⁵ Increasing the amount of Co(Pc) from 1 mol% to 6 mol% as well as increasing the temperature from rt to 80 °C allowed the reaction to proceed in high yield to form diphenylmethane (**3a'**).²² Notably, we observe no hydrodehalogenation under these conditions, perhaps because there is no acid (EtN(*i*-

Pr)₂•HOMs) present. We anticipate that a similar strategy could be employed to couple other less reactive electrophiles.

The likely intermediacy of a benzyl radical suggested to us that an enantioconvergent coupling of racemic secondary benzylic electrophile with an aryl bromide could be achieved.^{26, 27} Although yields were low with secondary benzylic alcohol derivatives, α -chloroethylbenzene (**7**) coupled efficiently under slightly modified standard conditions to form diarylethane **8**²⁸ in 41% isolated yield and 43% *ee* (Scheme 3).²⁹ This yield and enantioselectivity compares favorably to the results of Molander in the coupling of a racemic secondary benzylboron reagent with an aryl halide using the same chiral ligand (52% yield, 50% *ee*), suggesting a similar enantiodetermining step (radical capture by a chiral nickel complex).⁷ Together with the related highly enantioselective coupling of vinyl halides by Reisman,¹⁶ these results suggest highly enantioselective reactions to form diarylmethanes will soon be realized. Future studies will be aimed at improving the enantioselectivity and yield of these and related reactions.

Conclusions

In conclusion, Co(Pc) is a new co-catalyst for radical generation that is compatible with nickel-catalysis offering a different selectivity than other established co-catalysts while maintaining high functional-group compatibility. The first synthesis of diarylmethanes from benzyl mesylates and aryl iodides or bromides is possible because 1) [Ni⁰] reacts with Ar-X faster than with BnOMs, preventing bibenzyl formation and 2) Co(Pc) reacts with benzyl mesylate faster than with Ar-X, selectively forming benzyl radicals. We propose that these benzyl radicals react with (L)Ni^{II}(Ar)X to form product.^{2, 4, 6, 7} These results demonstrate the potential of Co(Pc) as a nucleophilic co-catalyst for radical cross-coupling.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgements

This work was supported by the NIH (R01 GM097243). LKGA is supported by an NSF Graduate Fellowship (NSF DGE-1419118). DJW is an Alfred P. Sloan Research Fellow. We thank Daniel A. Everson and Maxwell B. Hecht (Univ. of Rochester) for initial experiments with benzylic halides and mesylates; Sylvia Chen (Univ. of Rochester) for key experimental support and for assisting in the analysis of results; and Zulema Melchor (Univ. of Rochester) for experimental assistance.

References

1. a Everson DA, Shrestha R, Weix DJ. *J. Am. Chem. Soc.* 2010; 132:920. [PubMed: 20047282] b Everson DA, Jones BA, Weix DJ. *J. Am. Chem. Soc.* 2012; 134:6146. [PubMed: 22463689] c Everson DA, Weix DJ. *J. Org. Chem.* 2014; 79:4793. [PubMed: 24820397] d Knappke CEI, Grupe S, Gärtner D, Corpet M, Gosmini C, Jacobi von Wangelin A. *Chem. Eur. J.* 2014; 20:6828. [PubMed: 24825799]
2. Biswas S, Weix DJ. *J. Am. Chem. Soc.* 2013; 135:16192. [PubMed: 23952217]
3. For the generation of alkyl radicals from the reaction of nickel(I) with alkyl halides, see: Laskowski CA, Bungum DJ, Baldwin SM, Del Ciello SA, Iluc VM, Hillhouse GL. *J. Am. Chem. Soc.* 2013; 135:18272. [PubMed: 24237257]

4. For the reaction of an arylnickel(II) with an alkyl radical to form cross-coupled product, see: Breitenfeld J, Ruiz J, Wodrich MD, Hu X. *J. Am. Chem. Soc.* 2013; 135:12004. [PubMed: 23865460]
5. Zhao Y, Weix DJ. *J. Am. Chem. Soc.* 2014; 136:48. [PubMed: 24341892]
6. a Kalyani D, McMurtrey KB, Neufeldt SR, Sanford MS. *J. Am. Chem. Soc.* 2011; 133:18566. [PubMed: 22047138] b Zuo Z, Ahneman D, Chu L, Terrett J, Doyle AG, MacMillan DWC. *Science*. 2014; 345:437. [PubMed: 24903563]
7. Tellis JC, Primer DN, Molander GA. *Science*. 2014; 345:433. [PubMed: 24903560]
8. For the use of Co(Pc) to activate alkyl halides for Cr-mediated coupling with aldehydes, see references 9e and 9f.
9. a Alonso F, Beletskaya IP, Yus M. *Chem. Rev.* 2002; 102:4009. [PubMed: 12428984] b Espenson J, Shveima J. *J. Am. Chem. Soc.* 1973; 95:4468. c Espenson J, Sellers T. *J. Am. Chem. Soc.* 1974; 96:94. d Torii S, Inokuchi T, Yukawa T. *J. Org. Chem.* 1985; 50:5875. e Takai K, Nitta K, Fujimura O, Utimoto K. *J. Org. Chem.* 1989; 54:4732. f Kurosu M, Lin M-H, Kishi Y. *J. Am. Chem. Soc.* 2004; 126:12248. [PubMed: 15453741]
10. a Liegault B, Renaud J-L, Bruneau C. *Chem. Soc. Rev.* 2008; 37:290. Reviews. [PubMed: 18197345] b Kuwano R. *Synthesis*. 2009:1049.
11. A Scifinder Scholar database search for ArCH₂Ar (<https://scifinder.cas.org>, June 2014) noted 567,220 diarylmethanes that had reported biological studies and 324,596 patents. Activities included antitumor, CNS, cardiovascular, anti-infective, anti-inflammatory, antidiabetic, and antiobesity activity.
12. a Itami K, Mineno M, Kamei T, Yoshida J.-i. *Org. Lett.* 2002; 4:3635. [PubMed: 12375906] b Flaherty A, Trunkfield A, Barton W. *Org. Lett.* 2005; 7:4975. [PubMed: 16235936] c Schade MA, Metzger A, Hug S, Knochel P. *Chem. Commun.* 2008:3046. d Duplais C, Krasovskiy A, Wattenberg A, Lipshutz BH. *Chem. Commun.* 2010; 46:562. e Krasovskaya V, Krasovskiy A, Lipshutz BH. *Chem. Asian J.* 2011; 6:1974. [PubMed: 21626709] f Krasovskaya V, Krasovskiy A, Bhattacharjya A, Lipshutz BH. *Chem. Commun.* 2011; 47:5717.
13. a Kuwano R, Yokogi M. *Org. Lett.* 2005; 7:945. [PubMed: 15727481] b Singh R, Viciu MS, Kramareva N, Navarro O, Nolan SP. *Org. Lett.* 2005; 7:1829. [PubMed: 15844917] c Kofink CC, Knochel P. *Org. Lett.* 2006; 8:4121. [PubMed: 16928089] d Molander GA, Elia MD. *J. Org. Chem.* 2006; 71:9198. [PubMed: 17109547] e Amatore M, Gosmini C. *Chem. Commun.* 2008:5019. f McLaughlin M. *Org. Lett.* 2005; 7:4875. [PubMed: 16235911] g Bedford RB, Huwe M, Wilkinson MC. *Chem. Commun.* 2009:600. h Bedford RB, Gower NJ, Haddow MF, Harvey JN, Nunn J, Okopie RA, Sankey RF. *Angew. Chem., Int. Ed.* 2012; 51:5435. i Rao MLN, Dhanorkar RJ. *RSC Advances*. 2013; 3:6794. j Do H-Q, Chandrashekar ERR, Fu GC. *J. Am. Chem. Soc.* 2013; 135:16288. [PubMed: 24164502] k Maity P, Shacklady-McAtee DM, Yap GPA, Sirianni ER, Watson MP. *J. Am. Chem. Soc.* 2013; 135:280. [PubMed: 23268734] l Zhao F, Tan Q, Xiao F, Zhang S, Deng G-J. *Org. Lett.* 2013; 15:1520. [PubMed: 23556515] m Harris MR, Hanna LE, Greene MA, Moore CE, Jarvo ER. *J. Am. Chem. Soc.* 2013; 135:3303. [PubMed: 23414579] n Zhou Q, Srinivas HD, Dasgupta S, Watson MP. *J. Am. Chem. Soc.* 2013; 135:3307. [PubMed: 23425080] o Wisniewska HM, Swift EC, Jarvo ER. *J. Am. Chem. Soc.* 2013; 135:9083. [PubMed: 23751004] p Shacklady-McAtee DM, Roberts KM, Basch CH, Song Y-G, Watson MP. *Tetrahedron*. 2014; 70:4257. [PubMed: 25364060]
14. Commercial availability: 6,152 ArCH₂OH vs. 2,357 ArCH₂NH₂ vs. 605 ArCH₂Cl vs. 948 ArCH₂Br; data from eMolecules database (eMolecules, Inc.), accessed via REAXYS (Elsevier, Inc.) on April 21, 2014.
15. For one example of coupling benzyl chloride (2 equiv) with an aryl bromide that might not proceed through an organometallic intermediate, see: Amatore M, Gosmini C. *Chem. Eur. J.* 2010; 16:5848. [PubMed: 20379979]
16. Cherney, AH.; Reisman, SE. [Oct 5, 2014] *J. Am. Chem. Soc.* [Online early access]. DOI: 10.1021/ja508067c Published online Sept 22, 2014. <http://dx.doi.org/10.1021/ja508067c>
17. Co(Pc) reacted with benzyl mesylate to form benzyl radicals, quantitated as toluene (5, 4%) and bibenzyl (42%). See Table S1 in the Supporting Information.
18. Conditions: 1:1 PhI:BnBr, 7 mol% NiBr₂•3H₂O, 5 mol% dtbbpy, and 2 equiv Zn dust in DMA (0.25 M) at 60 °C for 10 min. BnBr is consumed, most of the iodobenzene remains.

19. Ligand-free nickel does not provide high yields (Table 2, entry 7) and excess nickel does not negatively impact reactions.^{1a}
20. See Supporting Information, Scheme S1 and Chart S1-S10 for further selectivity and optimization details.
21. a Tipson RS, Clapp MA, Cretcher LH. *J. Org. Chem.* 1947; 12:133.b Ito S, Fujiwara Y.-i, Nakamura E, Nakamura M. *Org. Lett.* 2009; 11:4306. [PubMed: 19731937] c Prinsell MR, Everson DA, Weix DJ. *Chem. Commun.* 2010; 46:5743.
22. We denote the same product made by a different set of substrates with a prime. For example, **3c** and **3c'** are made from a different disconnection. **3a** and **3a'** are made from different benzyl alcohol derivatives.
23. a Thiele K, Ahmed Q, Jahn U, Adrian RW. *Arzneim.-Forsch.* 1979; 29:711. [PubMed: 582967] b Mertins K, Iovel I, Kischel J, Zapf A, Beller M. *Adv. Synth. Catal.* 2006; 348:691.
24. See references 13c and 13f for examples of coupling benzyl phosphate esters with arylmetal reagents.
25. The relative leaving group ability can be estimated from the pK_a data available in the literature for MsOH (1.6 in DMSO) and (PhO)₂P(O)OH (3.9 in DMSO). See: Bordwell FG, Algrim D. *J. Org. Chem.* 1976; 41:2507. Christ P, Lindsay AG, Vormittag SS, Neudörfl J-M, Berkessel A, O'Donoghue AC. *Chem. Eur. J.* 2011; 17:8524. [PubMed: 21714022]
26. For examples of stereoselective or stereospecific coupling of benzylic substrates, see references 13j, 13k, 13m, 13n, 13o, 13p, and the references in the following endnote.
27. a Taylor BLH, Swift EC, Waetzig JD, Jarvo ER. *J. Am. Chem. Soc.* 2010; 133:389. [PubMed: 21155567] b Swift EC, Jarvo ER. *Tetrahedron.* 2013; 69:5799. [PubMed: 23956470]
28. Diarylethane **8** has been previously synthesized by the enantiospecific coupling of secondary benzylsilicon and benzylboron reagents with aryl iodides or aryl triflates. Yields over the two steps were 41-51% with a 34% - 83% *ee*. See: Hatanaka Y, Hiyama T. *J. Am. Chem. Soc.* 1990; 112:7793. Imao D, Glasspoole B, Laberge V. r. Crudden C. *J. Am. Chem. Soc.* 2009; 131:5024. [PubMed: 19301820] Glasspoole BW, Oderinde MS, Moore BD, Antoft-Finch A, Crudden CM. *Synthesis.* 2013; 45:1759.
29. Further study is needed to better understand reactions of secondary benzylic chlorides. For these electrophiles, a co-catalyst may not be required. In preliminary studies, reactions run without added Co(Pc) provided similar results to reactions run with Co(Pc). Unlike reactions with vinyl bromides (ref. 16), reactions with bis(oxazoline) ligands provided low yields of racemic material.






Figure 1. Comparison of radical co-generation methods in cross-coupling. An electrophile (Ar-X) reacts to form an arylmetal intermediate and the other substrate (R-Y) reacts to form a radical (R•).



Scheme 1. Diarylmethanes from cross-electrophile coupling: problem (A and B) and mechanism-based solution (C).¹⁸ Ligand = 4,4'-di-*tert*-butyl-2,2'-bipyridine.

**Scheme 3.**

Preliminary enantioconvergent coupling of a secondary benzyl chloride with an aryl bromide. Configuration assigned by comparison of the optical rotation to those reported in the literature.

Table 1

Cross-electrophile coupling of Bn-OH with Ar-X.^a

Entry	X	Catalyst ^b	Yield 3 (A%) ^c	3:4
1	Br	[Ni] only	3	1:1
2	Br	Co(Pc) only	0	ND:3
3	Br	[Ni], Co(Pc)	73	1:ND
4	Br	[Ni], CoCl ₂	4	4:1
5	Br	[Ni], NaI (25 mol%)	25	1:1
6	Br	[Ni], Co(Pc), Mn instead of Zn	64	4:1
7	Br	NiBr ₂ •3H ₂ O, Co(Pc)	1	1:6
8	Br	No [Ni] and no Co(Pc)	0	0:1
9 ^{d,e}	I	[Ni] only	71	7:1
10 ^e	I	Co(Pc) only	0	ND:1
11 ^{d,e}	I	[Ni], Co(Pc)	83	17:1
12	I	[Ni], Co(Pc)	83	83:1
13	I	[Ni], NaI (25 mol%)	64	8:1
14	I	[Ni], Co(Pc), Mn instead of Zn	42	1.4:1

^aReactions run at 0.25 M in DMA. BnOMs was formed *in situ* from BnOH, Ms₂O (1.44 equiv), and EtN(*i*-Pr)₂ (1.6 equiv). See Supporting Information.

^b[Ni] = 7 mol% NiBr₂•3H₂O and 5 mol% dtbbpy.¹⁹ Co(Pc) = 1 mol% cobalt phthalocyanine.

^cYields and ratios reported as GC Area%.

^dReaction run at 60 °C.

^eReaction complete within 1 h.

Table 2Reaction with benzyl diethyl phosphate, a less reactive electrophile.^a

Entry	Co(Pc) (mol %)	T (°C)	Yield ^b
1	3	rt	ND ^c
2	3	40	ND ^c
3	1	80	3 ^{c,d}
4	3	80	36 ^{c,d}
5	6	80	(70) ^d

^aReactions run at 0.25 M in DMA using pre-formed **6** for 15-22 h. See Supporting Information.^bYield of **3a'** from GC area% data. Yields in parentheses are isolated yields of purified product.^cSignificant amounts of starting materials remain.^dOnly byproduct observed by GC analysis was small amounts of benzene from hydrodehalogenation of Ph-Br.