



Published in final edited form as:

Nat Commun. ; 6: 5984. doi:10.1038/ncomms6984.

Hydrophobic mismatch sorts SNARE proteins into distinct membrane domains

Dragomir Milovanovic^{1,†}, Alf Honigmann^{2,†}, Seiichi Koike¹, Fabian Göttfert², Gesa Pähler⁴, Meike Junius⁵, Stefan Müller⁵, Ulf Diederichsen⁵, Andreas Janshoff⁴, Helmut Grubmüller³, Herre Jelger Risselada^{3,6}, Christian Eggeling^{2,7}, Stefan W. Hell², Geert van den Bogaart^{1,8,*}, and Reinhard Jahn^{1,*}

¹Department of Neurobiology, Max Planck Institute for Biophysical Chemistry, Göttingen, Germany ²Department of NanoBiophotonics, Max Planck Institute for Biophysical Chemistry, Göttingen, Germany ³Department of Theoretical and Computational Biophysics, Max Planck Institute for Biophysical Chemistry, Göttingen, Germany ⁴Institute for Physical Chemistry, Georg-August-University Göttingen, Germany ⁵Institute for Organic and Biomolecular Chemistry, Georg-August-University Göttingen, Germany ⁶Department of Chemistry, Leibniz Institute of Surface Modification, Leipzig, Germany ⁷MRC Human Immunology Unit, Weatherall Institute of Molecular Medicine, University of Oxford, United Kingdom ⁸Department of Tumor Immunology, Radboud University Medical Centre, Nijmegen, The Netherlands

Abstract

The clustering of proteins and lipids in distinct microdomains is emerging as an important principle for the spatial patterning of biological membranes. Such domain formation can be the result of hydrophobic and ionic interactions with membrane lipids as well as of specific protein-protein interactions. Here, using plasma membrane-resident SNARE proteins as model, we show that hydrophobic mismatch between the length of transmembrane domains and the thickness of the lipid membrane suffices to induce clustering of proteins. Even when the transmembrane domains differ in length by only a single residue, hydrophobic mismatch can segregate structurally closely homologous membrane proteins in distinct membrane domains. Domain formation is further fine-tuned by interactions with polyanionic phosphoinositides and homo- and heterotypic protein interactions. Our findings demonstrate that hydrophobic mismatch contributes to the structural organization of membranes.

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

*Correspondence to: mail@bogeert.com and rjahn@gwdg.de.

[†]D.M. and A.H. contributed equally to this work.

Author contributions: D.M, A.H, G.v.d.B, and R.J. designed the research and wrote the paper. D.M. did all the reconstitution experiments and analyzed the data. D.M. and S.K. performed cellular analysis. H.J.R. and H.G. performed the molecular dynamics simulations; G.P. and A.J. performed imaging ellipsometry measurements; S.M, M.J, U.D. synthesized the peptide. A.H, F.G, C.E. and S.W.H. contributed to the microscopy.

The Authors declare no conflict of interest.

Introduction

Since many years, the architecture of biological membranes has been subject of intense research. Presently, the concepts for the arrangement of membrane proteins, particularly in the plasma membrane of eukaryotic cells, are undergoing major changes. According to the classical Singer-Nicholson model membrane proteins are viewed as separate entities floating in the lipid bilayer like icebergs in a sea. In contrast, it is now appreciated that many (if not all) membrane proteins are not evenly distributed across the membrane but rather organized in microdomains with a high local protein concentration, ranging in size between 2 and 200 nm¹⁻⁴. Frequently, domain formation appears to be essential for the functions governed by these proteins, e.g. by forming localized hotspots for signaling or for exo- and endocytosis⁵⁻⁷.

SNARE proteins operating in exocytosis at the plasma membrane, in particular syntaxins 1 and 4, have served as convenient models for studying mechanisms responsible for protein clustering⁸⁻¹⁰. Syntaxins are tail-anchored membrane proteins with a single hydrophobic transmembrane domain (TMD) at the C-terminus¹¹. In the plasma membrane these proteins are organized in clusters with a diameter of around 70 nm that are dependent on the presence of cholesterol in the membrane, but are distinct from clusters formed by “classical raft” residents such as caveolin or proteins linked to a glycosylphosphatidylinositol (GPI) anchor^{5,12,13}. In addition, clustering is dependent on the presence of plasma membrane-specific polyphosphoinositides (PI(4,5)P₂ and PI(3,4,5)P₃) that bind to a conserved juxtamembrane polybasic motif^{6,10,14-18}. However, the physical principles underlying cluster formation within the bilayer are still unclear, with explanations including mechanisms as diverse as phase partitioning into cholesterol-enriched membrane rafts^{5,19-21}, decreased solubility caused by the presence of cholesterol^{15,16}, electrostatic interactions with the phosphoinositides^{10,18}, and homophilic as well as heterophilic interactions between the proteins themselves involving either the hydrophobic TMDs or the hydrophilic domains^{8,9,22}. In any case it is becoming apparent that membrane cholesterol plays a fundamental role in domain formation that is fine-tuned by the other factors but cannot be replaced by them. The causative mechanism of cholesterol is still unclear, with neither phase-partitioning (yielding phase-segregated areas of much bigger size which exclude many of the cholesterol-dependent proteins^{4,23}) nor specific binding of cholesterol to individual proteins (shown for only few of the cholesterol-dependent proteins) providing a satisfying explanation. A second important point is that closely homologous SNARE proteins such as syntaxin 1 and 4 are segregated into non-overlapping membrane domains⁸. Understanding how syntaxins segregate into distinct membrane clusters is essential for understanding their distinct roles in regulated (syntaxin 1) and constitutive (syntaxin 4) exocytosis.

In this study, we have investigated whether hydrophobic mismatch may be the underlying physical principle for many of the effects summarized above, with the effects of cholesterol being due to effects on membrane thickness. Hydrophobic mismatch means that the length of the hydrophobic part of a membrane-spanning macromolecule does not match the thickness of the hydrophobic core of the membrane. Such mismatch results in hydrophobic “defects” at the boundary between protein and membrane lipid, which impose an energy

penalty that can be minimized by clustering (Fig. 1a). The plasma membrane is a complex milieu with diverse thicknesses, amendable to quick changes, and we reasoned that the patterning of the plasma membrane might be caused by sequestering of lipids and proteins in membrane regions of matching hydrophobic thickness. Indeed, mismatch-driven clustering was previously shown to occur in a system employing synthetic peptides and artificial membranes^{24,25}. Furthermore, hydrophobic mismatch was invoked as possible mechanism for the retention of membrane proteins in early compartments of the secretory pathway (such as the endoplasmic reticulum (ER)) since membrane thickness increases between the ER and the plasma membrane²⁶. However, there appear to be many exceptions^{27,28}, and thus hydrophobic mismatch is presently not being considered as a relevant factor contributing to membrane patterning and protein sorting. In this study, we show that hydrophobic mismatch between the length of transmembrane domains and the thickness of the plasma membrane contributes to the clustering of proteins in the plasma membrane. We also show that hydrophobic mismatch can contribute to the segregation of structurally closely homologous SNARE proteins in distinct membrane domains even when the transmembrane domains differ in length by only a single residue.

Results

Clustering of syntaxin isoforms by hydrophobic mismatch

In order to dissect the protein distribution in different membrane environments, we started by addressing syntaxin clustering in artificial liposomes with defined membrane thicknesses. First, we determined the effect of acyl-chain length and cholesterol on membrane thickness. To this end, we prepared ~100 nm sized large unilamellar vesicles (LUVs) composed of unsaturated phosphatidylcholine (PC) with stepwise increase in the length of the acyl-chains (ranging from C14:1 to C20:1) either in the absence or in the presence of 30 mol% cholesterol. The thickness of these membranes was determined by imaging ellipsometry measurements (Fig. 1b), which is based on polarization changes of monochromatic light upon reflection on the bilayer²⁹. As expected³⁰, the membrane thickness increased by about 0.15 nm for each carbon unit added to the phospholipid acyl chains. Inclusion of 30 mol% cholesterol increased the membrane thickness by approximately 0.8 nm, independently of the acyl chain length (Fig. 1b), indicating that cholesterol is a main modulator of membrane thickness.

We choose the SNAREs syntaxin 1 and syntaxin 4 to study the influence of hydrophobic mismatch on the distribution of membrane proteins for two reasons. First, the lengths of the hydrophobic segments of these SNAREs (21–23 AA) appear to be shorter than that needed to fully span the average hydrophobic core of a plasma membrane. This is particularly evident from the crystal structure of the neuronal SNARE complex where the TMD segments of the SNAREs synaptobrevin 2 and syntaxin 1 seem indeed not sufficiently long to traverse the average thickness of plasma membrane of eukaryotic cells (~4 nm). In a simulation, this resulted in defects in lipid packing^{30,31} and suggested that these SNAREs might prefer, or even organize, membrane regions of lipids with matching thicknesses. Second, these two syntaxins segregate in separate clusters although they are homologous and structurally very similar to each other. While it was shown previously that segregation

depends at least in part on homophilic interactions between the cytoplasmic domains²², it is conceivable that the small differences in the length of the TMD segments (see below) may contribute to segregation. To isolate the effects on clustering within the membrane space from “secondary” effects caused by protein-protein interactions in the hydrophilic space, we employed truncation mutants of syntaxins 1 and 4 with their cytoplasmically oriented domains deleted.

We tested whether syntaxin clustering is dependent on membrane thickness. For this purpose we measured clustering of the syntaxin 1 truncation mutant (sx-1TM) by Förster resonance energy transfer (FRET) using an approach very similar to that described by Murray and Tamm^{15,16}. Two sx-1TM populations labeled with spectrally separated fluorophores were mixed and incorporated into the liposomes described above, resulting in a high FRET signal in case of cluster formation (Fig. 1c). We found that in the absence of cholesterol, the FRET efficiency of sx-1TM was lowest (i.e. least protein clustering) in membranes composed of C16:1 PC compared to C14:1, C18:1 and C20:1 PC (Fig. 1d). In the presence of cholesterol, clustering of syntaxin 1 was strongly enhanced (about 50% increase in FRET efficiency, Fig. 1d) for a given acyl chain length, but was similar when related to membranes with the same thickness without cholesterol. As an independent approach we also determined the lateral mobility of sx-1TM by fluorescence correlation spectroscopy (FCS) in stacked supported lipid bilayers. Lateral mobility is expected to inversely correlate with cluster formation. As shown in figure 1d, a profile very similar to the FRET measurements was obtained, with sx-1TM showing the highest mobility in membranes of C16:1 PC. Together, our data show that syntaxin 1 clustering was lowest in membranes composed of C16:1 PC, corresponding to a hydrophobic thickness of about 3.4 nm, well below the average thickness of a eukaryotic plasma membrane. Clustering increased with increasing membrane thickness independent of whether this increase was caused by longer acyl chains or by the inclusion of cholesterol in the membrane.

To directly visualize syntaxin clustering in dependence of cholesterol, we prepared supported artificial bilayers (C18:1 PC, with or without 30 mol% cholesterol) containing sx-1TM. In this experiment sx-1TM was labeled with the dye Atto647N to monitor its distribution, with the membrane being stained with the green fluorescent lipid analogue DiO (3,3'-dilinoleoyloxacarbocyanine) (Fig. 1e). Clustering was clearly observable in the cholesterol-containing membranes whereas it was much less conspicuous in the absence of cholesterol, in agreement with the results described above. Clearly, clustering is not due to cholesterol-induced phase separation of membrane lipids since these membranes did not contain any lipids with saturated fatty acids required for partitioning into Lo and Ld phases. Rather, our experiments reveal that the effect of cholesterol is caused by the increase of the membrane thickness, which results in clustering due to increased hydrophobic mismatch.

Interestingly, the TMD of syntaxin 4 is longer by 1–2 amino acids than that of syntaxin 1, and this difference in length is well conserved in mammalian species (Fig. 1f). Since the length of the TMD determines the hydrophobic matching with the surrounding lipid environment^{24,25}, it is expected that optimal matching (i.e. lowest clustering) requires a thicker bilayer for syntaxin 4 than syntaxin 1. Therefore, we compared clustering of syntaxin 1 with that of syntaxin 4 as a function of membrane thickness, using an analogous truncation

mutant for syntaxin 4 that lacked most of the cytoplasmic part (sx-4TM). Indeed, the local minimum of sx-4TM clustering was observed in C18:1 PC membranes instead of C16:1 PC membranes observed for sx-1TM (Fig. 1g). These FRET data fit well with a quadratic curve ($ax^2 + bx + c$). In this empirical model, $-b / (2a)$ reflects the acyl chain lengths with the lowest clustering which are 16.5 and 17.3 for sx-1TM and sx-4TM, respectively. Based on our imaging ellipsometry (Fig. 1b), these acyl chain lengths correspond to membrane thicknesses of 3.6 nm for sx-1TM and 3.7 nm for sx-4TM. Together, our data indicate that clustering of TMDs is indeed determined by the hydrophobic matching with the local lipid environment; with a length difference of even a single residue resulting in a shift towards an approximately 1 Å thicker membrane area. Despite this difference, both syntaxins are expected to cluster since the plasma membrane has an average thickness of around 4 nm³⁰, which would result in pronounced sequestering of these proteins to regions of decreased thickness.

Synergy of ionic interactions and hydrophobic mismatch

Next we investigated how clustering caused by hydrophobic mismatch is influenced by phosphoinositides. Syntaxin 1 is known to interact with PI(4,5)P₂ and/or PI(3,4,5)P₃ via a conserved polybasic motif directly adjacent to the hydrophobic TMD (Fig. 1f)^{6,10,15,16}. Furthermore, both phosphoinositides are highly accumulated in at least a fraction of syntaxin 1 clusters in the plasma membrane^{6,10,17,18}. Using two-color super-resolution STED (stimulated emission depletion) microscopy, we confirmed the enrichment of PI(4,5)P₂ in syntaxin 1 clusters within plasma membrane sheets prepared from neuroendocrine PC12 cells (Fig. 2a). Syntaxin 4 also contains a polybasic motif at the membrane interface and, indeed, PI(4,5)P₂ was also enriched in syntaxin 4 clusters (Fig. 2b). In the plasma membrane, the density of PI(4,5)P₂ clusters (13.9 ± 1.6 clusters/ μm^2) was approximately 3 times higher than the cluster density of syntaxin 1 (4.5 ± 0.4 clusters/ μm^2) and syntaxin 4 (5.4 ± 0.7 clusters/ μm^2), which is not surprising when considering that PI(4,5)P₂ interacts with many other proteins in cells.^{32,33} We then reconstituted sx-1TM labeled with Rhodamine Red (donor fluorophore) and sx-4TM labeled with Atto647N (acceptor fluorophore) in LUVs and measured interaction by FRET (Fig. 2c). The presence of 1 mol% PI(4,5)P₂ in LUVs composed of brain PC caused an increase of the FRET efficiency, indicating that the TMDs of the two syntaxin isoforms co-clustered in the membrane when PI(4,5)P₂ was present. This clustering corroborates with our previous findings showing that interactions with polyanionic phosphoinositides can cluster syntaxin 1^{10,18}. Clustering was also observed in the presence of both PI(4,5)P₂ and cholesterol (Fig. 2d) showing that interactions with phosphoinositides act synergistically with hydrophobic mismatch. To further dissect how electrostatic repulsion between the polybasic linker region affects clustering by hydrophobic mismatch, we repeated our FRET assay in the presence of high concentrations of NaCl. Membrane clustering of sx-1TM was strongly promoted when electrostatic interactions were screened by the presence of 1 M NaCl. This indicates that the repulsion of the polybasic linkers of syntaxins counteracts hydrophobic mismatch and thereby limits clustering even in the presence of PI(4,5)P₂ (Fig. 2e).

For a better understanding of the membrane clustering, we carried out coarse-grain molecular simulations. In these simulations clustering of sx-1TM in membranes composed

of longer (C20:1) and shorter (C12:1) acyl chains was observed already at very short simulation time scales (~100 μ s). The simulation qualitatively reproduced the experimentally observed minimum of sx-1TM clustering in C16:1 membranes (Supplementary Fig. 1). However, a detailed analysis revealed that hydrophobic mismatch is not the only mechanism driving clustering. In fact the mismatch between the sx-1TM peptide and the membrane seems to be always negative even in thin membranes (Supplementary Fig. 2a–b). In such thinner membranes, syntaxin clustering was primarily promoted by increased homotypic protein-protein interactions which for membranes with shorter acyl chains were facilitated by the much larger rotational tilting angle relative to sx-1TM tilting angles in thicker membranes (Supplementary Figs. 1b, 2c-d)³⁴. In membranes with longer acyl chains, protein clustering was mainly caused by the more pronounced hydrophobic mismatch (Supplementary Fig. 2a), which causes a corresponding free energy penalty ΔG (Supplementary Fig. 2b), proportional to the protein-lipid mismatch interface. While strictly a free energy difference, the effective TMD extension (d) of the mismatch interface is rather constant for a given membrane and, therefore, the length of the protein-lipid interface mainly determines this free energy. Accordingly, and intuitively, we will refer to $\Delta G/d$ as a ‘line tension’.

We then further characterized the influence of homotypic TMD interactions on SNARE clustering. It was demonstrated previously that syntaxin 1 TMDs homodimerize in membranes despite the electrostatic repulsion of the cationic linker. This homodimerization depends on specific protein-protein interactions in the hydrophobic phase that can be disrupted by alanine substitutions of three hydrophobic side chains within the TMD (M267A, C271A, I279A)³⁵. To examine whether homodimerization contributes to homophilic clustering during hydrophobic mismatch, we reconstituted the corresponding sx-1TM mutant peptides and measured clustering by FRET. In membranes composed of C14:1 PC and C18:1 lipids (i.e. both thicker and thinner than required for optimal hydrophobic matching of syntaxin 1), the dimerization mutant clustered comparable to wild-type (Fig. 2f), indicating that at these conditions homotypic interactions were not required for clustering. These findings agree with the results from our simulations, where although clustering of the dimerization mutant was reduced compared to wild-type sx-1TM, the pronounced rotational mobility in thin membranes facilitated clustering of even the sx-1TM mutant. Here, the larger oligomers were absent and the shape of the cluster distribution resembled the wild type size distribution in membranes with longer acyl chains (Supplementary Fig. 2c-d). Together, these results demonstrate that the observed minimum of clustering in membranes without cholesterol is caused by a competition between protein-protein interactions and hydrophobic mismatch. For membranes with more pronounced negative mismatch (such as the plasma membrane environment for syntaxins) clustering is mainly driven by the line tension that results from the increased hydrophobic mismatch free energy.

Segregation of syntaxins 1 and 4 by hydrophobic mismatch

In the final set of experiments we asked whether the small difference in the length of the TMDs between syntaxin 1 and 4 might result in at least partial segregation into separate clusters because of hydrophobic mismatch between the TMDs. Thus, difference in length of

the TMDs may contribute, in addition to the well-established homophilic interactions between SNARE motifs^{8,9,22}, to the conspicuous segregation of the two syntaxins into different clusters (Fig. 3a). To address this question, we first reconstituted sx-1TM and sx-4TM in liposomes composed of a mixture of PC with different acyl chain lengths (C14:1 to C20:1) and measured clustering by FRET. Clustering of sx-1TM with sx-4TM, but not of sx-1TM to sx-1TM and sx-4TM to sx-4TM, was lower compared to liposomes containing only C18:1 PC (Fig. 3b). This indicates that syntaxin TMDs preferentially cluster in regions containing lipids with matching hydrophobic thickness and a heterogeneous plasma membrane environment can drive segregation of membrane proteins with different lengths of TMDs in distinct, membrane domains.

We then asked if hydrophobic mismatch can also cause segregation of syntaxin 1 and 4 in the complex environment of a plasma membrane. To this end, we transfected PC12 cells with truncation mutants of both syntaxins (sx-1TM and sx-4TM, similar to the fragments used in abovementioned experiments) N-terminally fused to GFP and RFP, respectively. When membrane sheets from these cells were analyzed by two-color superresolution STED microscopy³⁶, segregation of the two mutants in separate clusters was observed (Fig. 3c-e). To test if this segregation was due to the difference in length of the TMDs, we generated syntaxin 1 TMDs that were either two amino acids longer (sx-1TM^{+VG}) or three amino acids shorter (sx-1TM^{-IFG}) than wild type. In line with the hydrophobic mismatch hypothesis, clusters of the shorter sx-1TM^{-IFG} strongly segregated from sx4-TM clusters (Fig. 3e). In contrast, the longer sx-1TM^{+VG}, with a similar length of TMD as sx-4TM, showed significantly more co-localization with sx-4TM. Thus, our data show that increasing or reducing hydrophobic mismatch by altering the length of TMDs by only a few residues contributes not only to protein clustering but also to segregation into separate clusters.

Discussion

In summary, we have shown that hydrophobic mismatch due to cholesterol mediated thickening of the membrane can drive clustering of proteins in distinct membrane domains and that this clustering can be modulated by homotypic interactions of the TMD and electrostatic protein-lipid interactions (Fig. 4). These data provide an alternative to previously established models of plasma membrane patterning^{4,37} allowing for drawing of three major conclusions. One, differences between membrane thickness and the length of the hydrophobic TMDs can drive clustering of membrane proteins. The resulting free energy penalty due to hydrophobic mismatch can be described in terms of a line-tension, which is minimized by protein-protein clustering. Such hydrophobic mismatch not only explains protein clustering in the plasma membrane, but also may explain cargo sorting of proteins into specific intracellular compartments^{26,38,39}. Indeed, theoretical calculations of hydrophobic mismatch carried out many years ago revealed that “embedded inclusions” (i.e. membrane proteins) perturb the membrane thickness resulting in a state with defined spacing between neighboring “inclusion” clusters^{40,41}. Membrane proteins can thus recruit lipids and other membrane proteins with matching hydrophobic thicknesses and are thereby able to pattern biological membranes⁴².

Two, inclusion of physiological concentrations of cholesterol (~30 mol%) results in a substantial thickening (by ~0.8 nm) of the membranes and consequently profoundly increases membrane cluster formation by hydrophobic mismatch. This agrees well with previous studies in which leucine-alanine rich repeat peptides were used as models for TMDs^{24,25}. Our results provide a mechanistic explanation for cholesterol induced syntaxin 1 domains reported by Murray and Tamm^{15,16} and also explain the cholesterol dependence of syntaxin 1 clustering observed in the plasma membrane^{5,21}. However, in contrast to former interpretations^{19,43}, we now show that cholesterol induced cluster formation is the result of hydrophobic mismatch and does not require partitioning into separate membrane phases or domains enriched in cholesterol and sphingolipids ('lipid rafts').

Three, clustering of proteins in the plasma membrane can be further modulated by electrostatic interactions with phosphoinositides and ions. Here, ions and the charged phosphoinositide head groups can balance the charges of the positive residues found on the juxtamembrane regions of many proteins, thereby overcoming electrostatic repulsion^{14,32}. We recently showed that polyanionic phosphoinositides can even reinforce the clusters by providing "charge bridges"^{10,18}. At least in the case of syntaxins, homophilic interactions between the TMDs can also promote clustering especially in thinner membranes where the increased rotational mobility facilitates these interactions. Based on our data, we propose a model (Fig. 4) where hydrophobic mismatch and electrostatic protein-protein and protein-lipid interactions act synergistically and are required for the stabilization of syntaxin clusters in the plasma membrane. Finally, homotypic interactions between the SNARE motifs and/or heterotypic interactions with known binding partners (e.g. SNAP 25, Munc 18a) can capitalize on the underlying physical principles, increasing specificity of segregation by hydrophobic mismatch due to differences in TMD-length^{9,22}.

Our findings are important for the understanding of the cellular organization of membrane trafficking. It is well established that SNARE clusters act as a functional platforms for vesicle docking and fusion, either directly^{5,44-46} or by providing pools of active SNAREs (discussed in⁴⁷). Membrane clusters of syntaxin 1 are involved in Ca²⁺-regulated secretion and are completely excluded from clusters of syntaxin 4 involved in constitutive exocytosis^{8,48}. Our results show that this segregation is at least partially caused by hydrophobic mismatch between the 1-2 residue longer transmembrane helix of syntaxin 4 compared to syntaxin 1. The recruitment of hydrophobically matching lipids and membrane proteins by these SNAREs may also have functional consequences for membrane fusion. It is conceivable that the increased line tension of the membrane clusters induces local membrane curvature and lowers the energy barrier for membrane fusion⁴⁹⁻⁵¹, which could reduce the number of SNARE complexes required for fusion⁵²⁻⁵⁴. Thus, an important concept raised in this study is that membrane proteins not only seem to partition into lipid-dependent membrane domains, but that they are in fact essential determinants of these lipid domains in the first place. Given that proteins are very abundant in the plasma membrane (20 – 25 volume% of membrane)^{55,56}, these protein domains warrant being further investigated as a distinct membrane phase.

Methods

Proteins and lipids

Syntaxin 1 TMD (residues 266–288; sx-1TM) from *Rattus norvegicus*, syntaxin 4 TMD (residues 262–297; sx-4TM) from *Homo sapiens*, and syntaxin 1 TMD mutant (sx-1TM with the following mutations: M267A, C271A, I279A) were synthesized using Fmoc solid phase synthesis. The fluorescent dyes Atto647N NHS-ester (Atto-Tec) and Rhodamine red succinimidyl ester (Life Technologies) were coupled to the N-termini of sx-TM. The detailed synthesis is described in¹⁰.

C18:1 (1,2-dioleoyl-*sn*-glycero-3-phosphocholine), C14:1 (1,2-dimyristoleoyl-*sn*-glycero-3-phosphocholine), C16:1 (1,2-dipalmitoleoyl-*sn*-glycero-3-phosphocholine), C20:1 (1,2-dieicosenoyl-*sn*-glycero-3-phosphocholine), brain PI(4,5)P₂, doPI(4,5)P₂ (1,2-dioleoyl-*sn*-glycero-3-phosphatidyl-(1'-myo-inositol-4',5'-bisphosphate)) and cholesterol were purchased from Avanti Polar Lipids. Bodipy-labeled PI(4,5)P₂ (bodipy-FL-PI(4,5)P₂, C16) was from Echelon Biosciences and Top-Fluor labeled PI(4,5)P₂ was from Avanti Polar Lipids. The lipophilic fluorescent probe DiO (3,3'-dilinoleyloxacarbocyanine perchlorate) was from Life Technologies.

Vesicle formation

Large unilamellar vesicles (LUV) were prepared from PC of different acyl-chain lengths (C14:1, C16:1, C18:1, and C20:1) with or without 30 mol% cholesterol and/or 1 mol% PI(4,5)P₂. Lipid mixtures were prepared at a total concentration of approximately 30 mM lipids in chloroform as described in⁵⁷. After removal of the chloroform with a rotary evaporator, the lipid film was resuspended to 40 mM in methanol and fluorescently labeled peptides were added in 2,2-trifluoroethanol. The organic solvents were then evaporated and resuspended to 8 mM total lipid concentration in 50 mM HEPES buffer with 150 mM KCl (pH 7.4) unless otherwise indicated (i.e. no salt or 1 M NaCl). The multilamellar vesicles were then extruded through polycarbonate filters with 100 nm pore diameter (Avanti Polar Lipids).

FRET measurements

For FRET analysis we used Rhodamine Red coupled to sx-TM (donor) and Atto647N coupled to sx-TM (acceptor). Protein-to-lipid ratio in our FRET measurements was 1:1000. Excitation was at 560 nm, and emission was collected from 570 to 700 nm with 1 nm slit widths on a FluoroMax-2 (Horiba). We corrected for cross-talk arising from acceptor excitation with samples containing only the acceptor fluorophore. The obtained FRET spectra were normalized to the maximum donor emission at 580 nm. The FRET efficiency was calculated as the ratio of emission intensities at 660 nm (acceptor maximum) over 580 nm (donor maximum)^{15,16}.

Protein reconstitutions in polymer supported membranes

Glass cover slides used in microscopy were prepared as described in⁵⁸ and the supported lipid bilayer was generated by spincoating. For the spincoating (at 100 xg) we prepared a 40 mM lipid mixture consisting of a 2:1 molar ratio of phospholipids to cholesterol⁵⁹. Molar

ratios of DiO and sx-1TM to phospholipids were 1:5,000 and 1:10,000 respectively. After spincoating 10 μ L of the lipid solution, the lipid film was rehydrated in 1 mL of 50 mM HEPES buffer with 150 mM KCl (pH at 7.4).

Cell culture and immunofluorescence

We used the pheochromocytoma cell line PC12 from *Rattus norvegicus*⁶⁰. Lipofectamin LTX reagents from Life Technologies were used for transfection and cells were analyzed 24 hours posttransfection. Native membrane sheets were generated by gentle sonication described in^{8,10} and sonication buffer containing 20 mM K-HEPES pH 7.4, 120 mM K-gluconate, 20 mM K-acetate, 2 mM ATP and 0.5 mM DTT. Antibodies used for immunohistochemistry were syntaxin 1 HPC-1 IgG1 (Sigma, clone HPC-1) and rabbit polyclonal antiserum (Synaptic Systems); syntaxin 4 mouse monoclonal IgG1 (Synaptic Systems, clone number 139.2); IgM antibodies against PI(4,5)P₂ (Echelon, clone Z-A045); mouse monoclonal IgG2a anti-mCherry (Abcam, clone 1C51); and rabbit polyclonal anti-EGFP (Abcam). Secondary antibodies against IgG and IgM were labeled with Alexa Fluor 488C₅-maleimide (Life Technologies) or KK114-maleimide (gift from Vladimir Belov, MPI-BPC, Göttingen, Germany, described in⁶¹).

For transfection of PC12 cells we used synthetic chimeric constructs (Genscript) in the KpnI-HindIII restriction sites of pCEP4. The sequences for syntaxin 1A (sequence from *Rattus norvegicus* 262–297) N-terminally tagged with mCherry or mEGFP are given in Supplementary Table 1. The constructs coding for mCherry-tagged Sx-1TM^{IFG} (residues 257–285) and sx1TM^{+VG} (residues 257–288 with two additional amino acids at the C-terminus: V289, G290) were generated from the wild-type construct by Quick Change mutagenesis (Agilent Technologies).

Fluorescence Correlation Spectroscopy

For FCS we used home-built confocal beam-scanning microscopy setup with two color excitation by pulsed-diode lasers at 485 nm (pulse length 80 ps; LDH-P-635, PicoQuant) and 635 nm (pulse length 80 ps LDH-P-485B, PicoQuant)⁶². Emission filters were 540 ± 20 for the green channel and 670 ± 30 for the red channel. We used a 100x oil objective with 1.42 NA (Leica Microsystems). Avalanche single photon counting detectors were used (APD, SPCM-AQR-13-FC, Perkin Elmer Optoelectronics). FCS curves were fitted to a model for two-dimensional diffusion and with the axial radii of the focal volumes ω_{xy} , defined as the point where the measured fluorescence drops e^2 times relative to the maximum, of 200 and 250 nm for Bodipy-FL and Atto647N, respectively.

Two color STED microscopy

The STED images were acquired on a home built setup as described in³⁶. The basic setup resembles a standard confocal microscope with pulsed excitation at 595 nm and 640 nm wavelength. The fluorescence was collected from 600–640 nm and 660–720 nm by avalanche photo diodes (Excelitas, USA and Micro Photon Devices, Italy), allowing good spectral separation of the dyes used. Superresolution was achieved by silencing the fluorophores in the periphery of the diffraction limited excitation spot via stimulated emission induced by the STED laser, a 775 nm wavelength, 20 MHz pulsed fiber laser (IPG

Photonics). In combination with a 2π vortex phase plate (RPC Photonics, USA) and a $\lambda/4$ plate the typical “doughnut” shaped focal intensity distribution with its central zero was produced. Pulse energies from 3 to 8 nJ in the objectives back aperture yield a resolution of down to 30 nm. Using the same STED beam for both dyes inherently ensures a colocalization accuracy far below the resolution limit. As we record both color channels quasi simultaneously we do not have to correct for drift or channel misalignment. The hardware and data acquisition was controlled by the software ImSpector (<http://www.imspector.de>). The density of clusters was analyzed using the particle analysis plugin in the Fiji software⁶³.

Ellipsometry measurements

Lipid stock solutions ($c_{\text{lipid}} = 1\text{--}10$ mg/ml) were prepared in chloroform and transformed into lipid films by removal of the solvent in a nitrogen stream followed by 3h drying in vacuum. Multilamellar vesicles (MLV) were produced by resuspending the lipid films in buffer (50 mM HEPES pH 7.4 with 3 mM Ca^{2+}) at a concentration of 1 mg/ml. MLVs were transformed into small unilamellar vesicles (SUV) by sonication (50 W, 0.4 s pulse, 30 min) in a vessel resonator (Sonoplus HD 2070). Average vesicle size was 30–50 nm as determined by dynamic light scattering⁶⁴. Si-Wafers were cleaned in $\text{H}_2\text{O}_2/\text{NH}_3/\text{H}_2\text{O}$ 1:1:5 at 70°C for 15 min and afterwards hydrophilized for 1 min in O_2 -plasma. For preparing the lipid bilayer, freshly prepared SUVs were spreaded for 10–30 min on a hydrophilized Si-Wafer at a concentration of 0.2 mg/ml in 50 mM HEPES pH 7.4 with 3 mM Ca^{2+} . Measurements were carried out in the same buffer in a closed fluid chamber. Ellipsometry experiments were performed using an imaging ellipsometer EP³-SW from Accurion as described previously^{29,65}. This method offers the possibility to measure thin layer thicknesses in real time within a convective flow at defined temperature. The principal angle *del* determined by this method is proportional to layer thicknesses for sufficiently thin dielectric layers ($h < 30$ nm). Absolute height changes resulting from spreaded solid supported membranes (SSM) were computed from the angle *del* which is linearly related to the height for thin layers ($1 \text{ nm} \approx 0.91^\circ \text{ del}$) and assuming a refractive index of 1.5 for the all used lipids.

Molecular dynamics

The molecular dynamics simulations were performed with the GROMACS simulation package, version 4.5.5⁶⁶. We used the MARTINI coarse-grained model to simulate the lipids, amino acids and polar solvents^{67,68}. Within this model an unsaturated bond is modeled by a decreased equilibrium bond-angle (120° versus 180°), increased flexibility and increased polarity. In all simulations, the system was coupled to a constant temperature bath with a relaxation time of 1.0 ps⁶⁹. We performed our simulations at a temperature of 310 K. Periodic boundary conditions were applied to simulate bulk behavior. The time step used in the simulation was 20 fs. The dielectric constant in the simulations was $\epsilon_r = 15$. The neighbor-list was updated every 10 simulation steps. The pressure was weakly coupled to 1 bar with a relaxation time of 0.5 ps. In analogy to the other studies done with the MARTINI model, the time scales quoted in this work were scaled by a factor of four to approximately correct for the faster diffusion rates of water and lipids in the coarse-grained model⁴⁸.

Syntaxin 1 TMD (residues 266–288; sx-1TM) was modeled using the MARTINI model for proteins, which qualitatively captures the chemical nature of each individual amino acid and includes the secondary structure. In our simulations, sx-1TM and linker region (residues 257–265) were modeled as an α -helix. The backbone was modeled as a stiff and conserved spring with a length of 3.6 nm (residue 266 to 288). This model allows us to qualitatively investigate the relationship between membrane thickness, free-energy of protein insertion (hydrophobic mismatch), and protein clustering.

The simulations used to study spontaneous sx-1TM clustering contained 81 copies of sx-1TM embedded in a $30 \times 30 \mu\text{m}$ planar bilayer containing 2,560 lipids and 43,000 solvent beads. The total charge of the system was kept at zero with Cl^- counterions. To mimic the cell membrane, all sx-1TM peptides were embedded with their *N*-termini facing the same side of the membrane. These simulations were run for 60 μs . Complementary we also performed 4 μs simulations with smaller $7 \times 7 \text{ nm}$ bilayer system (128 lipids) that contained a single sx-1TM. These systems were used to calculate the membrane thickness adaptation, and membrane insertion free-energy of sx-1TM. The cluster-sizes (oligomer size) were calculated over the last 20 μs , using a distance cutoff-based cluster algorithm. Neighboring sx-1TMs were considered clustered if any of the beads in the trans-membrane region (residues 266–288) were within a distance of 0.7 nm.

To provide a quantitative measure for the hydrophobic-mismatch, i.e. the protein-lipid interactions, we applied thermodynamic integration techniques (with 4 μs simulation for each lambda point) to calculate the relative free-energy of membrane insertion for sx-1TM as a function of membrane thickness/composition. The exact protocol can be found elsewhere²⁵.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank L. Tamm, V. Kiessling (Charlottesville, VA), S. Rizzoli (Göttingen, Germany), and S. Safran (Rehovot, Israel) for advice and critical discussions, F. Roder and J. Piehler for introducing us to the PSM technology and P. Thiesen for discussions on use of imaging ellipsometry.

This work was supported by the Deutsche Forschungsgemeinschaft (SFB803) and the US National Institutes of Health (P01 GM072694) to RJ and the European Research Council (FP7 Starting Grant 336479-MEMPART) to GvdB

References

1. Singer SJ, Nicolson GL. The fluid mosaic model of the structure of cell membranes. *Science*. 1972; 175:720–731. [PubMed: 4333397]
2. Mukherjee S, Maxfield FR. Membrane domains. *Annu. Rev. Cell Dev. Biol.* 2004; 20:839–866. [PubMed: 15473862]
3. Kusumi A, et al. Paradigm shift of the plasma membrane concept from the two-dimensional continuum fluid to the partitioned fluid: high-speed single-molecule tracking of membrane molecules. *Annu. Rev. Biophys. Biomol. Struct.* 2005; 34:351–378. [PubMed: 15869394]

4. Lingwood D, Simons K. Lipid rafts as a membrane-organizing principle. *Science*. 2010; 327:46–50. [PubMed: 20044567]
5. Lang T, et al. SNAREs are concentrated in cholesterol-dependent clusters that define docking and fusion sites for exocytosis. *EMBO J*. 2001; 20:2202–2213. [PubMed: 11331586]
6. Aoyagi K, et al. The Activation of Exocytotic Sites by the Formation of Phosphatidylinositol 4,5-Bisphosphate Microdomains at Syntaxin Clusters. *J. Biol. Chem*. 2005; 280:17346–17352. [PubMed: 15741173]
7. Knowles MK, et al. Single secretory granules of live cells recruit syntaxin-1 and synaptosomal associated protein 25 (SNAP-25) in large copy numbers. *Proc. Natl. Acad. Sci. U.S.A.* 2010; 107:20810–20815. [PubMed: 21076040]
8. Sieber JJ, Willig KI, Heintzmann R, Hell SW, Lang T. The SNARE Motif Is Essential for the Formation of Syntaxin Clusters in the Plasma Membrane. *Biophys. J*. 2006; 90:2843–2851. [PubMed: 16443657]
9. Barg S, Knowles MK, Chen X, Midorikawa M, Almers W. Syntaxin clusters assemble reversibly at sites of secretory granules in live cells. *Proc. Natl. Acad. Sci. U.S.A.* 2010; 107:20804–20809. [PubMed: 21076041]
10. van den Bogaart G, et al. Membrane protein sequestering by ionic protein-lipid interactions. *Nature*. 2011; 479:552–555. [PubMed: 22020284]
11. Hong W, Lev S. Tethering the assembly of SNARE complexes. *Trends Cell Biol*. 2014; 24:35–43. [PubMed: 24119662]
12. Bacia K, Schuette CG, Kahya N, Jahn R, Schwille P. SNAREs prefer liquid-disordered over ‘raft’ (liquid-ordered) domains when reconstituted into giant unilamellar vesicles. *J. Biol. Chem*. 2004; 279:37951–37955. [PubMed: 15226320]
13. Lang T. SNARE proteins and ‘membrane rafts’. *J. Physiol. (Lond.)*. 2007; 585:693–698. [PubMed: 17478530]
14. Do Heo W, et al. PI(3,4,5)P3 and PI(4,5)P2 Lipids target proteins with polybasic clusters to the plasma membrane. *Science*. 2006; 314:1458–1461. [PubMed: 17095657]
15. Murray DH, Tamm LK. Clustering of syntaxin-1A in model membranes is modulated by phosphatidylinositol 4,5-bisphosphate and cholesterol. *Biochemistry*. 2009; 48:4617–4625. [PubMed: 19364135]
16. Murray DH, Tamm LK. Molecular mechanism of cholesterol- and polyphosphoinositide-mediated syntaxin clustering. *Biochemistry*. 2011; 50:9014–9022. [PubMed: 21916482]
17. James DJ, Khodthong C, Kowalchuk JA, Martin TFJ. Phosphatidylinositol 4,5-bisphosphate regulates SNARE-dependent membrane fusion. *J. Cell Biol*. 2008; 182:355–366. [PubMed: 18644890]
18. Khuong TM, et al. Synaptic PI(3,4,5)P3 is required for Syntaxin1A clustering and neurotransmitter release. *Neuron*. 2013; 77:1097–1108. [PubMed: 23522045]
19. Chamberlain LH, Burgoyne RD, Gould GW. SNARE proteins are highly enriched in lipid rafts in PC12 cells: Implications for the spatial control of exocytosis. *Proc. Natl. Acad. Sci. U.S.A.* 2001; 98:5619–5624. [PubMed: 11331757]
20. Chamberlain LH, Gould GW. The vesicle- and target-SNARE proteins that mediate Glut4 vesicle fusion are localized in detergent-insoluble lipid rafts present on distinct intracellular membranes. *J. Biol. Chem*. 2002; 277:49750–49754. [PubMed: 12376543]
21. Rickman C, et al. t-SNARE protein conformations patterned by the lipid microenvironment. *J. Biol. Chem*. 2010; 285:13535–13541. [PubMed: 20093362]
22. Sieber JJ, et al. Anatomy and dynamics of a supramolecular membrane protein cluster. *Science*. 2007; 317:1072–1076. [PubMed: 17717182]
23. Simons K, Gerl MJ. Revitalizing membrane rafts: new tools and insights. *Nature Publishing Group*. 2010; 11:688–699.
24. Kaiser HJ, et al. Lateral sorting in model membranes by cholesterol-mediated hydrophobic matching. *Proc. Natl. Acad. Sci. U.S.A.* 2011; 108:16628–16633. [PubMed: 21930944]
25. Schäfer LV, et al. Lipid packing drives the segregation of transmembrane helices into disordered lipid domains in model membranes. *Proc. Natl. Acad. Sci. U.S.A.* 2011; 108:1343–1348. [PubMed: 21205902]

26. Sharpe HJ, Stevens TJ, Munro S. A Comprehensive Comparison of Transmembrane Domains Reveals Organelle-Specific Properties. *Cell*. 2010; 142:158–169. [PubMed: 20603021]
27. Borgese N, Colombo S, Pedrazzini E. The tale of tail-anchored proteins: coming from the cytosol and looking for a membrane. *J. Cell Biol.* 2003; 161:1013–1019. [PubMed: 12821639]
28. Bulbarelli A, Sprocati T, Barberi M, Pedrazzini E, Borgese N. Trafficking of tail-anchored proteins: transport from the endoplasmic reticulum to the plasma membrane and sorting between surface domains in polarised epithelial cells. *J. Cell. Sci.* 2002; 115:1689–1702. [PubMed: 11950887]
29. Faiss S, Schuy S, Weiskopf D, Steinem C, Janshoff A. Phase transition of individually addressable microstructured membranes visualized by imaging ellipsometry. *J. Phys. Chem. B*. 2007; 111:13979–13986. [PubMed: 18044861]
30. Mitra K, Ubarretxena-Belandia I, Taguchi T, Warren G, Engelman DM. Modulation of the bilayer thickness of exocytic pathway membranes by membrane proteins rather than cholesterol. *Proc. Natl. Acad. Sci. U.S.A.* 2004; 101:4083–4088. [PubMed: 15016920]
31. Stein A, Weber G, Wahl MC, Jahn R. Helical extension of the neuronal SNARE complex into the membrane. *Nature*. 2009; 460:525–528. [PubMed: 19571812]
32. McLaughlin S, Wang J, Gambhir A, Murray D. PIP2 and proteins: interactions, organization, and information flow. *Annu. Rev. Biophys. Biomol. Struct.* 2002; 31:151–175. [PubMed: 11988466]
33. Di Paolo G, De Camilli P. Phosphoinositides in cell regulation and membrane dynamics. *Nature*. 2006; 443:651–657. [PubMed: 17035995]
34. Gambin Y, et al. Lateral mobility of proteins in liquid membranes revisited. *Proc. Natl. Acad. Sci. U.S.A.* 2006; 103:2098–2102. [PubMed: 16461891]
35. Laage R, Rohde J, Brosig B, Langosch D. A conserved membrane-spanning amino acid motif drives homomeric and supports heteromeric assembly of presynaptic SNARE proteins. *J. Biol. Chem.* 2000; 275:17481–17487. [PubMed: 10764817]
36. Göttfert F, et al. Coaligned dual-channel STED nanoscopy and molecular diffusion analysis at 20 nm resolution. *Biophys. J.* 2013; 105:L01–L03. [PubMed: 23823248]
37. Ritchie K, Kusumi A. Role of the membrane skeleton in creation of microdomains. 2004; 37:233–245.
38. Munro S. Sequences within and adjacent to the transmembrane segment of alpha-2,6-sialyltransferase specify Golgi retention. *EMBO J.* 1991; 10:3577–3588. [PubMed: 1935890]
39. Munro S. An investigation of the role of transmembrane domains in Golgi protein retention. *EMBO J.* 1995
40. Dan N, Berman A, Pincus P, Safran SA. Membrane-induced interactions between inclusions. *J. Phys. II France*. 1994; 4:1713–1725.
41. Aranda-Espinoza H, Berman A, Dan N, Pincus P, Safran S. Interaction between inclusions embedded in membranes. *Biophys. J.* 1996; 71:648–656. [PubMed: 8842204]
42. Anderson RGW, Jacobson K. A role for lipid shells in targeting proteins to caveolae, rafts, and other lipid domains. *Science*. 2002; 296:1821–1825. [PubMed: 12052946]
43. Salaün C, Gould GW, Chamberlain LH. Lipid raft association of SNARE proteins regulates exocytosis in PC12 cells. *J. Biol. Chem.* 2005; 280:19449–19453. [PubMed: 15769746]
44. Zilly FE, Sørensen JB, Jahn R, Lang T. Munc18-bound syntaxin readily forms SNARE complexes with synaptobrevin in native plasma membranes. *PLoS Biol.* 2006; 4:e330. [PubMed: 17002520]
45. Honigsmann A, et al. Phosphatidylinositol 4,5-bisphosphate clusters act as molecular beacons for vesicle recruitment. *Nat. Struct. Mol. Biol.* 2013; 20:679–686. [PubMed: 23665582]
46. Ohara-Imaizumi M, et al. Site of docking and fusion of insulin secretory granules in live MIN6 cells analyzed by TAT-conjugated anti-syntaxin 1 antibody and total internal reflection fluorescence microscopy. *J. Biol. Chem.* 2004; 279:8403–8408. [PubMed: 14676208]
47. van den Bogaart G, Lang T, Jahn R. Microdomains of SNARE proteins in the plasma membrane. *Curr. Top. Membr.* 2013; 72:193–230. [PubMed: 24210431]
48. Predescu SA, Predescu DN, Shimizu K, Klein IK, Malik AB. Cholesterol-dependent syntaxin-4 and SNAP-23 clustering regulates caveolar fusion with the endothelial plasma membrane. *J. Biol. Chem.* 2005; 280:37130–37138. [PubMed: 16118213]

49. Risselada HJ, Bubnis G, Grubmüller H. Expansion of the fusion stalk and its implication for biological membrane fusion. *Proc. Natl. Acad. Sci. U.S.A.* 2014; 111:11043–11048. [PubMed: 25024174]
50. Boucrot E, et al. Membrane fission is promoted by insertion of amphipathic helices and is restricted by crescent BAR domains. *Cell.* 2012; 149:124–136. [PubMed: 22464325]
51. Kozlov MM, et al. Mechanisms shaping cell membranes. *Curr. Opin. Cell Biol.* 2014; 29C:53–60. [PubMed: 24747171]
52. van den Bogaart G, et al. One SNARE complex is sufficient for membrane fusion. *Nat. Struct. Mol. Biol.* 2010; 17:358–364. [PubMed: 20139985]
53. Mohrmann R, de Wit H, Verhage M, Neher E, Sørensen JB. Fast vesicle fusion in living cells requires at least three SNARE complexes. *Science.* 2010; 330:502–505. [PubMed: 20847232]
54. Hernandez JM, Kreutzberger AJB, Kiessling V, Tamm LK, Jahn R. Variable cooperativity in SNARE-mediated membrane fusion. *Proc. Natl. Acad. Sci. U.S.A.* 2014; 111:12037–12042. [PubMed: 25092301]
55. Takamori S, et al. Molecular anatomy of a trafficking organelle. *Cell.* 2006; 127:831–846. [PubMed: 17110340]
56. Dupuy AD, Engelman DM. Protein area occupancy at the center of the red blood cell membrane. *Proc. Natl. Acad. Sci. U.S.A.* 2008; 105:2848–2852. [PubMed: 18287056]
57. Schuette CG, et al. Determinants of liposome fusion mediated by synaptic SNARE proteins. *Proc. Natl. Acad. Sci. U.S.A.* 2004; 101:2858–2863. [PubMed: 14981239]
58. Roder F, et al. Reconstitution of membrane proteins into polymer-supported membranes for probing diffusion and interactions by single molecule techniques. *Anal. Chem.* 2011; 83:6792–6799. [PubMed: 21838222]
59. Mennicke U, Salditt T. Preparation of solid-supported lipid bilayers by spin-coating. *Langmuir.* 2002; 18:8172–8177.
60. Heumann R, Kachel V, Thoenen H. Relationship between NGF-mediated volume increase and ‘priming effect’ in fast and slow reacting clones of PC12 pheochromocytoma cells. *Exp. Cell Res.* 1983; 145:179–190. [PubMed: 6303817]
61. Kolmakov K, et al. Red-emitting rhodamine dyes for fluorescence microscopy and nanoscopy. *Chemistry.* 2010; 16:158–166. [PubMed: 19950338]
62. Eggeling C, et al. New insight into lipid-protein membrane organization and its functionality with super-resolution STED microscopy. *Biophys. J.* 2013; 104:5.
63. Schindelin J, et al. Fiji: an open-source platform for biological-image analysis. *Nat. Meth.* 2012; 9:676–682.
64. Castorph S, et al. Synaptic vesicles studied by dynamic light scattering. *Eur. Phys. J E.* 2011; 34:1–11. [PubMed: 21706281]
65. Schuy S, et al. Structure and thermotropic phase behavior of fluorinated phospholipid bilayers: a combined attenuated total reflection FTIR spectroscopy and imaging ellipsometry study. *J. Phys. Chem. B.* 2008; 112:8250–8256. [PubMed: 18563929]
66. Hess B, Kutzner C, van der Spoel D, Lindahl E. GROMACS 4: Algorithms for highly efficient, load-balanced, and scalable molecular simulation. *J. Chem. Theory Comput.* 2008; 4:435–447. [PubMed: 26620784]
67. Monticelli L, et al. The MARTINI coarse-grained force field: extension to proteins. *J. Chem. Theory Comput.* 2008; 4:819–834. [PubMed: 26621095]
68. Marrink SJ, Risselada HJ, Yefimov S, Tieleman DP, de Vries AH. The MARTINI force field: coarse grained model for biomolecular simulations. *J. Phys. Chem. B.* 2007; 111:7812–7824. [PubMed: 17569554]
69. Berendsen HJC, Postma JPM, van Gunsteren WF, DiNola A, Haak JR. Molecular dynamics with coupling to an external bath. *J. Chem. Phys.* 1984; 81:3684.




Figure 1. Clustering of syntaxin isoforms by hydrophobic mismatch

(a) Positive and negative hydrophobic mismatch caused by differences between membrane thickness and TMD length. (b) Bilayer thickness determined by imaging ellipsometry of supported lipid bilayers composed of C14:1, C16:1, C18:1 and C20:1 PC with and without 30 mol% cholesterol (three independent experiments \pm SD). Solid lines show linear regression analyses (slopes of 0.15 and 0.25 for without and with cholesterol, respectively). (c) Scheme of the clustering assay for sx-1TM in 100 nm sized liposomes using FRET. A mixture of sx-1TM was used that were N-terminally labeled with Rhodamine Red and

Atto647N. **(d)** Clustering determined by FRET using liposomes composed of PC of increasing acyl chain lengths, without (green) and with (pink) 30 mol% cholesterol. Independently, normalized lateral diffusion coefficients of sx-1TM labeled with Atto647N were determined by FCS (blue). Error bars: range from two independent reconstitutions, three technical repeats each. **(e)** Clustering of sx-1TM (labeled with Rhodamine Red) in supported lipid bilayers (C18:1 PC) in the absence (top) and presence (bottom) of 30 mol% cholesterol measured by confocal microscopy. The membrane was visualized with the lipophilic dye DiO. (Scale bars, 2 μm) **(f)** Domain organization and alignment of the TMD regions of syntaxin 1 and syntaxin 4. The TMDs and adjacent polybasic patches are marked in magenta and red, respectively. **(g)** Clustering of human sx-1TM (green) and sx-4TM (magenta) measured by FRET without cholesterol as in panel **d**. Solid lines show fits with quadratic curves. Error bars: range from two independent reconstitutions, three technical repeats each.




Figure 2. Ionic interactions and hydrophobic mismatch act synergistically on syntaxin clustering (a,b) Both syntaxin 1 and syntaxin 4 clusters colocalize with PI(4,5)P₂. Plasma membrane sheets derived from PC12 cells were immunostained for PI(4,5)P₂ (green), syntaxin 1 (a; red) and syntaxin 4 (b; red) and imaged by two color STED microscopy. The graphs show the fluorescence intensity profiles as indicated in the figures (syntaxin 1 and 4: red profiles; PI(4,5)P₂: green profiles); yellow bars depict the positions of the domains. (c,d) Both PI(4,5)P₂ and cholesterol enhance co-clustering of sx-1TM and sx-4TM. FRET was measured after reconstitution in large unilamellar vesicles containing sx-1TM labeled with

Rhodamine Red (donor fluorophore) and sx-4TM with Atto647N (acceptor) composed of porcine brain PC without or with 1 mol% PI(4,5)P₂, and/or 30 mol% cholesterol (\pm range from two independent reconstitutions, three technical repeats each). (e) Clustering of sx-1TM by FRET in the presence or absence of 150 mM or 1 M NaCl and in DOPC liposomes in the absence or presence of 3 mol% PI(4,5)P₂. (f) Clustering of sx-1TM (green) and the sx-1TM oligomerization mutant (purple), monitored by FRET, in C14:1 PC or C18:1 PC liposomes. Error bars: range from two independent reconstitutions, three technical repeats each. Scale bars, 2 μ m.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 3. Differences in the length of the TMDs contribute to segregation of syntaxin 1 and 4 to distinct clusters in the plasma membrane

(a) Two-color STED microscopy of PC12 cell sheets immunostained for syntaxin 1 and syntaxin 4 shows segregation of endogenous proteins into separate clusters. (b) Reduced co-clustering of sx-1TM and sx-4TM in membranes composed of a mixture of PC with different acyl chain lengths. FRET assay is similar to Fig. 1c, but now measuring clustering of TMDs in liposomes composed of either C18:1 PC or an equimolar mixture of C14:1, C16:1, C18:1 and C20:1 PC. All liposomes contained 1 mol% PI(4,5)P₂ and 30 mol%

cholesterol (\pm range from two independent reconstitutions, three technical repeats each). **(c)** Same as in panel **a** but now using PC12 cell sheets derived from cells expressing truncation mutants of syntaxin 1 and 4 (sx-1TM; sx-4TM) that are fused to mCherry and EGFP, respectively, and immunostained with antibodies against mCherry and EGFP. **(d)** Control experiment of PC12 cells co-expressing sx-1TM tagged with either mCherry or EGFP, showing colocalization (Scale bar, 2 μ m). **(e)** Overlap of clusters in membrane sheets from PC12 cells transfected with various sx-1TM and sx-4TM mutants (all mCherry and EGFP tagged, respectively) measured by determining the Pearson-correlation coefficient. Sx-1FL and Sx-4FL; full length constructs of syntaxin 1 and syntaxin 4, respectively (from panel **a**). Each analysis included at least ten sheets from three independent transfections (***) $p < 0.001$, two sided, unpaired *t-test*; error bars show SEM). Scale bars, 2 μ m.




Figure 4. Synergistic model of syntaxin clustering in the membrane

Syntaxin membrane clustering is induced by a combination of hydrophobic mismatch (increased by cholesterol-induced membrane thickening) and electrostatic interactions with ions and PI(4,5)P₂. The membrane clusters are further refined by protein-protein interactions in the aqueous space.