



Published in final edited form as:

Nat Commun. ; 6: 7223. doi:10.1038/ncomms8223.

Piezo1 ion channel pore properties are dictated by C-terminal region

Bertrand Coste^{1,2,5,*}, Swetha E. Murthy^{2,5}, Jayanti Mathur³, Manuela Schmidt^{2,4}, Yasmine Mechioukhi¹, Patrick Delmas¹, and Ardem Patapoutian^{2,3}

¹Aix Marseille Université, CNRS, CRN2M-UMR7286, 13344 Marseille France

²Howard Hughes Medical Institute, Molecular and Cellular Neuroscience, The Scripps Research Institute, La Jolla, California 92037, USA

³Genomics Institute of the Novartis Research Foundation, San Diego, California 92121, USA

Abstract

Piezo1 and Piezo2 encode mechanically activated cation channels that function as mechanotransducers involved in vascular system development and touch sensing, respectively. Structural features of Piezos remain unknown. Mouse Piezo1 is bioinformatically predicted to have 30–40 transmembrane (TM) domains. Here, we find that nine of the putative inter-transmembrane regions are accessible from the extracellular side. We use chimeras between mPiezo1 and dPiezo to show that ion permeation properties are conferred by C-terminal region. We further identify a glutamate residue within a conserved region adjacent to the last two putative TM domains of the protein, that when mutated, affects unitary conductance and ion selectivity, and modulates pore block. We propose that this amino acid is either in the pore or closely associates with the pore. Our results describe important structural motifs of this channel family and lay the groundwork for a mechanistic understanding of how Piezos are mechanically gated and conduct ions.

Mechanotransduction is the process by which mechanical stimuli are converted into biological activity. Piezos are mechanically activated (MA) cation channels conserved through evolution and act as mechanotransducers in various biological processes. The single Piezo gene in flies is involved in nociception¹; zebrafish and mouse Piezo2, in touch sensation^{2–6}; zebrafish Piezo1, in red blood cell volume regulation⁷; and mouse Piezo1, in

Users may view, print, copy, and download text and data-mine the content in such documents, for the purposes of academic research, subject always to the full Conditions of use:http://www.nature.com/authors/editorial_policies/license.html#terms

*Correspondence: bertrand.coste@univ-amu.fr (B.C.).

⁴Present Address: Somatosensory Signaling Group; Max Planck Institute of Experimental Medicine, 37075 Goettingen, Germany

⁵These authors contributed equally to this work

Author contributions

B.C. and S.M. performed electrophysiological experiments. M.S. designed and initiated immunostaining experiments. J.M. generated chimera, mutant and tagged constructs and performed immunostaining experiments. Y.M. provided technical help. B.C., S.M. and A.P. designed experiments and wrote the manuscript. All authors discussed the paper.

Additional information

Supplementary Information includes 8 figures and 3 tables.

Competing financial interests: The authors declare no competing financial interests.

vascular development^{8,9}. In humans, mutations that alter channel gating of Piezo1 and 2 are linked to various disorders with dominant inheritance^{10–12}. Piezo proteins contain over 2000 amino acid residues with an estimated 30 to 40 transmembrane (TM) segments and are likely to form homo-tetramers in a complex weighing over 1.2 million daltons^{13,14}. Piezos lack homology with other proteins, and their structural features remain unknown. The large size and numerous hydrophobic domains of Piezos constitute technical challenges for structural analysis of the intact channel¹⁵. Basic questions regarding Piezo topology and the location of the ion permeation pathway remain unanswered, and yet these questions are crucial for a mechanistic understanding of how the channel is gated by mechanical forces, and how human disease-related point mutations affect channel function¹⁶.

We aimed to determine the topology of these large proteins and delineate the amino acid residues involved in the ion permeation pathway. Here, we provide experimental evidence to confirm the position of the N- and C- terminals, and 13 of the putative inter-hydrophobic loop regions of the protein. We show that the C-terminal region of the Piezo protein encompasses the pore. Within this region, we identify a glutamate residue involved in ion conduction properties. Our results lay the groundwork for understanding how Piezos are mechanically gated and conduct ions, and how Piezo mutations affect human biology.

Results

Transmembrane topology of Piezo channels

To characterize the transmembrane topology of mPiezo1 we combined bioinformatics analysis, immunostaining to detect extracellular tags inserted in predicted loop regions, and detection of intracellular phosphorylation sites by mass spectrometry (Fig 1). First, we used hydrophobicity plots, transmembrane segment predictions, and sequence alignment of functionally tested Piezo proteins (human, mouse and fly Piezos) to generate a predicted transmembrane topology of mPiezo1. This virtual topology was then tested experimentally by inserting Myc tags at mPiezo1 termini and in each predicted loop. In some cases, more than one tag was tested per predicted loop. Each of these constructs was subjected to an immunostaining protocol to test whether an anti-Myc antibody could recognize the Myc epitope in mPiezo1 in non-permeabilized cells, suggesting an extracellular topology (Fig. 1a and Supplementary Fig. 1). A negative signal would suggest an intracellular epitope or one that is masked by the cell membrane or another such mechanism¹⁷. Therefore, only tags that gave a positive signal were used to predict the topology. A total of 48 Myc-constructs were designed. 45 of these constructs showed staining with the Myc-antibody after permeabilization, among which only 10 constructs were positive for staining without permeabilization, and therefore predicted to be present extracellularly (Supplementary Fig. 2 and Supplementary Table 1). 2 among the 10 Myc constructs were 4 amino acids apart and can account only for a single extracellular loop, therefore 9 extracellular loops are labelled. The orientation of these extracellular loops agreed with the hydropathy/hydrophilicity plot predictions. Insertion of Myc tags at the N- or C-terminal, and at 7 out of the 9 extracellular loop positions did not affect channel function, and resulted in MA currents of amplitude comparable to WT currents. Whole-cell currents could not be detected from constructs with Myc tags at amino acid position 2071, 2075 or 2336, suggesting that tag insertions there

compromised channel function. Next, using mass spectrometry we analyzed phosphorylated sites within mPiezo1 purified protein to identify intracellular loops in the protein. A total of 24 phosphorylated peptides were detected corresponding to phosphorylation of 23 positions (Supplementary Table 2). Combining Myc-immunostaining and phosphorylation sites together with the bioinformatics analysis allowed us to construct a model topology which predicts 38 transmembrane domains with intracellular N- and C-termini (Fig. 1b). However, this might be an over-estimation of transmembrane domains, since the prediction heavily relies on bioinformatics, and because we were unable to get experimental evidence for many of the predicted extracellular loops. If we instead consider that Myc tags detected only after permeabilization may be indicative of intracellular or transmembrane positions, then a second model with less TM domains can be proposed (Fig. 1c). In this model, among the 35 Myc tags detected only after permeabilization, only one is predicted to be extracellular. Indeed, Myc 1126 is expected to be present in the extracellular loop labeled by Myc 1071 in both models because no predicted TM domain or hydrophobic domain is present between Myc 1071 and Myc 1126. Therefore, absence of Myc 1126 staining without permeabilization is likely due to accessibility issue. In this second model mPiezo1 is predicted to have minimum of 18 TM domains. In addition, if we further assume that the inter-hydrophobic loops with phosphorylated residues are the only 4 intracellular loops, then only 10 of the hydrophobic regions would be counted as transmembrane domains. This extreme scenario would predict a topology with a minimum of 10 TM domains.

Pore related properties of mPiezo1/dPiezo chimeras

We next set out to identify residues involved in the ion permeation pathway of the channel. mPiezo1 and dPiezo differ in their biophysical pore properties, such as sensitivity to the polycationic pore blocker ruthenium red (RR) and single-channel conductance¹⁴. We therefore generated chimeras between mPiezo1 and dPiezo based on our predicted mPiezo1 topology, and assayed them for these pore-related properties (Fig. 2). All chimeras were expressed in HEK293T cells and tested in whole-cell configuration to determine RR sensitivity of MA currents, and in cell-attached configuration to determine unitary conductance of stretch-activated channels.

Replacing the C-terminal half of mPiezo1 amino acid sequence with dPiezo (chimera mP1¹⁻¹³¹⁵/dP¹³¹⁷⁻²⁵⁴⁸) generated a non-functional channel but the reciprocal substitution (chimera dP¹⁻¹³¹⁶/mP1¹³¹⁶⁻²⁵⁴⁷) gave rise to a chimeric channel with MA currents sensitive to 30 μ M RR, similar to mPiezo1 but not to dPiezo (Fig. 2a–c and Supplementary Table 3). Single-channel conductance of this chimeric channel was also similar to mPiezo1 (Fig. 2d–f and Supplementary Table 3), demonstrating that biophysical pore properties of Piezo channels are determined by the C-terminal half of the protein sequence. We subsequently generated more chimeras by swapping shorter C-terminal regions between the two proteins. Chimera mP1¹⁻¹⁷¹⁴/dP¹⁷⁵³⁻²⁵⁴⁸ was not functional, and chimera dP¹⁻¹⁹²⁹/mP1¹⁹⁷⁴⁻²⁵⁴⁷ could not be generated, but the complementary chimeras, dP¹⁻¹⁷⁵²/mP1¹⁷¹⁵⁻²⁵⁴⁷ and mP1¹⁻¹⁹⁷³/dP¹⁹³⁰⁻²⁵⁴⁸, were functional and analyzed. Chimera dP¹⁻¹⁷⁵²/mP1¹⁷¹⁵⁻²⁵⁴⁷, displayed RR sensitivity and unitary conductance similar to mPiezo1, whereas chimera mP1¹⁻¹⁹⁷³/dP¹⁹³⁰⁻²⁵⁴⁸ had channel properties comparable to dPiezo (Fig. 2 and Supplementary Table 3). These results suggest that domain(s) required for determining

conductance and pore block are encoded by the region of the protein sequence starting from mPiezo1 amino acids 1974. Interestingly, this segment of dPiezo and mPiezo1 sequences are 40% identical compared to 24% identity for the entire sequences. We also observed that whole-cell inactivation properties were likely dependent on this C-terminal region, as chimera dP¹⁻¹³¹⁶/mP¹³¹⁶⁻²⁵⁴⁷ and dP¹⁻¹⁷⁵²/mP¹⁷¹⁵⁻²⁵⁴⁷ display inactivation kinetics comparable to mPiezo1, whereas mP¹⁻¹⁹⁷³/dP¹⁹³⁰⁻²⁵⁴⁸ displays faster inactivation similar to dPiezo (Supplementary Table 3).

Neutralization of acidic residues in the C-terminal region

Results from mouse-fly chimeras that replaced shorter regions of the C-terminus were inconclusive, as chimeras were either non-functional or displayed weak functionality preventing their characterization. Therefore, we performed site-directed mutagenesis to identify the molecular determinants that control ion permeation properties of Piezo channels. Single point mutations in pore-forming domains of channels alter ion selectivity, single-channel conductance, and pore block, and this approach has been widely used to prove that a protein is truly a pore forming subunit¹⁸⁻²⁴. Studies of classical Na⁺ and Ca²⁺ channels have shown that divalent/monovalent selectivity depends on charge density in the selectivity filter, the narrowest region that lines the pore. For example, the high calcium selectivity of Ca²⁺ channels depends on the high negative charge density in their selectivity filter¹⁸. Similarly, TRPV1-4 channels, which are non-selective cation channels with a P_{Ca}/P_{Na} ratio in the 1-10 range²⁵, contain a single acidic residue in their selectivity filter²⁶. The neutralization of this acidic residue in TRPV1 and TRPV4 reduces the channel permeability for Ca²⁺ and its affinity for the pore blocker RR^{27,28}. Since mPiezo1 channels are non-selective cation channels¹³, we hypothesized that acidic residues may be a component of the putative selectivity filter controlling ion permeation properties in mPiezo1. To this end, based on the data from our last chimeric channel (mP¹⁻¹⁹⁷³/dP¹⁹³⁰⁻²⁵⁴⁸), we performed an alanine scan of acidic residues located in inter- or intra- hydrophobic domains encompassed by residues 1974-2547. Since there are no acidic residues within or adjacent to the last two predicted TM domains we restricted our scan to the 15 Aspartate and Glutamate residues of mPiezo1 1974-2188 region (Fig. 3a). Of note, only 4 of these 15 acidic residues are not conserved in dPiezo; one of them (D2013) is replaced by another acidic residue (E), and the remaining three (D1987, D2014 and E2170) by neutral residues (N, N and Q, respectively).

Of the 15 individual alanine mutants, D1975A and D2034A were non-functional mutants, but the neutralization of aspartate residues with asparagine substitution resulted in functional clones. Therefore, we used D1975N and D2034N for further characterization. None of the mutations significantly affected the channel's sensitivity to 30μM RR (60 to 79% block at -80 mV) (Fig. 3b and Supplementary Table 3). Remarkably, one of the 15 mutations, E2133A, caused a strong reduction of single-channel current amplitude (Fig. 3c-e and Supplementary Table 3). The unitary conductance of E2133A mutant is half the unitary conductance of WT channels (14.4 ± 0.5 and 29.1 ± 0.4 pS, respectively; mean \pm s.e.m.). This suggests that E2133 is located in or near the mPiezo1 ion-conducting pathway.

Mutation of conserved residues surrounding E2133

Alignment of Piezo protein sequences from various species highlights conservation of the region around mPiezo1 E2133 (Supplementary Fig. 4). We therefore tested if any of these conserved residues affect channel conductance. Removing Ca^{2+} from the recording solution abolishes Ca^{2+} block of these channels, resulting in a larger single-channel conductance and hence a higher signal to noise ratio. Indeed, mPiezo1 unitary conductance is doubled in Na^+ - or K^+ -based solution compared to in standard pipette solution¹⁴. Under these conditions, E2133A unitary conductance is reduced by 2-fold compared to WT channels (26.8 ± 0.6 and 58.6 ± 1.2 pS, respectively) (Supplementary Fig. 5a–c and Supplementary Table 3). Individual alanine substitution of conserved residues upstream and downstream of E2133 resulted either in non-functional channels (P2129A, L2131A, R2135A and W2140A) or in channels with unitary conductance similar to WT channels (F2130A; 56.3 ± 1.7 pS, V2132A; 59.5 ± 1.7 pS and L2134A; 59.4 ± 2.1 pS; mean \pm s.e.m.). Additionally, we observed that in the absence of divalent ions, D2139A and D2144A mutations, which are downstream of E2133, result in slightly increased single-channel conductance compared to WT (64.2 ± 1.2 and 69.6 ± 1.8 pS, respectively; mean \pm s.e.m.). We next tested ion selectivity of these alanine mutants by measuring current-voltage relationship in 150 mM CsCl / 100 mM CaCl_2 asymmetric conditions (Supplementary Fig. 5d). Only E2133A mutation showed a significant shift in reversal potential relative to WT (Supplementary Fig. 5e). These results show that among the conserved residues in this region, E2133 is an important determinant of channel conductance and ion selectivity. mPiezo1 E2133 is surrounded by hydrophobic domains. Based on the predicted topologies presented in Figure 1, this residue could be present in the membrane facing the extracellular side, in an intracellular loop, or in a structure analogous to a re-entrant loop (Fig. 4a).

E2133 mutations influence mPiezo1 pore properties

We next tested if changing the size/charge of the residue at position 2133 modulates channel conductance, ion selectivity, or RR sensitivity (Fig. 4 and 5). Three additional mutations were generated: E2133Q neutralizes the charge while keeping a similar side chain length, E2133D keeps the negative charge but reduces the side chain length, and E2133K replaces the negative charge with a positive one. In Na^+ -based pipette solution, all these substitutions significantly altered unitary conductance compared to WT channels. Unitary conductance of E2133Q was similar to E2133A mutants, whereas unitary conductance of E2133D and E2133K increased or drastically decreased compared to WT (77.5 ± 3.0 and 20.0 ± 1.3 pS, respectively; mean \pm s.e.m.) (Fig. 4b–d and Supplementary Table 3). To determine if these mutations affected ion selectivity of the channel, recordings were again made in 150 mM CsCl / 100 mM CaCl_2 asymmetric conditions. Under these conditions, E2133Q and E2133K mutations significantly shift the channel's reversal potential to the left (5.0 ± 0.5 and 2.6 ± 0.4 mV, respectively, vs. 10.3 ± 0.6 mV for WT; mean \pm s.e.m.) (Fig. 4e–f). These results show that E2133 is a determinant of ion selectivity, and the shift in reversal potential could be caused either due to a decrease in calcium permeability or due to an increase in chloride permeability. Therefore, we explored these two possibilities by determining chloride and calcium permeability for E2133Q and E2133K mutants. We found that mPiezo1 permeates chloride with $P_{\text{Cl}}/P_{\text{Na}}$ of 0.14, and that E2133K mutation significantly increases this ratio by 1.5 fold compared to WT (Supplementary Fig. 6a–b). This shows that introducing a positive

charge at position 2133 decreases mPiezo1 preference for cations over anions. On the contrary, chloride permeability of E2133Q mutant was similar to WT and therefore cannot explain the difference in reversal potential observed in the 150 mM CsCl / 100 mM CaCl₂ condition, suggesting that Ca²⁺ permeability was altered in this mutant. Indeed, we found a P_{Ca}/P_{Cs} ratio of 1.21 for mPiezo1 which is decreased by 1.3 fold in E2133Q mutant (Supplementary Fig. 6c–d). These results highlight the importance of the charge present at position 2133 for mPiezo1 conductance and ion selectivity.

RR is a voltage-dependent extracellular pore blocker of mPiezo1¹⁴. Mutation of acidic residues in the selectivity filter of TRPV channels affects sensitivity to RR^{26–28}. Therefore, we tested if RR sensitivity was modulated by E2133Q, D, and K substitutions (Fig. 5). At –80 mV, 30 μM RR blocked MA currents from E2133Q and E2133D channels by 74.6 ± 6.2 and 85 ± 2 % respectively, compared to 71.4 ± 4.4 % for WT channels (mean ± s.e.m.). E2133K, on the other hand, was only slightly blocked (8.1 ± 7.1 %) by 30 μM RR. RR dose-response curves measured at –80 mV gave IC₅₀ values of 5.62 ± 1.27 μM and 1.26 ± 0.28 μM for E2133Q and E2133D, respectively, compared to 4.71 ± 0.83 μM for WT (mean ± s.e.m.), illustrating that RR sensitivity of E2133Q is similar to WT whereas E2133D channels are slightly more sensitive. Remarkably, RR IC₅₀ for E2133K channels was about 80-fold higher than WT (Fig. 5b). Neutralization of the E2133 residue did not relieve RR block; therefore, the glutamate is unlikely to be the RR binding site. The positively charged lysine substitution, however, could alter the local charge in the vicinity of the putative RR binding site, thereby altering the electrostatic interaction between RR and the channel²⁹. Additional studies in this region will determine the molecular mechanism of RR binding and inhibition in Piezos.

To test if the E2133K substitution, which had the most prominent consequence on pore-properties, could also alter pressure sensitivity of the channel, we compared the current-pressure relationships of WT and E2133K channels in cell-attached mode. We did not detect a change in pressure sensitivity, suggesting that although E2133K affects pore permeation properties it does not alter the channel's mechanical sensitivity (Supplementary Fig. 7).

Conserved glutamate determines Piezo2 pore properties

We next tested if the conserved glutamate residue is also functionally required to determine pore properties in another Piezo ion channel by mutating the homologous residue in mPiezo2, E2416, into a Lysine (Fig. 6a). We focused on mouse Piezo2 instead of *Drosophila* Piezo since the fly channel is not blocked by RR, and its small single-channel currents are challenging to record (a 2-fold decrease in conductance would be difficult to validate). We recorded stretch-activated single-channel currents from HEK293T cells expressing mPiezo2 and found the unitary conductance in Na⁺-based pipette solution condition to be 56.9 ± 1.3 pS. Under these conditions, the unitary conductance of mPiezo2 E2416K mutant was reduced by 1.7 fold relative to WT (32.6 ± 1.0 pS; mean ± s.e.m.) (Fig. 6b–d and Supplementary Table 3). Additionally, E2416K mutation also alters ion selectivity of the channel (Supplementary Fig. 8), decreasing cation over anion preference illustrated by a 3-fold increase of P_{Cl}/P_{Na} ratio (0.08 and 0.25 for WT and E2416K, respectively) (Fig. 6e–f). We next tested the extent of RR block of MA whole-cell currents from WT and

E2416K mPiezo2 expressing cells. Compared to WT, E2416K mutant channels were significantly less sensitive to 30 μ M RR (70.8 ± 1.4 and 49.6 ± 4.6 % blocks, respectively; mean \pm s.e.m.) (Fig. 6g–h and Supplementary Table 3). Therefore, similar to mPiezo1, lysine substitution of E2416 in mPiezo2 alters conductance, ion selectivity and pore block of the channel. These results suggest a general requirement for this conserved glutamate residue in controlling pore properties and indicates that the hydrophobic regions adjacent to the last 2 predicted TM domains of the channel could be either part of the pore domain or closely associated with the pore across Piezo channels.

Discussion

In this study, we describe the first structural motifs of Piezo channel family. We combined bioinformatics, intramolecular Myc-tag and phosphorylated site detection to begin to unlock the transmembrane topology of mPiezo1 protein. By developing a chimeric approach between mouse Piezo1 and drosophila Piezo, we show that the portion of mPiezo1 from 1974 to C-terminus is essential for ion permeation properties. Further, we show that a conserved glutamate residue within hydrophobic regions neighboring the last 2 putative TM domains alter pore-dependent properties, suggesting that this residue is either within or in close proximity to the mPiezo1 pore-forming domain.

We provide the first experimental evidence validating certain regions of bioinformatically predicted mPiezo1 topology. We show that mPiezo1 has intracellular N- and C-terminals, and verify the orientation of 13 inter-hydrophobic domain loops of the protein. Piezo proteins are encoded by more than 2000 residues and lack homology with any other protein¹³. The large size of the protein has proved challenging in delineating the exact topology of this non-selective cation channel. Several bioinformatics predictions estimate a topology consisting of anywhere between 30 to 40 TMs domains^{11,16,30–36}. We experimentally addressed the transmembrane topology of Piezo proteins by generating 48 Myc insertion mutations in mPiezo1 predicted loops. Immuno-detection of these tags in non-permeabilized conditions was used to probe extracellular loops. We combined this approach with the detection of 23 phosphorylation sites by mass spectrometry to identify intracellular loops. Our results verify that 9 of these putative loops have an extracellular orientation, and that along with the N- and C- terminal of the protein, 4 loops are intracellularly located. Although this advances our knowledge on topology, we cannot predict exact number of transmembrane domains. Based on this data, a model that predicts 38 TM domains best agrees with the bioinformatics prediction of mPiezo1 topology. However, we do not have experimental data proving that many of the loops that are proposed to be present on the extracellular side in this model are indeed accessible from outside the cell. This is not completely unexpected, since Myc tags inserted in small extracellular loops might not be accessible to the Myc Antibody in live cells. But it is also possible that the lack of labeling could point to these loops being intracellular. Catering to this possibility, if we assume that all Myc tags not observed in live unpermeabilized cells are intracellular or within transmembrane regions, then mPiezo1 could have only 18 TM domains. Furthermore, if we assume that only regions that are phosphorylated form intracellular loops, then a topology with a minimum of 10 putative TM domains could be proposed. The actual number of TM domains could be anywhere in between the two extreme cases. Regardless, our results here

represent the first experimentally based information on Piezo topology, and will be invaluable for future studies addressing structural features of this channel.

We generated chimeric proteins between mPiezo1 and dPiezo and assayed their pore-related properties¹⁴. We show that the mPiezo1 C-terminal domains encoded by the residues 1974–2547 confer unitary conductance, ion selectivity and RR sensitivity. Therefore, we conclude that C-terminal portion of mPiezo1, which is 1/5th of the whole protein, includes most if not all of the pore domain. Interestingly, according to our predicted mPiezo1 topology, almost 70% of the reported hPiezo1 and hPiezo2 mutations map to this region, including hPiezo1 (T2127M) and hPiezo2 (T2356M) which are in close proximity to our identified glutamate residue (see below)¹⁶, emphasizing the importance of this region in channel function.

Following-up on our chimeric channel screen, we employed site-directed mutagenesis of acidic residues within hydrophobic regions encoded by residues 1974–2188 to show that neutralization of a single glutamate residue results in a 2-fold decrease in unitary conductance. Additional substitutions of aspartate, lysine or glutamine instead of the glutamate residue altered unitary conductance, ion selectivity and sensitivity to RR, thereby suggesting that E2133 contributes to mPiezo1 pore properties. The role of this glutamate residue in dictating ion permeation properties of Piezos was further confirmed when the lysine substitution of the homologous residue in mPiezo2 (E2416K) also modulated unitary conductance, cation over anion selectivity and RR sensitivity.

Alignment of Piezo proteins from distant species such as parasites shows that mPiezo1 E2133 glutamate residue is the only charged residue in a conserved stretch of amino acids that form a PF(X₂)E(X₆)W motif³⁴. We found that P and W residues are indeed essential for channel function, as mutations at these positions result in non-functional channels. The impact of the glutamate residue on pore-properties and its location in a highly conserved motif is suggestive of its presence in a structure similar to a selectivity filter present in most members of known ion channel families. However, only lysine substitution of the E2133 altered mPiezo1 RR sensitivity, suggesting that this residue is not the RR binding site. Furthermore, the magnitude of the impact the lysine substitution has on ion-selectivity and single channel conductance is rather mild for a model where the selectivity filter of mPiezo1 or 2 channels would constitute a ring of E2133 or E2416 residues from associated subunits. This suggests that the glutamate residue may not lie in the selectivity filter but could be located close enough to the pore to allosterically modulate its properties.

The E2133 residue lies in a region flanked by hydrophobic residues. However, the different topology models differ in the prediction of how these hydrophobic domains are oriented. It is possible that the hydrophobic domains form full transmembrane segments, and that E2133 could be accessible from or close to the extracellular side (Fig 1b). Alternatively, it might be located closer to the intracellular side, potentially as part of a re-entrant loop (Fig 1c). The true orientation of this E2133 region relative to the rest of the channel will be elucidated by future structural analysis.

Although mutating the glutamate residue impacts pore properties in mouse Piezos, this residue is conserved between mouse (E2133/2416 in mPiezo1/2) and fly Piezos (E2091 in

dPiezo) and therefore cannot explain their pore-related differences. Therefore, other unidentified differences between mouse and fly Piezos in the C-terminal region must explain the large variations in their biophysical pore-properties.

Overall, we provide strong evidence that the protein region corresponding to mPiezo1 residues 1974 to C-terminus encompass the domains that form the pore of the channel, and that mPiezo1 E2133 residue (or mPiezo2 E2416) is at least in the vicinity of the pore permeation pathway and can influence Piezo pore properties.

Methods

mPiezo1 topology prediction

Using the sequence of mammalian Piezos, topology predictions were made using Hopp-Woods hydrophilicity/antigenicity plots as well as Kyte Doolittle hydrophobicity plots. A combination of these results was used to predict the final topology. Our approach was validated to predict the topology of various TRP channels and CLC channels³⁸. For Hopp-Woods plots we used a window size of 7 to predict highly hydrophilic/antigenic regions. For Kyte Doolittle hydrophobicity plots we used a window size of 17 to predict TM regions and a window size of 9 to predict surface regions over the full-length amino acid sequence. For ambiguous regions, fragments of the amino acid sequence were analyzed separately using the same parameters. The results were then compared with putative transmembrane domains predicted by TMHMM2 and Phobius algorithms. A total of 38 transmembrane regions were predicted. Myc tags were inserted in predicted inside and outside loops near hydrophilic residues, preferably in the middle of the loop. All Myc constructs were generated using the QuickChange II XL site directed mutagenesis kit (Agilent Technologies). Myc tags were also introduced at the N and C-terminal regions. All constructs were verified by sequencing.

HEK293T cells were plated on poly-D-lysine coated MatTek-dishes which were additionally coated with laminin (10 or 20 $\mu\text{g/ml}$). Cells were transfected with each construct separately (2 μg cDNA each) using Fugene HD (Promega). 24–48 hrs after transfection live labeling was carried out by incubating transfected HEK293T cells with Myc 9E11 (1:50; Santa Cruz Biotechnology) at 37°C for 20 mins or 1 hr. After 5 washes with warm medium, cells were incubated with secondary antibodies conjugated to Alexa Fluor 546 (1:200; Lifetechnologies) for 10–20 min at RT. Cells were washed 5 times with PBS, fixed with 2% PFA/PBS for 20–30 min and imaged at an Olympus (Tokyo, Japan) Fluoview 500 confocal microscope by illumination with the HeNe green 543 nm laser.

Immunostaining after fixation and permeabilization of cells was carried out essentially as described¹³. In brief cells were fixed with 4% PFA/PBS for 10 min, washed 5 times with PBS, permeabilized in PBS containing 0.4% Triton X-100 and blocked with 10% normal goat serum in PBS followed by incubation with antibodies. The Myc 9E11 (1:100; Santa Cruz Biotechnology) antibody was used to confirm predicted outside and inside loops and Piezo1 antibodies (1:100) to confirm Piezo1 expression. After three washes in PBS secondary antibodies conjugated to Alexa Fluor 546 and 633 (1:200; Lifetechnologies) were applied. Immunostainings were imaged at an Olympus (Tokyo, Japan) Fluoview 500 confocal microscope by sequential illumination using the HeNe green 543 nm laser and the

HeNe red 633 nm laser. The live labelling and permeabilized staining was repeated 2 or more times for each construct to confirm results.

Detection of phosphorylation sites

mPiezo1-GST proteins were purified from HEK293T cells transfected with mPiezo1-GST cDNAs¹⁴. After incubation with cell lysates overnight at 4 °C, the glutathione beads were washed four times in a buffer containing 25 mM NaPIPES, 140 mM NaCl, 0.6% CHAPS, 0.14% phosphatidylcholine (PC), 2.5 mM dithiothreitol (DTT), and a cocktail of protease inhibitors and eluted with 100 mM glutathione in a buffer containing 25 mM NaPIPES, 50 mM Tris, 0.6% CHAPS, 0.14% PC, 2.5 mM DTT and a cocktail of protease inhibitors. The eluant was dialysed against a buffer containing 25 mM NaPIPES, 0.6% CHAPS, 0.14% PC, 2.5 mM DTT and a cocktail of protease inhibitors. The protein samples were denatured, reduced, and alkylated with 2-iodoacetamide prior to precipitation with Acetonitrile. The pellet was re-dissolved in 8M Urea and ProteaseMax detergent, then diluted to 1M Urea and digested with trypsin or chymotrypsin. A portion of each digest was analyzed by nanoscale liquid chromatography-mass spectrometry (nanoLC/MS) on the Orbitrap Velos (Thermo scientific) without further processing. The remainder of the digests was enriched for phosphopeptides using TiO₂ beads and also analyzed by nanoLC/MS on the Orbitrap Velos. The data was searched using a small database (about 450 proteins) to identify peptides from mPiezo1 (~73% sequence coverage) and sites of phosphorylation and the search results combined in Scaffold 3. The search results were filtered for high confidence (95% at the peptide level) and MS/MS indicating sites of phosphorylation were inspected.

Generation of chimeras and mutants

mPiezo1 and dPiezo chimeras were generated by swapping specific regions in the C-terminal end of the protein. The beginning of a swapped region was chosen at a stretch of amino acids that were conserved between mPiezo1 and dPiezo. These junction sites are marked in Supplementary Fig 2. Domain swapping was carried out using the QuickChange II XL site directed mutagenesis kit from Agilent technologies. Mega-primers were designed flanking the region of interest with overlapping base pairs between the donor and the recipient cDNA. The mega-primer was generated by PCR amplification and gel purification. The PCR product was then used as a primer in the mutagenesis reaction with the recipient cDNA as the template. PCR cycling conditions were according to the manufacturer's instructions. Reactions were transformed into XL-gold competent cells and colonies screened. All positive chimera clones were verified by full length DNA sequencing.

Point mutants were also generated using the QuickChange II XL site directed mutagenesis kit according to the manufacturer's instructions, and confirmed by full length DNA sequencing. dPiezo chimeras were cloned into pIres2-EGFP vector and all mPiezo1 chimeras and mutants were cloned into pcDNA3.1(-) vector which was modified to include an IRES-EGFP element. mPiezo2 (E2416K) was cloned in pCMV-Sport6 vector.

Cell culture and transient transfection

Human Embryonic Kidney 293T (HEK293T) cells were grown in Dulbecco's Modified Eagle Medium containing 4.5 mg.ml⁻¹ glucose, 10% fetal bovine serum, 50 units.ml⁻¹

penicillin and 50 $\mu\text{g}\cdot\text{ml}^{-1}$ streptomycin. Cells were plated onto 12-mm round glass poly-D-lysine coated coverslips placed in 24-well plates and transfected using lipofectamine 2000 (Invitrogen) according to the manufacturer's instruction. All plasmids were transfected at a concentration of 600 to 1000 $\text{ng}\cdot\text{ml}^{-1}$. mPiezo2(E2416K) was co-transfected with GFP at a concentration of 300 $\text{ng}\cdot\text{ml}^{-1}$. Cells were recorded from 12–48 hours post transfection.

Electrophysiology

Patch-clamp experiments were performed in standard whole-cell or cell-attached mode using Axopatch 200B amplifier (Axon Instruments). Currents were sampled at 20 kHz and filtered at 2 kHz. Leak currents before mechanical stimulations were subtracted off-line from the current traces. Voltages were not corrected for a liquid junction potential (LJP) except for ion selectivity experiments (Figures 4E–G, 5E–F, S6D–E and S7). LJP was calculated using Clampex 10.3 software. All experiments were done at room temperature.

Solutions

For whole-cell patch clamp recordings, recording electrodes had a resistance of 2–3 $\text{M}\Omega$ when filled with internal solution composed of (in mM) 133 CsCl, 1 CaCl₂, 1 MgCl₂, 5 EGTA, 10 HEPES (pH 7.3 with CsOH), 4 MgATP and 0.4 Na₂GTP. The extracellular solution was composed of (in mM) 133 NaCl, 3 KCl, 2.5 CaCl₂, 1 MgCl₂, 10 HEPES (pH 7.3 with NaOH) and 10 glucose. For ion selectivity experiments, internal solution used was (in mM) 150 CsCl, 10 HEPES (pH 7.3 with CsOH) and extracellular solution consisted of (in mM) 100 CaCl₂ and 10 HEPES (pH 7.3 with CsOH). $P_{\text{Cl}}/P_{\text{Na}}$ was measured in extracellular solution composed of (in mM) 30 NaCl, 10 HEPES and 225 Sucrose (pH 7.3 with NaOH) and intracellular solution consisted of (in mM) 150 NaCl and 10 HEPES (pH 7.3 with NaOH). $P_{\text{Ca}}/P_{\text{Cs}}$ was measured in extracellular solution composed of (in mM) 50 Ca-gluconate, 0.5 CaCl₂, 10 HEPES and 170 Sucrose (pH 7.3 with CsOH), and intracellular solution composed of (in mM) 149 Cs-methanesulfonate, 1 CsCl and 10 HEPES (pH 7.3 with CsOH). For ruthenium red (RR) experiments, 10 or 100 mM RR stock solution was prepared in water and diluted to working concentrations in extracellular solution.

For cell attached patch clamp recordings, external solution used to zero the membrane potential consisted of (in mM) 140 KCl, 1 MgCl₂, 10 glucose and 10 HEPES (pH 7.3 with KOH). Recording pipettes were of 2–3 $\text{M}\Omega$ resistance when filled with standard solution composed of (in mM) 130 mM NaCl, 5 KCl, 1 CaCl₂, 1 MgCl₂, 10 TEA-Cl and 10 HEPES (pH 7.3 with NaOH) or divalent-free pipette solution consisted of (in mM) 150 mM NaCl, 10 HEPES (pH 7.3 with NaOH).

Whole-cell mechanical stimulation

For whole-cell recordings, mechanical stimulation was achieved using a fire-polished glass pipette (tip diameter 3–4 μm) positioned at an angle of 80° relative to the cell being recorded. Downward displacement of the probe towards the cell was driven by Clampex-controlled piezo-electric crystal microstage (E625 LVPZT Controller/Amplifier; Physik Instrumente). The probe had a velocity of 1 $\mu\text{m}/\text{ms}$ during the ramp phase of the command for forward movement and the stimulus was applied for 150 ms. To assess the mechanical sensitivity of a cell, a series of mechanical steps in 0.5 or 1 μm increments was applied every

10–20 s. For RR experiments, once an efficient stimulus eliciting MA current for a cell was achieved, repeated stimuli at that distance were then given every 20–30 seconds. Cells were continuously perfused and at least 3 consecutive stable recording traces were achieved before switching between solutions. Current parameters were measured from 3 or more averaged traces. For I–V relationship recordings, voltage steps were applied 700 ms before mechanical stimulation (150 ms) from a holding potential of –60 mV. Voltage steps were given from –80 mV to +80 in 20 mV increments. The starting position of the stimulation probe relative to the cell is not tightly controlled and varies between cells. Therefore responses were recorded at different probe displacements from cell to cell allowing recording of MA currents with similar amplitude.

Permeability ratio measurements

Reversal potential for each cell in the mentioned solution was determined by interpolation of the respective current-voltage data. Permeability ratios were calculated by using the following Goldman-Hodgkin-Katz (GHK) equations:

P_{Cl}/P_{Na} ratios:

$$E_{rev} = \frac{RT}{F} \ln \frac{P_{Na} [Na]_o + P_{Cl} [Cl]_i}{P_{Na} [Na]_i + P_{Cl} [Cl]_o}$$

P_{Ca}/P_{Cs} ratios:

$$E_{rev} = \frac{RT}{F} \ln \left(\sqrt{\frac{4P_{Ca} [Ca]_o}{P_{Cs} [Cs]_i} + \frac{1}{4}} - \frac{1}{2} \right)$$

where, RT/F has the value 25.69 at 25°C.

Unitary conductance measurement

Stretch-activated currents were recorded in the cell-attached patch clamp configuration. Membrane patches were stimulated with 500 ms negative pressure pulses through the recording electrode using Clampex controlled pressure clamp HSPC-1 device (ALA-scientific). Since the single-channel amplitude is independent of the pressure intensity, the most optimal pressure stimulation was used to elicit responses that allowed single-channel amplitude measurements. These stimulation values were largely dependent on the number of channels in a given patch of the recording cell. Single-channel amplitude at a given potential was measured from trace histograms of 5 to 20 repeated recordings. Histograms were fitted with Gaussian equations using Clampfit 10.3 software or multi-peak fitting analysis of IGOR Pro software. Single-channel slope conductance for each individual cell was calculated from linear regression curve fit to single-channel I–V plots. We sometimes observed sub-conductance states in single-channel recording of chimeric and mutant channels, but we focused the analysis on the state that was prevalent.

Concerning mPiezo2 WT and mutant, stretch-activated currents were detected in only ~50% of HEK293T transfected cells but never in cells expressing ires2-EGFP only. This frequency is lower than the one of mPiezo1 and related mutants (~90% positive patches). It may be due to either lower expression of mPiezo2 related constructs or to unknown functional difference between mPiezo1 and 2. In standard pipette solution mPiezo2 unitary conductance is 23.7 ± 0.6 pS (Supplementary Table 3).

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We thank Tim Jegla for discussions on pore prediction, Allain Francisco for help with molecular biology, Dan Mason from Genomics Institute of the Novartis Research Foundation for assistance with mass spectrometry experiments, and Jorg Grandl and Nancy Hong for critical reading of the manuscript. We thank Bailong Xiao for technical help and for sharing unpublished data. This work was supported by grants from Agence Nationale de la Recherche (ANR-12-PDOC-0005-01) to B.C. and by National Institutes of Health (NS083174) to A.P., who is also an investigator of the Howard Hughes Medical Institute.

References

1. Kim SE, Coste B, Chadha A, Cook B, Patapoutian A. The role of *Drosophila* Piezo in mechanical nociception. *Nature*. 2012; 483:209–212. [PubMed: 22343891]
2. Faucherre A, Nargeot J, Mangoni ME, Jopling C. piezo2b regulates vertebrate light touch response. *J Neurosci*. 2013; 33:17089–17094. [PubMed: 24155313]
3. Ikeda R, et al. Merkel cells transduce and encode tactile stimuli to drive abeta-afferent impulses. *Cell*. 2014; 157:664–675. [PubMed: 24746027]
4. Maksimovic S, et al. Epidermal Merkel cells are mechanosensory cells that tune mammalian touch receptors. *Nature*. 2014
5. Woo SH, et al. Piezo2 is required for Merkel-cell mechanotransduction. *Nature*. 2014
6. Ranade SS, et al. Piezo2 is the major transducer of mechanical forces for touch sensation in mice. *Nature*. 2014; 516:121–125. [PubMed: 25471886]
7. Faucherre A, Kissa K, Nargeot J, Mangoni ME, Jopling C. Piezo1 plays a role in erythrocyte volume homeostasis. *Haematologica*. 2014; 99:70–75. [PubMed: 23872304]
8. Li J, et al. Piezo1 integration of vascular architecture with physiological force. *Nature*. 2014
9. Ranade SS, et al. Piezo1, a mechanically activated ion channel, is required for vascular development in mice. *Proc Natl Acad Sci U S A*. 2014
10. Albuisson J, et al. Dehydrated hereditary stomatocytosis linked to gain-of-function mutations in mechanically activated PIEZO1 ion channels. *Nat Commun*. 2013; 4:1884. [PubMed: 23695678]
11. Bae C, Gnanasambandam R, Nicolai C, Sachs F, Gottlieb PA. Xerocytosis is caused by mutations that alter the kinetics of the mechanosensitive channel PIEZO1. *Proc Natl Acad Sci U S A*. 2013; 110:E1162–1168. [PubMed: 23487776]
12. Coste B, et al. Gain-of-function mutations in the mechanically activated ion channel PIEZO2 cause a subtype of Distal Arthrogyriposis. *Proc Natl Acad Sci U S A*. 2013; 110:4667–4672. [PubMed: 23487782]
13. Coste B, et al. Piezo1 and Piezo2 are essential components of distinct mechanically activated cation channels. *Science*. 2010; 330:55–60. [PubMed: 20813920]
14. Coste B, et al. Piezo proteins are pore-forming subunits of mechanically activated channels. *Nature*. 2012; 483:176–181. [PubMed: 22343900]
15. Kamajaya A, Kaiser JT, Lee J, Reid M, Rees DC. The Structure of a Conserved Piezo Channel Domain Reveals a Topologically Distinct beta Sandwich Fold. *Structure*. 2014

16. Volkers L, Mechoukhi Y, Coste B. Piezo channels: from structure to function. *Pflugers Arch*. 2014
17. Schmidt M, Dubin AE, Petrus MJ, Earley TJ, Patapoutian A. Nociceptive signals induce trafficking of TRPA1 to the plasma membrane. *Neuron*. 2009; 64:498–509. [PubMed: 19945392]
18. Hille, B. *Ion channels of excitable membranes*. 3. Sinauer; 2001.
19. Voets T, Nilius B. The pore of TRP channels: trivial or neglected? *Cell Calcium*. 2003; 33:299–302. [PubMed: 12765676]
20. Brelidze TI, Niu X, Magleby KL. A ring of eight conserved negatively charged amino acids doubles the conductance of BK channels and prevents inward rectification. *Proc Natl Acad Sci U S A*. 2003; 100:9017–9022. [PubMed: 12843404]
21. Hansen SB, Wang HL, Taylor P, Sine SM. An ion selectivity filter in the extracellular domain of Cys-loop receptors reveals determinants for ion conductance. *J Biol Chem*. 2008; 283:36066–36070. [PubMed: 18940802]
22. Kellenberger S, Auberson M, Gautschi I, Schneeberger E, Schild L. Permeability properties of ENaC selectivity filter mutants. *The Journal of general physiology*. 2001; 118:679–692. [PubMed: 11723161]
23. Kang L, Gao J, Schafer WR, Xie Z, Xu XZ. C. elegans TRP family protein TRP-4 is a pore-forming subunit of a native mechanotransduction channel. *Neuron*. 2010; 67:381–391. [PubMed: 20696377]
24. Yan Z, et al. Drosophila NOMPC is a mechanotransduction channel subunit for gentle-touch sensation. *Nature*. 2013; 493:221–225. [PubMed: 23222543]
25. Owsianik G, Talavera K, Voets T, Nilius B. Permeation and selectivity of TRP channels. *Annual review of physiology*. 2006; 68:685–717.
26. Liao M, Cao E, Julius D, Cheng Y. Structure of the TRPV1 ion channel determined by electron cryo-microscopy. *Nature*. 2013; 504:107–112. [PubMed: 24305160]
27. Garcia-Martinez C, Morenilla-Palao C, Planells-Cases R, Merino JM, Ferrer-Montiel A. Identification of an aspartic residue in the P-loop of the vanilloid receptor that modulates pore properties. *J Biol Chem*. 2000; 275:32552–32558. [PubMed: 10931826]
28. Voets T, et al. Molecular determinants of permeation through the cation channel TRPV4. *J Biol Chem*. 2002; 277:33704–33710. [PubMed: 12093812]
29. MacKinnon R, Miller C. Mutant potassium channels with altered binding of charybdotoxin, a pore-blocking peptide inhibitor. *Science*. 1989; 245:1382–1385. [PubMed: 2476850]
30. Andolfo I, et al. Multiple clinical forms of dehydrated hereditary stomatocytosis arise from mutations in PIEZO1. *Blood*. 2013; 121:3925–3935. [PubMed: 23479567]
31. Bagriantsev SN, Gracheva EO, Gallagher PG. Piezo proteins: regulators of mechanosensation and other cellular processes. *J Biol Chem*. 2014; 289:31673–31681. [PubMed: 25305018]
32. Delmas P, Coste B. Mechano-gated ion channels in sensory systems. *Cell*. 2013; 155:278–284. [PubMed: 24120130]
33. Nilius B, Honore E. Sensing pressure with ion channels. *Trends Neurosci*. 2012; 35:477–486. [PubMed: 22622029]
34. Prole DL, Taylor CW. Identification and analysis of putative homologues of mechanosensitive channels in pathogenic protozoa. *PloS one*. 2013; 8:e66068. [PubMed: 23785469]
35. Xiao R, Xu XZ. Mechanosensitive channels: in touch with Piezo. *Curr Biol*. 2010; 20:R936–938. [PubMed: 21056836]
36. Zarychanski R, et al. Mutations in the mechanotransduction protein PIEZO1 are associated with hereditary xerocytosis. *Blood*. 2012; 120:1908–1915. [PubMed: 22529292]
37. McHugh BJ, et al. Integrin activation by Fam38A uses a novel mechanism of R-Ras targeting to the endoplasmic reticulum. *J Cell Sci*. 2010; 123:51–61. [PubMed: 20016066]
38. Dutzler R, Campbell EB, Cadene M, Chait BT, MacKinnon R. X-ray structure of a Cl⁻ chloride channel at 3.0 Å reveals the molecular basis of anion selectivity. *Nature*. 2002; 415:287–294. [PubMed: 11796999]




Figure 1. Transmembrane topology of mPiezo1

(a) Representative images (3 experimental replicates) of Myc labeling in mPiezo1-Myc transfected HEK293T cells. Myc tags were inserted at position 897 (upper panels) or 753 (lower panels). Immunostaining was performed prior (left panels) or after (right panels) cell permeabilization. Scale bar: 20 μ m. (b-c) Schematic of transmembrane topology of mPiezo1 protein showing positions of Myc-tags detected extracellularly (green) and phosphorylated sites detected by mass-spectrometry (blue). Panel (c) also shows positions of Myc-tags detected only after permeabilization (orange).




Figure 2. Ruthenium red sensitivity and single-channel properties of mPiezo1 and dPiezo chimeras

(a) Schematic representations illustrating portions of the protein sequence from mPiezo1 (black) or dPiezo (red) for each construct. (b) Representative (from 5–8 experimental replicates) MA current traces at -80 mV from cells transfected with constructs shown in (a) before, during or after application of 30 μM RR. Each trace is an average of 3–5 trials. Probe stimulation displacements are indicated. (c) Percent inhibition of MA currents in cells transfected with specified constructs in the presence of 30 μM RR (n= 8, 5, 6, 5 and 5; mean ± s.e.m.). One-way ANOVA with Dunn's comparison to mPiezo1 or dPiezo, * $P < 0.05$, ** $P < 0.01$. (d) Representative (from 5–7 experimental replicates) stretch-activated channel openings elicited at -180 mV from cells transfected with specified constructs. dPiezo and mP¹¹⁻¹⁹⁷³/dP¹⁹³⁰⁻²⁵⁴⁸ traces are displayed after applying 1kHz digital filter. (e) Average I-V relationships of stretch-activated single channels in cells transfected with specified constructs. Single-channel amplitude was determined as the amplitude difference in Gaussian fits of full-trace histograms. (f) Unitary conductance of stretch-activated channels from cells transfected with specified constructs. Conductance is calculated from the slope of linear regression line of individual cell single-channel I-V relationships (n= 7, 5, 5, 5 and 6; mean ± s.e.m.). One-way ANOVA with Dunn's comparison to mPiezo1 or dPiezo, * $P < 0.05$, ** $P < 0.01$.




Figure 3. Neutralization of mPiezo1 acidic residues

(a) Protein sequence of mPiezo1 C-terminal region beginning at residue 1971. Mutated acidic residues and other acidic residues are highlighted (red and orange, respectively). Grey bars indicate hydrophobic regions. (b) Average block of MA currents at -80 mV by 30 μM RR in cells transfected with mPiezo1 WT and specified mutants (n = 2–6, mean ± s.e.m.). (c) Unitary conductance of stretch-activated channels in cells transfected with WT mPiezo1 and specified mutants. Conductance is calculated from the slope of linear regression line of individual cell single-channel I-V relationships (n = 3–13, mean ± s.e.m.; One-way ANOVA with Dunn's comparison to WT, **P<0.01). (d) Representative (from 13 and 5 experimental replicates) stretch-activated channel openings elicited at specified potentials from mPiezo1 WT and E2133A transfected cells. (e) Average I-V relationships of stretch-activated single channels in mPiezo1 WT and E2133A transfected cells (n = 13 and 5, respectively; mean ± s.e.m.). Single-channel amplitude was determined as the amplitude difference in Gaussian fits of full-trace histograms.




Figure 4. Mutations of a conserved Glutamate residue alter mPiezo1 pore properties

(a) Protein sequence surrounding E2133 with putative topology models of mPiezo1 C-ter region. E2133 is highlighted (orange) and green bars indicate hydrophobic regions. (b) Representative (from 4–7 experimental replicates) stretch-activated channel openings at –80 mV from cells transfected with mPiezo1 WT, E2133Q, E2133D and E2133K. Stimulation intensities are –15, –10, –20 and –50 mm Hg, respectively. (c) Average I-V relationships of stretch-activated single channels in cells transfected with mPiezo1 WT, E2133Q, E2133D and E2133K (n = 7, 5, 5 and 4, respectively; mean ± s.e.m.). Single-channel amplitude was determined as the amplitude difference in Gaussian fits of full-trace histograms. (d) Unitary conductance calculated from the slope of linear regression line of individual cells (mean ± s.e.m.; One-way ANOVA with Dunn’s comparison to WT, **P<0.01). (e) Representative (from 6–9 experimental replicates) traces of MA currents recorded with 150 mM CsCl based intracellular solution and 100 mM CaCl₂ extracellular solution from mPiezo1 WT (black), E2133Q (grey), E2133D (red) and E2133K (blue) transfected cells. Currents are elicited from –69.6 to +50.4 mV, 20 mV. Scale bars: 100 pA, 50 ms. Probe stimulation displacements are 3, 4, 6 and 7 μm, respectively. (f) Average I-V relationships of MA

currents recorded from mPiezo1 WT, E2133Q, E2133D and E2133K expressing cells with 150 mM CsCl based intracellular solution and 100 mM CaCl₂ extracellular solution (n = 8, 6, 6 and 9, respectively). Inset: Average reversal potentials from individual cells corresponding to panel (f) experiments (WT: 10.3 ± 0.6 mV; E2133Q: 5.0 ± 0.5 mV; E2133D: 12.1 ± 1.0 mV and E2133K: 2.6 ± 0.4 mV; mean \pm s.e.m.; One-way ANOVA with Dunn's comparison to WT, **P<0.01). Panels (b), (c) and (d) experiments were performed in cell-attached configuration with Na⁺ as the only permeating cation in the recording pipette.




Figure 5. RR sensitivity of E2133 mutants

(a) Representative (from 3–10 experimental replicates) MA current traces from cells transfected with mPiezo1 WT, E2133Q, E2133D and E2133K at -80 mV before, during or after application of 30 μM RR. Each trace is an average of 2–4 trials. Probe stimulation displacements are 4.5, 5, 6 and 8 μm, respectively. Scale bars: 300 pA, 100 ms. (b) RR concentration-inhibition curves on MA currents at -80 mV in cells transfected with specified constructs (n = 2–10 per data point; mean ± s.e.m.). One-way ANOVA with Dunn's comparison to WT done separately for each RR concentration, **P<0.01, *P<0.05.




Figure 6. Lysine substitution of E2416 alters pore properties of mPiezo2 channels
(a) Conservation of mPiezo1 E2133 at position 2416 on mPiezo2 protein. **(b)** Representative (from 9 and 6 experimental replicates) stretch-activated channel openings at -80mV from cells transfected with mPiezo2 WT and E2416K. Stimulation intensities are -15 and -20 mm Hg, respectively. **(c)** Average I-V relationships of stretch-activated single channels in mPiezo2 WT and E2416K transfected cells (n = 9 and 6, respectively; mean ± s.e.m.). Single-channel amplitude was determined as the amplitude difference in Gaussian fits of full-trace histograms. **(d)** Single-channel conductance calculated from the slope of linear regression line of individual cell single-channel I-V relationships (mean ± s.e.m.; Mann-Whitney test, ***P<0.001). **(e)** Average I-V relationships of MA currents recorded from mPiezo2 WT and E2416K expressing cells with 150 mM NaCl based intracellular solution and 30 mM NaCl extracellular solution. Top left inset: typical recording traces for WT (black) and E2416K (green) from -71.7 to +48.3 mV, 20 mV. Scale bars: 100 pA, 50 ms. Probe stimulation displacements are 9 and 8 μm, respectively. Bottom right inset: Average reversal potential (mean ± s.e.m.; n = 6 and 5, respectively; Mann-Whitney test **P<0.01). **(f)** P_{Cl⁻}/P_{Na⁺} permeability ratios of MA currents from cell transfected with mPiezo2 WT, and

E2416K (mean \pm s.e.m.; n = 6 and 5, respectively; Mann-Whitney test $**P < 0.01$). **(g)** Representative (from 5 experimental replicates) MA current traces from mPiezo2 WT and E2416K transfected cells at -80 mV before, during or after application of 30 μ M RR. Each trace is an average of 2–4 trials. Probe stimulation displacements are 5 and 7 μ m, respectively. Scale bars: 100 pA, 100 ms. **(h)** Average block of MA currents at -80 mV by 30 μ M RR in mPiezo2 WT and E2416K transfected cells (n = 5 for each; mean \pm s.e.m.; Mann-Whitney test, $**P < 0.01$). Panel **(b)**, **(c)** and **(d)** experiments were performed in cell-attached configuration with Na⁺ as the only permeating cation in the recording pipette.