



HHS Public Access

Author manuscript

BBA Clin. Author manuscript; available in PMC 2016 December 01.

Published in final edited form as:

BBA Clin. 2015 December 1; 4: 81–84. doi:10.1016/j.bbacli.2015.08.001.

No Evidence of Antibodies against GAD65 and Other Specific Antigens in Children with Autism

Simran Kalra¹, Peter D. Burbelo², Ahmad Bayat³, Kathryn H. Ching³, Audrey Thurm¹, Michael J. Iadarola³, and Susan E. Swedo¹

¹Pediatrics and Developmental Branch, National Institutes of Mental Health, National Institutes of Health, Bethesda, MD.

²Dental Clinical Research Core, National Institute of Dental and Craniofacial Research, National Institutes of Health, Bethesda, MD.

³Department of Perioperative Medicine, Clinical Center, National Institutes of Health, Bethesda, MD.

Abstract

Background—The presence of autoantibodies has been proposed as evidence for a role of autoimmunity in autism. This report investigates the prevalence of autoantibodies in children with autism using the luciferase immunoprecipitation systems (LIPS) immunoassay technology. A panel of autoantibody targets against several known and candidate neurological autoantigens, autoimmune-associated autoantigens and viruses was employed.

Methods—Serological analysis was performed on typically developing children (n = 55), developmentally delayed children without autism (n = 24) and children diagnosed with autism (n=104). Autoantibodies were measured against glutamic acid decarboxylase-65 (GAD65), a CNS autoantigen proposed to be associated with autism and against Ro52, glial fibrillary acidic protein, tyrosine hydroxylase, aquaporin-4, and gamma-enolase, the mouse mammary tumor virus and the xenotropic murine leukemia virus. Antibody levels and seropositivity prevalence were analyzed for statistically significant differences between the three groups.

Results—The majority of the children (98%) were seronegative for all targets in the antigen panel. No GAD65 seropositive children were detected in the cohort. Several low level seropositive sera against several of the protein targets were identified in isolated children in each of the three groups, but there was no difference in prevalence.

Conclusion—Using this panel of antigens and a sensitive, robust assay, no evidence of unusual immunoreactivity was detected in children with autism, providing evidence against a role of autoimmunity against several previously implicated proteins in autism spectrum disorder pathogenesis.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conflict of Interest

The authors declare no conflict of interest.

General significance—The idea that autoantibodies represent an underlying cause or are biomarkers for autism pathophysiology is not supported by this report.

1. Introduction

Autism spectrum disorder (ASD) is a behaviorally defined neurodevelopmental disorder [1]. The deficits in social-communication and presence of restricted interests and repetitive behaviors result in lifelong impairments and disability. ASD has been reported to affect as many as 1 of 88 children in the US [2]. A variety of genetic and environmental triggers have been proposed to be involved in causing autism [3, 4].

One focus of ASD pathophysiology involves a dysfunctional immune response, which is based in part on the controversial findings of autoantibodies in early fetal brain development or during the first few years of a child's life [5]. Likely contributing to often contradictory and conflicting autoantibody results has been the use of immunoassay methodologies measuring autoantibodies against undefined antigens such as by immunohistochemistry of brain tissue and Western blot of brain extracts using human serum [6-10]. Along these lines, a study by Singer et al found that more autistic children demonstrated increased staining intensity on Western blots corresponding to a 100 kDa band in the caudate, putamen and prefrontal cortex and for a 73 kDa band in the cerebellum and cingulate gyrus compared to controls [6]. Another group found immunoreactivity against a 52 kDa cerebellar protein as the major autoantigen species in ASD [7]. However, other studies examining autoantibodies in ASD and controls found no difference in immunoreactivity using Western blot analysis [9] and immunohistochemistry of simian brain slices [10]. Thus, the relevance, if any, of autoantibodies in ASD based on these methodologies remains unclear. Additionally, specific antigen immunoassays have been used to measure autoantibodies in ASD including autoantibodies against target proteins such as glial fibrillary acidic protein (GFAP), myelin basic protein (MBP) and glutamic acid decarboxylase-65 (GAD65) [11-15]. Singh et al. reported that ASD subjects had a higher prevalence of autoantibodies against GFAP compared to controls [14], yet another group found no association of autoantibodies against GFAP with autism [11]. Similar inconsistent reports have been reported for MBP [12, 15]. Although autoantibodies against GAD65, an enzyme responsible for producing the inhibitory transmitter gamma-aminobutyric acid, have also been reported in subjects with autism [13], these findings have yet to be reproduced by others. Together the inconsistencies of the findings highlight the need for employing more powerful immunoassay methodologies to clarify the frequency of autoantibodies in ASD.

Unlike ELISAs, fluid-phase immunoassays are generally regarded as the most sensitive and specific immunoassay format for identifying autoantibody responses and for the diagnosis of autoimmune diseases [16]. Luciferase Immunoprecipitation Systems (LIPS) is a fluid-phase immunoassay employing defined recombinant proteins expressed as an in-frame fusion with the low molecular weight light-emitting luciferase protein obtained from *Renilla reniformis*. LIPS permits robust detection of antibodies against a variety of infectious and autoimmune targets or panels of antigens from such diseases [17]. LIPS has shown high diagnostic performance for the diagnosis of autoimmune diseases that includes type I diabetes [18, 19], Sjögren's syndrome [20, 21], systemic lupus erythematosus [22], myositis [23], anti-

cytokine associated diseases [24, 25] and several autoimmune neurological diseases such as Stiff Person Syndrome [26-29]. In the current study, LIPS was used to investigate whether ASD children had previously reported autoantibodies against GAD65. In addition, antibodies were evaluated against several other autoimmune-associated autoantigens, candidate neurological autoantigens, and viral proteins. For most antigens in this study, detection of autoimmune reactivity has been previously verified in earlier reports using appropriate patient groups in which (a) the target antigen or autoantigens are known and (b) the cohort contains patients positive for the clinical diagnosis and positive for the target antibodies in their serum.

2. Methods

The cohort of children with autism and controls were evaluated at the National Institute of Mental Health, National Institutes of Health, Bethesda, MD under institutional review board approved protocol NCT00298246: Clinical and Immunological Investigations of Subtypes of Autism. The autistic children (n=104; mean age = 4.3yr; SD = 1.3; range 2.2-7.4 yrs), were diagnosed by Diagnostic and Statistical Manual of Mental Disorders, 4th ed. (DSM IV) [30] after administration of the Autism Diagnostic Observation Schedule [31] and the Autism Diagnostic Interview (ADI-R), a semi-structured parent interview concerning all domains of impairment in autism [32]. Typically developing children (n = 55) were used as a control group, (mean age = 3.3yr; SD = 1.4; range = 1.3-7.6 yrs). A control group of developmentally delayed children without autism (n=24) was also employed and was approximately matched on a developmental level to the autistic group (mean age = 4.3 yr; SD = 1.2; range = 2.7-7.6 yrs). Lastly, following antibody evaluation of the cohort, one sample was excluded from the analysis. This sample, from a subject with ASD, had unusual immunoreactivity against five of the six proteins. It appeared that the polyreactive serum recognized the *Renilla* luciferase which is the reporter enzyme in LIPS antibody test and so, the sample was excluded from consideration, leaving 103 samples in the ASD group.

Previously described LIPS tests were employed to detect autoantibodies against GAD65, Ro52, GFAP, aquaporin-4 (AQP-4), and tyrosine hydroxylase (TH) [18, 19, 22]. Three additional new antigen constructs for gamma-enolase (γ -enolase, neuron specific enolase), the gag protein from xenotropic murine leukemia virus (XMRV) and p24 from mouse mammary tumor virus (MMTV) were generated in the pREN2 vector [26] as C-terminal *Renilla* luciferase antigen fusions. γ -Enolase, was chosen as a potential target based on its high expression in neurons and reports of it being an autoantigen target in post-streptococcal autoimmune CNS disease [33]. MMTV is a potential zoonotic infection in humans [34] and XMRV infection was reported to cause chronic fatigue [35]. Both the γ -enolase and MMTV p24 were amplified by PCR using commercial cDNA plasmids. The XMRV gag protein was generated by synthetic gene synthesis (Blue Heron, Bothell, WA) using optimized human codons. The three new expression constructs were verified by sequencing and details of their construction are available upon request.

LIPS testing was performed as previously described [18, 19]. Light units were measured in a Berthold LB 960 Centro luminometer (Berthold Technologies, Germany) using coelenterazine substrate mix (Promega, Madison, WI). In some cases, control sera samples

from type I diabetes patients and systemic lupus erythematosus were used as known positive controls. Seropositivity status for GAD65, Ro52, GFAP, AQP-4 and TH was based on known cut-offs. For the three new autoantigens (γ -enolase, gag XMRV and p24 MMTV), cut-off values were assigned based on the mean plus three standard deviations of the typically developing children controls. Researchers were blinded to patient diagnosis during analysis.

The non-parametric Mann-Whitney *U* statistical test was used for comparison of antibody levels in the three different groups. For comparing the seroprevalence of antibody responses in the different groups, contingency tables were generated and analyzed using the Fischer's exact test for statistical significance.

3. Results

Based on the report that children with ASD have GAD65 autoantibodies [13], LIPS was used to measure autoantibodies against this target in a cohort with ASD. As described in the material and methods, three sera from subjects with Type I diabetes, an autoimmune disease known to generate anti-GAD65 autoantibodies, were employed as positive controls (17-19). As expected, the three type I diabetes sera showed highly elevated levels of GAD65 autoantibodies all above the established cut-off (**Fig. 1A**). However, testing of serum from the typically developed children ($n = 55$), developmentally delayed children ($n = 24$) and children with ASD ($n=103$) demonstrated no seropositive autoantibodies to GAD65: all measured LU values were below the cut-off (**Fig 1A**). To determine if the ASD children might have a lupus-like autoimmune phenotype, the LIPS Ro52 autoantibody test was employed (20-23). In contrast to three positive control samples from subjects with systemic lupus erythematosus, none of the children in the ASD cohort showed seropositivity with Ro52. All values were below the known cut-off (**Fig 1B**). It should be noted that LIPS tests for both GAD65 and Ro52 detected antibody levels in the positive control samples that were often 10-1000-fold higher than the control and ASD samples. Therefore, any subtle low positivity, if present in the cohort, should have been detected (**Fig. 1**). These findings rule out the possibility that GAD65 and Ro52 autoantibodies are biomarkers in ASD.

Autoantibodies against several other known or potential autoantigen targets including GFAP, TH, γ -enolase and AQP-4 (the autoantigen in neuromyelitis optica), were also measured by LIPS in the three groups. In certain individuals, occasional sporadic, low-level seropositive autoantibodies against the four proteins were detected. For example, two of the 55 typically developing, two of 25 developmental delayed and four of the 103 ASD children were seropositive, exhibiting low levels of autoantibodies to GFAP (**Table I**). However, statistical analysis of the GFAP seropositivity by Fischer's exact testing with contingency tables revealed no significant differences in seroprevalence between the three groups. Similar analyses for autoantibodies against AQP-4, TH, and γ -enolase also revealed occasional weak seropositivity in the children, but no significant increase in seropositivity was detected in ASD (**Table I**).

Based on the known ability of LIPS to detect robust antibodies to the capsid proteins from HIV [36] and HTL-I retroviruses [37], we explored the possibility of detecting antibodies

against two capsid proteins derived from MMTV and gag of XMRV in the cohort. Based on a cut-off derived from the healthy control children, no significant immunoreactivities were detected in any of the ASD or other children (**Table I**).

4. Discussion

In this study we tested the hypothesis that certain autoantibodies play a role in autism. First, we attempted to reproduce previous reports showing a high prevalence of GAD-65 autoantibodies in autistic children. For our investigation, a validated GAD-65 LIPS test that performs with the same high sensitivity and specificity as the gold-standard radiolabel binding immunoassay was employed [18]. Using this LIPS assay, positive control clinical serum samples were used to confirm the capacity of LIPS to detect autoantibodies against GAD65. However, no seropositive GAD65 autoantibodies were detected in any of the children from the autism cohort, which was unlike the earlier study that detected 15% seropositivity in an ASD cohort [13]. One likely explanation for this discrepancy in GAD65 prevalence in ASD is that the previous study used a solid-phase ELISA. This test format, in which target antigen proteins are bound to the ELISA plate, is known to be susceptible to non-specific reactivities [16]. Thus, it is possible that the reactivity to GAD65 in earlier ELISA-based studies was a result of such non-specific signals.

Antibodies against several other autoantigens also show no association with ASD. The lack of autoantibodies against Ro52, a commonly detectable target of autoantibodies in lupus [22], rules out a lupus-like neurological phenotype in ASD. This finding also eliminates the possibility that Ro52 is the 52 kDa target antigen seen in ASD sera by Western blotting [7]. Additionally, no evidence was found supporting a role of increased prevalence of GFAP autoantibodies in ASD, which is in agreement with a previous study [11]. Specifically, 1% and 2%, of the ASD and control groups, respectively, showed GFAP seropositivity, but the absolute levels were low. Thus, both a neuronal and a glial autoantibody either were not present or their prevalence was not elevated in ASD subjects. No differences between the three groups were found for the prevalence of autoantibodies against AQP-4, tyrosine hydroxylase or neuron-specific γ -enolase. The lack of autoantibodies against AQP-4, an astrocytic protein involved with maintenance of the blood brain barrier and a pathognomonic autoantibody in neuromyelitis optica rules out this target as a potential source of autoimmune blood brain barrier disruption in ASD. Moreover, seropositivity to the neuron-specific γ -enolase was found in 1% and 2%, of the ASD and control groups, respectively, but again the absolute levels were low. Also, no difference was seen between the three groups for TH autoantibodies. Thus, two more neuronal autoantibody targets either were not present or their prevalence was not elevated in ASD subjects. TH is particularly relevant since it is the first enzyme in the biosynthetic pathway for the catecholamines dopamine and norepinephrine, which are major neurotransmitters in circuits subserving affect and attention. Similar to our GFAP observations there are occasional sporadic low level signals to several of these targets that likely reflect uncharacterized immune responses.

Two possible viral targets, MMTV and XMRV, also showed no significant immunoreactivity in the cohort. The fact that we did not detect antibodies against XMRV in autism using LIPS is also in agreement with two other reports [38, 39], which used mainly

polymerase chain reaction molecular methods for detection. Thus, three independent reports using different methodologies do not support detection of the XMRV artifact in ASD. Despite no evidence for a role of XMRV or MMTV, additional studies are needed to determine if other infectious agents, or the body's response to such infectious agents, might play a role.

From the results presented here, we conclude that autoantibodies against GAD65 and the other targets that we tested are unlikely to be significant contributors to autism spectrum disorder. However, given the limited number of antigens examined, it is possible there are other neurological antigens and target proteins which might be targets of humoral responses in ASD. It would be worthwhile to explore the immunoreactivity reported in ASD by several groups that employed Western blot [7] and immunohistochemical techniques [7, 40] by using high sensitivity LIPS assays. These investigations would first require that the antigens be identified, in order that defined recombinant proteins might be used to construct quantitative immunoassays for analysis of a large number of controls and ASD subjects. The presence of an autoimmune component in ASD is an important question to resolve in a definitive fashion. The data presented in this report suggest that newer more specific and defined methodologies can make a substantial contribution to realizing this goal.

5. Conclusions

This study employed a fluid-phase immunoassay, LIPS, which shows high diagnostic performance for autoantibodies present in multiple autoimmune disorders and for infectious agents. Using this assay, we found no evidence of autoantibodies against GAD65 and several other autoantigens in children with autism spectrum disorder. No antibodies were detected against two retroviruses, MMTV or XMRV. These results suggest that autoimmunity against the tested neuronal, astrocytic, and other proteins are not likely to be causative factors or diagnostic biomarkers in autism.

Acknowledgments

This work was supported by the intramural research programs of the Clinical Center, National Institutes of Health, National Institute of Dental and Craniofacial Research, and the National Institute of Mental Health protocol 06-M-0102; NCT00298246. We thank current and former members of the Pediatrics and Developmental Neuroscience Branch, and extend sincere gratitude to the children and families who participated in this research.

References

1. Association A. P.. Diagnostic and statistical manual of mental disorders : DSM-5. 5th edn.. American Psychiatric Association; Washington, D.C.: 2013.
2. Investigators A. a. D. D. M. N. S. Y. P., Prevention C. f. D. C. a. Prevalence of autism spectrum disorders--Autism and Developmental Disabilities Monitoring Network, 14 sites, United States, 2008. *MMWR Surveill. Summ.* 2012; 61:1–19.
3. Devlin B, Scherer SW. Genetic architecture in autism spectrum disorder. *Curr. Opin. Genet. Dev.* 2012; 22:229–237. [PubMed: 22463983]
4. Lai MC, Lombardo MV, Baron-Cohen S. Autism. *Lancet.* 2014; 383:896–910. [PubMed: 24074734]
5. Gesundheit B, Rosenzweig JP, Naor D, Lerer B, Zachor DA, Prochazka V, Melamed M, Kristt DA, Steinberg A, Shulman C, et al. Immunological and autoimmune considerations of Autism Spectrum Disorders. *J. Autoimmun.* 2013; 44:1–7. [PubMed: 23867105]

6. Singer HS, Morris CM, Williams PN, Yoon DY, Hong JJ, Zimmerman AW. Antibrain antibodies in children with autism and their unaffected siblings. *J. Neuroimmunol.* 2006; 178:149–155. [PubMed: 16842863]
7. Wills S, Cabanlit M, Bennett J, Ashwood P, Amaral DG, Van de Water J. Detection of autoantibodies to neural cells of the cerebellum in the plasma of subjects with autism spectrum disorders. *Brain Behav. Immun.* 2009; 23:64–74. [PubMed: 18706993]
8. Mazur-Kolecka B, Cohen IL, Gonzalez M, Jenkins EC, Kaczmarek W, Brown WT, Flory M, Frackowiak J. Autoantibodies against neuronal progenitors in sera from children with autism. *Brain Dev.* 2014; 36:322–329. [PubMed: 23838310]
9. Morris CM, Zimmerman AW, Singer HS. Childhood serum anti-fetal brain antibodies do not predict autism. *Pediatr. Neurol.* 2009; 41:288–290. [PubMed: 19748049]
10. Rossi CC, Van de Water J, Rogers SJ, Amaral DG. Detection of plasma autoantibodies to brain tissue in young children with and without autism spectrum disorders. *Brain Behav. Immun.* 2011; 25:1123–1135. [PubMed: 21420487]
11. Kirkman NJ, Libbey JE, Sweeten TL, Coon HH, Miller JN, Stevenson EK, Lainhart JE, McMahon WM, Fujinami RS. How relevant are GFAP autoantibodies in autism and Tourette Syndrome? *J. Autism Dev. Disord.* 2008; 38:333–341. [PubMed: 17578659]
12. Libbey JE, Coon HH, Kirkman NJ, Sweeten TL, Miller JN, Stevenson EK, Lainhart JE, McMahon WM, Fujinami RS. Are there enhanced MBP autoantibodies in autism? *J. Autism Dev. Disord.* 2008; 38:324–332. [PubMed: 17588145]
13. Rout UK, Mungan NK, Dhossche DM. Presence of GAD65 autoantibodies in the serum of children with autism or ADHD. *Eur. Child Adolesc. Psychiatry.* 2012; 21:141–147. [PubMed: 22323074]
14. Singh VK, Warren R, Averett R, Ghaziuddin M. Circulating autoantibodies to neuronal and glial filament proteins in autism. *Pediatr. Neurol.* 1997; 17:88–90. [PubMed: 9308986]
15. Singh VK, Warren RP, Odell JD, Warren WL, Cole P. Antibodies to myelin basic protein in children with autistic behavior. *Brain Behav. Immun.* 1993; 7:97–103. [PubMed: 7682457]
16. Liu E, Eisenbarth GS. Accepting clocks that tell time poorly: fluid-phase versus standard ELISA autoantibody assays. *Clin. Immunol.* 2007; 125:120–126. [PubMed: 17904423]
17. Burbelo PD, Lebovitz EE, Notkins AL. Luciferase immunoprecipitation systems for measuring antibodies in autoimmune and infectious diseases. *Transl. Res.* 2015; 165:325–335. [PubMed: 25241936]
18. Burbelo PD, Hirai H, Issa AT, Kingman A, Lernmark A, Ivarsson SA, Notkins AL, Iadarola MJ. Comparison of radioimmunoprecipitation with luciferase immunoprecipitation for autoantibodies to GAD65 and IA-2beta. *Diabetes Care.* 2010; 33:754–756. [PubMed: 20086252]
19. Burbelo PD, Lebovitz EE, Bren KE, Bayat A, Paviol S, Wenzlau JM, Barriga KJ, Rewers M, Harlan DM, Iadarola MJ. Extraprostatic autoantibody profiles in type I diabetes. *PLoS One.* 2012; 7:e45216. [PubMed: 23028856]
20. Ching KH, Burbelo PD, Gonzalez-Begne M, Roberts ME, Coca A, Sanz I, Iadarola MJ. Salivary anti-Ro60 and anti-Ro52 antibody profiles to diagnose Sjogren's Syndrome. *J. Dent. Res.* 2011; 90:445–449. [PubMed: 21212317]
21. Burbelo PD, Leahy HP, Issa AT, Groot S, Baraniuk JN, Nikolov NP, Illei GG, Iadarola MJ. Sensitive and robust luminescent profiling of anti-La and other autoantibodies in Sjogren's syndrome. *Autoimmunity.* 2009; 42:515–524. [PubMed: 19657778]
22. Ching KH, Burbelo PD, Tipton C, Wei C, Petri M, Sanz I, Iadarola MJ. Two major autoantibody clusters in systemic lupus erythematosus. *PLoS One.* 2012; 7:e32001. [PubMed: 22363785]
23. Gan L, O'Hanlon TP, Gordon AS, Rider LG, Miller FW, Burbelo PD. Twins discordant for myositis and systemic lupus erythematosus show markedly enriched autoantibodies in the affected twin supporting environmental influences in pathogenesis. *BMC Musculoskelet. Disord.* 2014; 15:67. [PubMed: 24602337]
24. Browne SK, Burbelo PD, Chetchotisakd P, Suputtamongkol Y, Kiertiburanakul S, Shaw PA, Kirk JL, Jutivorakool K, Zaman R, Ding L, et al. Adult-onset immunodeficiency in Thailand and Taiwan. *N. Engl. J. Med.* 2012; 367:725–734. [PubMed: 22913682]

25. Burbelo PD, Browne SK, Sampaio EP, Giaccone G, Zaman R, Kristosturyan E, Rajan A, Ding L, Ching KH, Berman A, et al. Anti-cytokine autoantibodies are associated with opportunistic infection in patients with thymic neoplasia. *Blood*. 2010; 116:4848–4858. [PubMed: 20716769]
26. Burbelo PD, Groot S, Dalakas MC, Iadarola MJ. High definition profiling of autoantibodies to glutamic acid decarboxylases GAD65/GAD67 in stiff-person syndrome. *Biochem. Biophys. Res. Commun.* 2008; 366:1–7. [PubMed: 18047830]
27. Ching KH, Burbelo PD, Kimball RM, Clawson LL, Corse AM, Iadarola MJ. Recombinant expression of the AChR- α 1 subunit for the detection of conformation-dependent epitopes in Myasthenia Gravis. *Neuromuscul. Disord.* 2011; 21:204–213. [PubMed: 21195619]
28. Nakane S, Higuchi O, Koga M, Kanda T, Murata K, Suzuki T, Kurono H, Kunimoto M, Kaida K, Mukaino A, et al. Clinical features of autoimmune autonomic ganglionopathy and the detection of subunit-specific autoantibodies to the ganglionic acetylcholine receptor in Japanese patients. *PLoS One*. 2015; 10:e0118312. [PubMed: 25790156]
29. Higuchi O, Hamuro J, Motomura M, Yamanashi Y. Autoantibodies to low-density lipoprotein receptor-related protein 4 in myasthenia gravis. *Ann. Neurol.* 2011; 69:418–422. [PubMed: 21387385]
30. Association A. P.. *Diagnostic and statistical manual of mental disorders : DSM-IV*. 4th edn.. American Psychiatric Association; Washington, DC: 1994.
31. Lord, C.; Rutter, M.; DiLavore, PC.; Risi, S. *Autism Diagnostic Observation Schedule (ADOS)*. Western Psychological Services; Los Angeles, California: 1999.
32. Lord C, Rutter M, Le Couteur A. *Autism Diagnostic Interview-Revised: a revised version of a diagnostic interview for caregivers of individuals with possible pervasive developmental disorders*. *J Autism Dev Disord.* 1994; 24:659–685. [PubMed: 7814313]
33. Dale RC, Candler PM, Church AJ, Wait R, Pocock JM, Giovannoni G. Neuronal surface glycolytic enzymes are autoantigen targets in post-streptococcal autoimmune CNS disease. *J. Neuroimmunol.* 2006; 172:187–197. [PubMed: 16356555]
34. Voisset C, Weiss RA, Griffiths DJ. Human RNA “rumor” viruses: the search for novel human retroviruses in chronic disease. *Microbiol. Mol. Biol. Rev.* 2008; 72:157–196. table of contents. [PubMed: 18322038]
35. Lombardi VC, Ruscetti FW, Das Gupta J, Pfof MA, Hagen KS, Peterson DL, Ruscetti SK, Bagni RK, Petrow-Sadowski C, Gold B, et al. Detection of an infectious retrovirus, XMRV, in blood cells of patients with chronic fatigue syndrome. *Science*. 2009; 326:585–589. [PubMed: 19815723]
36. Burbelo PD, Bayat A, Rhodes CS, Hoh R, Martin JN, Fromentin R, Chomont N, Hutter G, Kovacs JA, Deeks SG. HIV antibody characterization as a method to quantify reservoir size during curative interventions. *J. Infect. Dis.* 2014; 209:1613–1617. [PubMed: 24286982]
37. Burbelo PD, Meoli E, Leahy HP, Graham J, Yao K, Oh U, Janik JE, Mahieux R, Kashanchi F, Iadarola MJ, et al. Anti-HTLV antibody profiling reveals an antibody signature for HTLV-I-associated myelopathy/tropical spastic paraparesis (HAM/TSP). *Retrovirology*. 2008; 5:96. [PubMed: 18937847]
38. Satterfield BC, Garcia RA, Gurrieri F, Schwartz CE. PCR and serology find no association between xenotropic murine leukemia virus-related virus (XMRV) and autism. *Mol. Autism*. 2010; 1:14. [PubMed: 20946639]
39. Lintas C, Guidi F, Manzi B, Mancini A, Curatolo P, Persico AM. Lack of infection with XMRV or other MLV-related viruses in blood, post-mortem brains and paternal gametes of autistic individuals. *PLoS One*. 2011; 6:e16609. [PubMed: 21373179]
40. Wills S, Rossi CC, Bennett J, Martinez Cerdeno V, Ashwood P, Amaral DG, Van de Water J. Further characterization of autoantibodies to GABAergic neurons in the central nervous system produced by a subset of children with autism. *Mol. Autism*. 2011; 2:5. [PubMed: 21521495]




Figure 1. Lack of autoantibodies against GAD65 and Ro52 in the ASD cohort

Autoantibody levels were evaluated for (A) GAD65 and (B) Ro52 in the cohort. Additional positive control serum samples from type I diabetes and systemic lupus erythematosus subjects were included for the detection of GAD65 and Ro52 autoantibodies, respectively. The light units (LU) plotted on the y-axis using a log₁₀ scale measure the autoantibody levels for each sample. The dotted lines represent the cut-off values for determining seropositivity for each autoantigen.

Table 1

Prevalence of antibodies in the autism cohort

Antigen	Typically Developing (N=55) Seropositivity	Developmental Delay (N=25) Seropositivity	ASD (N=103) Seropositivity *
GAD-65	0/55	0/25	0/103
Ro52	0/55	0/25	0/103
GFAP	2/55	2/25	4/103
AQP-4	3/55	0/25	5/103
TH	1/55	0/25	1/103
γ - enolase	1/55	0/25	2/103
XMRV gag	1/55	0/25	0/103
MMTV p24	1/55	0/25	0/103

* Fischer exact tests revealed no statistical difference in prevalence of antibodies against the eight targets in ASD compared to typically developing and developmental delay children.