



Published in final edited form as:

Clin Transl Sci. 2015 December ; 8(6): 740–745. doi:10.1111/cts.12346.

Unique DNA methylation patterns in offspring of hypertensive pregnancy

Colleen G. Julian¹, Brent S. Pedersen¹, Carlos Salinas Salmon², Ivana V. Yang¹, Marcelino Gonzales², Enrique Vargas², Lorna G. Moore³, and David A. Schwartz¹

¹Department of Medicine, University of Colorado School of Medicine, Aurora, Colorado, USA

²Bolivian Institute of High Altitude Biology, La Paz, Bolivia

³Department of Obstetrics and Gynecology, University of Colorado School of Medicine, Aurora, Colorado, USA 80045

Abstract

Epigenomic processes are believed to play a pivotal role for the effect of environmental exposures in early life to modify disease risk throughout the lifespan. Offspring of women with hypertensive complications of pregnancy (HTN_{PREG}) have an increased risk of developing systemic and pulmonary vascular dysfunction in adulthood. In this preliminary report, we sought to determine whether epigenetic modifications of genes involved in the regulation of vascular function were present in HTN_{PREG} offspring. We contrasted DNA methylation and gene expression patterns of peripheral blood mononuclear cells obtained from young male offspring of HTN_{PREG} (n=5) to those of normotensive controls (n=19). In HTN_{PREG} offspring we identified six differentially methylated regions (DMRs) including three genes (*SMOC2*, *ARID1B* and *CTRHC1*) relevant to vascular function. The transcriptional activity of *ARID1B* and *CTRHC1* was inversely related to methylation status. HTN_{PREG} offspring had higher systolic pulmonary artery pressure (sP_{PA}) vs. controls. Our findings demonstrate that epigenetic marks are altered in offspring of HTN_{PREG} with a modest elevation of sP_{PA} and introduce novel epigenomic targets for further study. On the basis of these findings we speculate that epigenomic mechanisms may be involved in mediating the effect of HTN_{PREG} to raise the risk of vascular disease later in life.

Keywords

epigenetics; preeclampsia; gestational hypertension; developmental programming

Introduction

Intrauterine exposures influence physiological function and disease susceptibility throughout the lifespan, as demonstrated by epidemiologic and experimental animal studies conducted over the past two decades. For instance, maternal hypertensive disorders of pregnancy

Corresponding Address: Colleen Glyde Julian, PhD, Department of Medicine, University of Colorado Denver, 12700 E 19th Avenue, Mailstop 8611, 3rd Floor Research Complex 2, Aurora, CO 80045, colleen.julian@ucdenver.edu, Phone: 303-724-6443, Fax: 303-724-1799.

(HTN_{PREG}) predispose offspring to systemic cardiovascular disease (1, 2) and pulmonary vascular dysfunction (3) in adulthood. Because epigenetic marks are responsive to environmental stimuli (4–6) and mediate gene expression patterns that control embryonic development and organogenesis (7), epigenomic processes are believed to play a central role in the developmental programming of adult disease. In support of this view, DNA methylation changes induced by intrauterine exposures can modify gene expression and physiological function in ways that are of clinical importance (4, 6, 8).

Locus-specific DNA methylation marks, some within genes known to influence vascular development and integrity, have been identified in placental tissue or cord blood derived from HTN_{PREG} (e.g., (9, 10)). Therefore, it is logical to speculate that epigenomic mechanisms may mediate the increased risk of vascular dysfunction in offspring of HTN_{PREG}. To our knowledge no study has queried genome-wide DNA methylation patterns in young adult offspring of HTN_{PREG} to identify the potential role of epigenetics for compromised vascular function in these individuals. Accordingly, in this preliminary report we sought to identify differentially methylated genomic regions (DMRs) in offspring of HTN_{PREG} vs. offspring of normotensive controls.

Understanding the role of epigenetics in mediating the effects of intrauterine exposures to modify long-term disease susceptibility has important clinical applications with respect to the identification of environmental triggers for adult-onset disease and novel therapeutic targets. DNA methylation is a particularly attractive candidate for therapeutic intervention because methylation status is amenable to pharmacologic modification.

Materials and Methods

Study Design

Subjects and sample sizes—Subjects included 24 Andean males between the ages of 18 and 25 living in La Paz-El Alto, Bolivia (3600 – 4100 m). Of these, five were born to women whose pregnancy was complicated by hypertension (HTN_{PREG}), and 19 were born to women having a normotensive pregnancy (controls). Additional exclusion criteria were a history of cardiac or pulmonary disease, evidence of cardiopulmonary disease on clinical exam, chronic bronchitis or frequent smoking (> 4 cigarettes per week). Conducting the overarching study in Bolivia was beneficial because more than 1.5 million persons live in the La Paz-El Alto metropolitan area which is situated at a sufficient altitude to increase the incidence of hypoxia-related pregnancy complications, including HTN_{PREG} (11, 12). An additional advantage of the high-altitude model is that because chronic hypoxia exposes predisposition to pulmonary vascular dysfunction (4) high altitude would be expected to improve the detection of elevated sP_{PA} and its association with methylation status. Physiologic studies and sample collection were performed at the Bolivian Institute of High Altitude Biology (3600 m) in La Paz, Bolivia. On the first visit peripheral blood samples were collected from an antecubital vein by routine venipuncture and the subject participated in a detailed questionnaire to obtain information regarding residential and health history. Cardiopulmonary evaluations were performed on subsequent visits. Medical record reviews and an in-depth interview with the subjects' mother were conducted to obtain information regarding pregnancy, delivery, and perinatal characteristics. The Colorado Multiple

Institutional Review Board and Colegio Medico, its Bolivian counterpart, approved all survey and study procedures.

Systolic Pulmonary Artery Pressure (sP_{PA})—Transthoracic echocardiography (GE Vivid I Ultrasound 2007) was used to estimate sP_{PA} as previously described (13).

Quantification of tricuspid regurgitation was measured with the use of continuous wave Doppler and averaged from three complete waveforms of good quality. Maximal tricuspid regurgitation jet flow velocity (TR_{max}) was used to estimate sP_{PA} using a modified Bernoulli equation as previously reported (3).

Medical Records Review and Maternal Interview—Perinatal data were obtained by structured interview of the subject's mother and a review of medical records from the hospital or clinic where the subject was born. Data collected were: a) maternal health and reproductive history, b) pregnancy or delivery complications, c) fetal complications, and d) newborn characteristics and complications. Hypertensive pregnancy was determined by a diagnosis of preeclampsia or gestational hypertension as noted in the medical records or as obtained from maternal interview. Specific questions regarding HTN_{PREG} included, "Were you told you had preeclampsia or gestational hypertension during your pregnancy?", "Were you told you had high blood pressure multiple times during your pregnancy?" and, to distinguish between preexisting hypertension and HTN_{PREG}, "Did you have high blood pressure before your pregnancy?". Gestational hypertension and preeclampsia were grouped into a single HTN_{PREG} category for this preliminary test since both conditions impair fetal growth and influence vascular function in adulthood (2, 14).

Differences between HTN_{PREG} and controls with respect to demographic variables and physiologic characteristics were determined using independent T-tests or chi-squared tests as appropriate. Data are reported as the mean \pm SEM or the proportion and 95% CI.

Sampling and Isolation of Genomic DNA (gDNA) and Total RNA—Peripheral blood samples were collected from an antecubital vein using standard phlebotomy, placed into a BD Vacutainer Cell Preparation Tube containing sodium citrate and Ficoll Hypaque density fluid. PBMCs were isolated according to manufacturer guidelines, resuspended in RNeasy Lysis Buffer (Qiagen) and stored at -80°C until analysis. gDNA and RNA were isolated using the Qiagen AllPrep DNA/RNA Mini Kit. gDNA was purified using the Qiagen QIAEX II Kit and quantitated with the NanoDrop ND-100 UV-Vis spectrophotometer (NanoDrop Technologies).

DNA Methylation Arrays, Analysis, and Interpretation—DNA methylation was assessed using Illumina's Infinium HumanMethylation450 BeadChip arrays. This methylation platform permits the interrogation of >485k methylation sites at single-nucleotide resolution and has been validated with respect to technical performance and functional relevance. Methylated cytosines were bisulfite converted using the EZ DNA Methylation Gold Kit (Zymo Research), amplified by WGA and then enzymatically fragmented. 4 μl of bisulfite converted DNA at 50ng/ μl was hybridized to the BeadChip. Quantitative methylation scores were generated for each of the 485k loci for each sample using Illumina's GenomeStudio Methylation module. Signal intensities of methylated and

unmethylated probes were exported from the GenomeStudio interface, along with detection p-values representing the likelihood of detection relative to background. Individual data points with a p-value outside of detection criteria ($p > 0.05$) were treated as missing data. Data were normalized using the SWAN method (15) within the R package minfi, and M-values were calculated. Methylation differences between HTN_{PREG} and controls were identified using limma (16) and multcomp (17) packages in R. From the resultant p-values, differentially methylated regions (DMRs) between HTN_{PREG} and controls were identified using Comb-p, an analytical tool developed to analyze spatially correlated p-values (18). To define DMRs, Comb-p uses a sliding window approach to identify genomic regions with multiple low p-values after accounting for local dependencies and multiple testing using a false discovery rate cut off for multiple comparisons. DMRs were ranked by p-value, and annotated to the nearest gene. Significance was defined as an $\alpha < 5\%$ after correcting for the number of possible DMRs. Because methylation marks can influence the transcriptional activity of adjacent genes, we inspected genes located within 1Mb upstream or downstream of intergenic DMRs for additional vascular-related candidates.

Compared to associations based on the differential methylation of single CpG sites, DMR analysis has greater specificity and power to detect functionally relevant methylation-phenotype associations (19). For instance, evidence indicates that methylation status may be profoundly affected by sequence content and genetic variation, including single nucleotide polymorphisms (20, 21). Because DMRs require multiple adjacent probes to be differentially methylated, the Comb-p method reduces the influence of sequence-specific bias. Prior investigations have used similar methods to identify clinically relevant methylation-phenotype associations (22).

Influence of DMRs on Gene Expression—RT-PCR was used to determine the influence of the DMRs identified on gene expression. cDNA was generated using random primers with Invitrogen's Superscript III First-Strand Synthesis System SuperMix. PCR reactions were prepared using TaqMan Fast Advanced Master Mix and TaqMan Gene Expression Assays (Applied Biosystems) as specified by the manufacturer. RT-PCR was performed and analyzed on the ViiA 7 Real-Time PCR System with ViiA 7 1.22 software (Applied Biosystems). The quantification of target gene transcription relative to that of GAPDH was assessed using the 2^{-CT} method. Unpaired Student's t tests were used to identify differential expression between HTN_{PREG} and controls using a significance threshold of $P < 0.05$. Data are expressed as means \pm SD.

Results

HTN_{PREG} and controls were equivalent with respect to age, height, weight, resting heart rate and S_aO_2 . When compared to controls, HTN_{PREG} had higher estimated sP_{PA} ($p = 0.007$) (Table 1).

Comparing HTN_{PREG} with control subjects, we identified six genome-wide significant DMRs (Figure 1). Each DMR was associated with a unique gene: namely, collagen triple helix repeat containing 1 (*CTHRC1*), tripartite motif containing 31 (*TRIM31*), AT rich interactive domain 1B (*ARID1B*), SPARC related modular calcium binding 2 (*SMOC2*),

leucine-rich repeats and IQ motif containing 3 (*LRRIQ3*) or long intergenic non-protein coding RNA 226 (*LINC00226*).

Of the six DMRs, three were hypomethylated and three were hypermethylated in HTN_{PREG} vs. controls. In each case, the directionality of the differentially methylated sites composing the DMR was consistent. Methylation differences between HTN_{PREG} and controls at each DMR averaged 6.7% (range, 2.6 to 11.2%; Table 2). Three DMRs contained differentially methylated sites within gene bodies (intron, exon, 3' untranslated region [UTR] and 5' UTR), and one DMR (*CTHRC1*) showed differential methylation within 200–1,500 bases from a transcriptional start site (promoter) (Table 2). Several genes with potential relevance for vascular function, including metastasis associated 1 (*MTA1*) and cytosine-rich protein2 (*CRIP2*), and thrombospondin 2 (*THBS2*) are located within 1Mb upstream or downstream of the intergenic *LINC00226* and *SMOC2* DMRs, respectively. No vascular-related candidate genes were identified within the 1Mb band on either side of the *ARID1B* DMR.

We then evaluated the association between DMRs and the expression of genes with a biologically plausible relationship to vascular function (*ARID1B*, *CTHRC1* and *SMOC2*). Compared to controls *ARID1B* gene expression was impaired in HTN_{PREG} offspring (P=0.025), while *CTHRC1* tended to be higher (P=0.08) and *SMOC2* was equivalent between groups (P=NS). Notably, *ARID1B* and *CTHRC1* expression were inversely related to methylation status. Based on existing literature, the remaining DMRs either do not have a clear relationship to vascular function (*TRIM31*) or have not been well described (*LRRIQ3* and *LINC00226*) and were therefore not considered for expression studies.

Discussion

Epigenomic processes are believed to play a pivotal role for mediating the effect of environmental exposures in early life on disease risk later in life. Numerous studies have shown that HTN_{PREG} is associated with systemic or pulmonary vascular abnormalities in adulthood (1–3). Here we identified locus-specific DMRs in offspring of HTN_{PREG} with elevated sP_{PA}. Two of the DMRs identified in vascular-related genes, *ARID1B* and *CTHRC1*, were also differentially expressed, suggesting the potential functional relevance of the epigenetic modifications identified. Our findings support the hypothesis that epigenetic mechanisms may be involved in mediating the effect of HTN_{PREG} to increase transgenerational vascular disease risk and introduce novel epigenomic targets for further experimental study.

ARID1B is the largest component of the switching defective/sucrose non-fermenting (SWI/SNF) ATP-dependent chromatin-modeling complex. Because SWI/SNF chromatin-remodeling complexes interact with numerous transcription factors to regulate gene expression and are required for the hypoxic induction of several genes including erythropoietin and vascular endothelial growth factor (VEGF) (e.g., (23)), we consider that this gene may be of particular relevance for the vascular dysfunction observed in offspring of HTN_{PREG}. Indeed, the SWI/SNF complex is essential for cellular proliferation and differentiation in multiple cardiac cell lines, cardiomyocyte development, vasculogenesis and vascular smooth muscle proliferation (reviewed in (24)). For instance, components of

the SWI/SNF complex (Brg1/Brm) contribute to the pathogenesis of endothelial dysfunction and atherosclerosis via the transcriptional regulation of inflammation-induced cell adhesion molecules (25). Consistent with the inverse methylation-expression relationship observed here, *ARID1B* is subject to methylation-induced transcriptional silencing. Specifically, hypermethylation upstream of the *ARID1B* initiation codon impairs ARID1B expression in pancreatic cancer cells (26).

Two of the DMRs identified in HTN_{PREG} offspring, *CTHRC1* and *SMOC2*, are important for vascular remodeling and angiogenesis in the context of vascular injury. Specifically, *CTHRC1* is transiently overexpressed by adventitial fibroblasts and smooth muscle of the neointima in the carotid artery after balloon-induced injury (27). *CTHRC1* overexpression in fibroblasts and smooth muscle cells enhances migratory ability, reduces collagen matrix deposition, and temporally coincides with extensive adventitial remodeling, suggesting its potential involvement in the regulation of vascular remodeling (27). The expression of *SMOC2*, a novel member of the SPARC family of matricellular proteins that regulate interactions between cells and the extracellular matrix, is upregulated in response to vascular injury (28). *SMOC2* overexpression in HUVECS also reduces the expression of thrombospondin 1 (*THBS1*), an angiogenic inhibitor, and enhances mitogenesis and cellular proliferation in the presence of angiogenic growth factors including vascular endothelial growth factor (VEGF) (29). *THBS2* neighbors the *SMOC2* intergenic DMR and, similar to *THBS1*, has potent antiangiogenic actions by antagonizing VEGF activity and via its direct influence on endothelial cell proliferation, survival and migration [reviewed in (30)]. This raises the possibility that the intergenic *SMOC2* DMR identified may be acting at a distance on other genes important for vascular function. Further supporting the angiogenic potential of *SMOC2* and its potential biological importance for vascular disease, *in vitro* Matrigel plug invasion assays in which HUVECS were transduced with Ad-*SMOC2* showed more projections and greater cell-network area compared to Ad-GFP transduced cells (29).

Although *LINC00226* itself is not well-described several genes within 1Mb of the intergenic *LINC00226* are involved in the regulation of vascular function. *CRIP2* is a cytoskeletal protein that influences the phenotypic modulation of vascular smooth muscle cells (VSMCs) in response to vascular injury (31). For instance, *CRIP2* prevents excessive VSMC migration and proliferation that are associated with increased myocardial infarction risk (31). *MTA1* is associated with increased metastatic potential and angiogenesis in numerous cancers, including malignant esophageal, lung, breast and prostate tumors, but also acts as a proangiogenic agent in normal, healthy tissue due, in part, to its ability to upregulate the expression of VEGF and its membrane-bound receptor (Flt-1) [reviewed in (32)].

There are several limitations to our study. A recurrent challenge in epigenomics is the inability to distinguish whether differential methylation patterns are a cause or consequence of the phenotype of interest, or whether their effect is neutral. Additionally, the DMRs we identified may have been influenced by other exposures occurring during development or inherent intra-individual differences. Despite this challenge, our findings offspring provide novel insight with respect to the potential importance of epigenomic dysregulation in individuals exposed to maternal hypertension during gestation and can be used to guide future research in this area. Similarly, the elevated sP_{PA} we observed in HTN_{PREG} offspring

is genetic in nature or may result from developmental exposures. In support of the latter possibility, offspring born to preeclamptic mothers have impaired pulmonary and systemic vessel function whereas their siblings born after a normotensive pregnancy do not (3). Another consideration is that we relied primarily on maternal recall of HTN_{PREG} or a clinical diagnosis as noted in the medical record, and lacked repeated blood pressure and proteinuria measurements. Maternal recall of HTN_{PREG} has recently been validated with respect to its concordance with medical records, and the ability to accurately discriminate between gestational hypertension and preeclampsia (33). It is also worth noting that we used PBMCs as a proxy to explore biological processes occurring in or influencing the vasculature. There are several advantages to the use of PBMCs. First, PBMCs are the most transcriptionally active cells in the blood and their gene expression patterns influence systemic processes including the regulation of vascular function. Second, because PBMCs can be obtained using minimally invasive methods the potential for their use in the clinic or for prospective epigenomic studies beginning in early life is high. Finally, epigenomic profiles of PBMCs have successfully identified pathological processes important for various diseases such as asthma (22).

In summary, our observations raise the possibility that intrauterine exposure to HTN_{PREG} modifies epigenetic marks in ways that may be of pathological importance for the development of vascular dysfunction in later life. While further human and experimental animal studies are required to evaluate the role of epigenetics for the effect of HTN_{PREG} to modify transgenerational vascular disease risk, the present study has identified several targets meriting further research.

Acknowledgments

We would like to acknowledge all the individuals who participated in this project as research subjects and the physicians and technical staff at the Bolivian Institute of High Altitude Biology. Support for data collection and analysis was obtained from grants NIH-HL079647 (L.G.M.), R03TW007957 from the Fogarty International Center (L.G.M. and C.G.J.), NIH-HL095393 (D.A.S.), NIH-HL101715, (D.A.S.), NIH-ES18181 (D.A.S.) and the NIH/NCATS Colorado CTSA UL1 TR001082 (CMH-Pilot, C.G.J.). Dr. Julian is supported by the NIH BIRCWH program (5 K12HD057022-07).

References

1. Kajantie E, Eriksson JG, Osmond C, Thornburg K, Barker DJ. Pre-eclampsia is associated with increased risk of stroke in the adult offspring: the Helsinki birth cohort study. *Stroke*. 2009; 40(4): 1176–80. [PubMed: 19265049]
2. Geelhoed JJ, Fraser A, Tilling K, Benfield L, Davey Smith G, Sattar N, Nelson SM, Lawlor DA. Preeclampsia and gestational hypertension are associated with childhood blood pressure independently of family adiposity measures: the Avon Longitudinal Study of Parents and Children. *Circulation*. 2010; 122(12):1192–9. [PubMed: 20823385]
3. Jayet PY, Rimoldi SF, Stuber T, Salmon CS, Hutter D, Rexhaj E, Thalmann S, Schwab M, Turini P, Sartori-Cucchia C, Nicod P, Villena M, Allemann Y, Scherrer U, Sartori C. Pulmonary and systemic vascular dysfunction in young offspring of mothers with preeclampsia. *Circulation*. 2010; 122(5):488–94. [PubMed: 20644018]
4. Rexhaj E, Bloch J, Jayet PY, Rimoldi SF, Dessen P, Mathieu C, Tolsa JF, Nicod P, Scherrer U, Sartori C. Fetal programming of pulmonary vascular dysfunction in mice: role of epigenetic mechanisms. *American journal of physiology Heart and circulatory physiology*. 2011; 301(1):H247–52. [PubMed: 21536851]

5. Watson JA, Watson CJ, McCann A, Baugh J. Epigenetics, the epicenter of the hypoxic response. *Epigenetics : official journal of the DNA Methylation Society*. 2010; 5(4):293–6.
6. Hollingsworth JW, Maruoka S, Boon K, Garantziotis S, Li Z, Tomfohr J, Bailey N, Potts EN, Whitehead G, Brass DM, Schwartz DA. In utero supplementation with methyl donors enhances allergic airway disease in mice. *J Clin Invest*. 2008; 118(10):3462–9. [PubMed: 18802477]
7. Li E. Chromatin modification and epigenetic reprogramming in mammalian development. *Nature reviews Genetics*. 2002; 3(9):662–73.
8. Patterson AJ, Xiao D, Xiong F, Dixon B, Zhang L. Hypoxia-derived oxidative stress mediates epigenetic repression of PKCepsilon gene in foetal rat hearts. *Cardiovasc Res*. 2012; 93(2):302–10. [PubMed: 22139554]
9. Blair JD, Yuen RK, Lim BK, McFadden DE, von Dadelszen P, Robinson WP. Widespread DNA hypomethylation at gene enhancer regions in placentas associated with early-onset pre-eclampsia. *Mol Hum Reprod*. 2013; 19(10):697–708. [PubMed: 23770704]
10. Ching T, Ha J, Song MA, Tiirikainen M, Molnar J, Berry MJ, Towner D, Garmire LX. Genome-scale hypomethylation in the cord blood DNAs associated with early onset preeclampsia. *Clin Epigenetics*. 2015; 7(1):21. [PubMed: 25806090]
11. Julian CG, Vargas E, Armaza JF, Wilson MJ, Niermeyer S, Villena M, Moore LG. High-altitude ancestry protects against hypoxia-associated reductions in fetal growth. *Arch Dis Child Fetal Neonatal Ed*. 2007; 92(5):F372–7. [PubMed: 17329275]
12. Keyes LE, Armaza JF, Niermeyer S, Vargas E, Young D, Villena M, Moore LG. Intrauterine growth restriction, preeclampsia and intrauterine mortality at high altitude in Bolivia. *Pediatr Res*. 2003; 54(1):20–5. [PubMed: 12700368]
13. Julian CG, Gonzales M, Rodriguez A, Bellido D, Salinas Salmon C, Ladenburger A, Reardon L, Vargas E, Moore LG. Perinatal hypoxia increases susceptibility to high-altitude polycythemia and attendant pulmonary vascular dysfunction. *American journal of physiology Heart and circulatory physiology*. 2015:ajpheart 00296.
14. Hypertension in pregnancy. . Report of the American College of Obstetricians and Gynecologists' Task Force on Hypertension in Pregnancy. *Obstet Gynecol*. 2013; 122(5):1122–31. [PubMed: 24150027]
15. Stegle O, Parts L, Durbin R, Winn J. A Bayesian framework to account for complex non-genetic factors in gene expression levels greatly increases power in eQTL studies. *PLoS computational biology*. 2010; 6(5):e1000770. [PubMed: 20463871]
16. Smyth GK. Linear models and empirical bayes methods for assessing differential expression in microarray experiments. *Statistical applications in genetics and molecular biology*. 2004; 3:Article3. [PubMed: 16646809]
17. Hothorn T, Bretz F, Westfall P. Simultaneous inference in general parametric models. *Biometrical journal Biometrische Zeitschrift*. 2008; 50(3):346–63. [PubMed: 18481363]
18. Pedersen BS, Schwartz DA, Yang IV, Kechris KJ. Comb-p: software for combining, analyzing, grouping and correcting spatially correlated P-values. *Bioinformatics*. 2012; 28(22):2986–8. [PubMed: 22954632]
19. Jaffe AE, Murakami P, Lee H, Leek JT, Fallin MD, Feinberg AP, Irizarry RA. Bump hunting to identify differentially methylated regions in epigenetic epidemiology studies. *Int J Epidemiol*. 2012; 41(1):200–9. [PubMed: 22422453]
20. Bell JT, Pai AA, Pickrell JK, Gaffney DJ, Pique-Regi R, Degner JF, Gilad Y, Pritchard JK. DNA methylation patterns associate with genetic and gene expression variation in HapMap cell lines. *Genome biology*. 2011; 12(1):R10. [PubMed: 21251332]
21. Kerkel K, Spadola A, Yuan E, Kosek J, Jiang L, Hod E, Li K, Murty VV, Schupf N, Vilain E, Morris M, Haghighi F, Tycko B. Genomic surveys by methylation-sensitive SNP analysis identify sequence-dependent allele-specific DNA methylation. *Nat Genet*. 2008; 40(7):904–8. [PubMed: 18568024]
22. Yang IV, Pedersen BS, Liu A, O'Connor GT, Teach SJ, Kattan M, Misiak RT, Gruchalla R, Steinbach SF, Szeffler SJ, Gill MA, Calatroni A, David G, Hennessy CE, Davidson EJ, Zhang W, Gergen P, Togias A, Busse WW, Schwartz DA. DNA methylation and childhood asthma in the inner city. *J Allergy Clin Immunol*. 2015

23. Sena JA, Wang L, Hu CJ. BRG1 and BRM chromatin-remodeling complexes regulate the hypoxia response by acting as coactivators for a subset of hypoxia-inducible transcription factor target genes. *Mol Cell Biol.* 2013; 33(19):3849–63. [PubMed: 23897427]
24. Bevilacqua A, Willis MS, Bultman SJ. SWI/SNF chromatin-remodeling complexes in cardiovascular development and disease. *Cardiovasc Pathol.* 2014; 23(2):85–91. [PubMed: 24183004]
25. Fang F, Chen D, Yu L, Dai X, Yang Y, Tian W, Cheng X, Xu H, Weng X, Fang M, Zhou J, Gao Y, Chen Q, Xu Y. Proinflammatory stimuli engage Brahma related gene 1 and Brahma in endothelial injury. *Circ Res.* 2013; 113(8):986–96. [PubMed: 23963727]
26. Khurshed M, Kolla JN, Kotapalli V, Gupta N, Gowrishankar S, Uppin SG, Sastry RA, Koganti S, Sundaram C, Pollack JR, Bashyam MD. ARID1B, a member of the human SWI/SNF chromatin remodeling complex, exhibits tumour-suppressor activities in pancreatic cancer cell lines. *Br J Cancer.* 2013; 108(10):2056–62. [PubMed: 23660946]
27. Pyagay P, Heroult M, Wang Q, Lehnert W, Belden J, Liaw L, Friesel RE, Lindner V. Collagen triple helix repeat containing 1, a novel secreted protein in injured and diseased arteries, inhibits collagen expression and promotes cell migration. *Circ Res.* 2005; 96(2):261–8. [PubMed: 15618538]
28. Nishimoto S, Hamajima Y, Toda Y, Toyoda H, Kitamura K, Komurasaki T. Identification of a novel smooth muscle associated protein, smap2, upregulated during neointima formation in a rat carotid endarterectomy model. *Biochim Biophys Acta.* 2002; 1576(1–2):225–30. [PubMed: 12031507]
29. Rocnik EF, Liu P, Sato K, Walsh K, Vaziri C. The novel SPARC family member SMOC-2 potentiates angiogenic growth factor activity. *J Biol Chem.* 2006; 281(32):22855–64. [PubMed: 16774925]
30. Lawler PR, Lawler J. Molecular basis for the regulation of angiogenesis by thrombospondin-1 and -2. *Cold Spring Harb Perspect Med.* 2012; 2(5):a006627. [PubMed: 22553494]
31. Chen CH, Ho HH, Wu ML, Layne MD, Yet SF. Modulation of cysteine-rich protein 2 expression in vascular injury and atherosclerosis. *Mol Biol Rep.* 2014; 41(11):7033–41. [PubMed: 25034893]
32. Nagaraj SR, Shilpa P, Rachaiah K, Salimath BP. Crosstalk between VEGF and MTA1 signaling pathways contribute to aggressiveness of breast carcinoma. *Mol Carcinog.* 2015; 54(5):333–50. [PubMed: 24265228]
33. Carter EB, Stuart JJ, Farland LV, Rich-Edwards J, Zera C, McElrath TF, Seely EW. [118-POS]: Validation of a maternal recall questionnaire for pregnancy complications associated with increased future risk for cardiovascular disease. *Pregnancy Hypertens.* 2015; 5(1):63.




Figure 1.

Differentially methylated regions (DMRs) were identified in young male Andean residents of high altitude (18–25 yrs) born to women with hypertensive complications of pregnancy (HTN_{PREG}) vs. women who remained normotensive throughout pregnancy (controls).

Shown is a Manhattan plot of the adjusted p-values for the comparison between HTN_{PREG} vs. controls. Each dot represents a p-value for a probe on the Illumina 450k array that has been adjusted by the significance of neighboring probes according to their correlation using Comb-p (18). Probes that comprise the DMRs are identified by red dots.

Table 1

Subject characteristics

	HTN _{PREG} Offspring (n = 5)	Controls (n = 19)	P-value
Age, yrs	21.4 ± 2.5	21.8 ± 1.5	NS
Height, cm	164.8 ± 3.1	167.8 ± 6.1	NS
Weight, kg	63.1 ± 3.4	62.8 ± 7.5	NS
Heart rate, bpm	64 ± 5	63 ± 10	NS
S _a O ₂ , %	90.4 ± 3.0	91.5 ± 2.4	NS
RVSP, mmHg	40.6 ± 1.1	37.4 ± 4.2	0.007
DLCO _{corr} , % predicted	121 ± 19	142 ± 29	NS
FEV ₁ /FVC	86.8 ± 5.8	85.8 ± 7.5	NS

Data are reported as the mean ± SD. Abbreviations: arterial oxygen saturation (S_aO₂), right ventricular systolic pressure (RVSP), pulmonary CO diffusion capacity (DLCO) corrected for altitude and hemoglobin, forced expiratory volume at one second (FEV₁) and forced vital capacity (FVC).

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 2

Differentially methylated regions (DMRs) identified in young male highlanders (18–25 yrs) born to women with hypertensive pregnancy (HTNPREG) vs. normotensive controls

DMR Gene Symbol Genomic Location	Ref Gene Feature(s)	Ref Gene Distance	Slk P-value	Slk-Sidak P-value	Probes	deltaBeta
CTHRC1 Chr8: 104383714 – 104383798	TSS, exon, 5'UTR	0	9.82E-07	0.006	cg04219247	-0.015
					cg20447655	-0.040
					cg20392240	-0.017
					cg11889769	-0.024
					cg21643403	-0.033
LINC00226 * Chr14: 106806142 – 106806179	intergenic	61176	1.97E-06	0.025	cg23404902	0.038
					cg01842774	0.072
TRIM31 Chr6: 30071502 - 30071612	intron	0	8.55E-08	0.0004	cg00354641	-0.034
					cg27123071	-0.028
					cg13436843	-0.086
					cg09735274	-0.087
ARID1B * Chr6: 159983262 - 156983315	intergenic	-115748	2.20E-07	0.002	cg05282260	0.047
					cg08839808	0.097
					cg00269725	0.072
SMOC2 * Chr6: 169284284 – 169284344	intergenic	215610	1.81E-08	0.001	cg17720554	0.052
					cg10213762	0.087
					cg12550496	0.100
LRR1Q3 Chr1: 74663582 - 74663750	intron, exon, 5'UTR	0	1.81E-08	5.11E-05	cg16356856	-0.106
					cg13900770	-0.085
					cg06034096	-0.094
					cg00402366	-0.162

* Denotes hypermethylated genes; the remainder are hypomethylated.

Genomic regions were defined according to the UCSC RefGene group. Intergenic = site not annotated in a gene, TSS = transcription start site at 200–1500 bp, 5' region = 5'UTR and 1st exon, Intragenic = gene body including introns and exons and, 3' region = 3'UTR, UTR – untranslated region. Individual sites that compose the differentially methylated regions and their percent methylation change are shown in the *Probes* and *deltaBeta* columns, respectively.