



Published in final edited form as:

J Intern Med. 2017 January ; 281(1): 75–85. doi:10.1111/joim.12544.

Ten-year alcohol consumption typologies and trajectories of C-reactive protein, interleukin-6 and interleukin-1 receptor antagonist over the following 12 years: a prospective cohort study

Steven Bell¹, Gautam Mehta², Kevin Moore², and Annie Britton¹

¹Research Department of Epidemiology and Public Health, University College London

²UCL Institute of Liver and Digestive Health, Royal Free Campus, University College London, London, UK

Abstract

Background—Moderate alcohol consumption is thought to confer cardiometabolic protective effects. Inflammatory pathways are hypothesised to partly underlie this association.

Objectives—The aim of this study was to examine the association between typologies of alcohol consumption and markers of inflammation, and their rate of change over time.

Methods—Data were collected from 8209 participants (69% men, mean age 50 years [SD 6.1]) of the British Whitehall II study. Alcohol consumption typologies were defined using up to three measures during an approximately 10-year period spanning from 1985 to 1994 as (i) stable non-drinkers, (ii) stable moderate drinkers (referent), (iii) stable heavy drinkers, (iv) non-stable drinkers and (v) former drinkers. C-reactive protein (CRP), interleukin (IL)-6 and IL-1 receptor antagonist (IL-1 RA) were measured up to three times in the following 12 years.

Results—Stable moderate drinkers had lower levels of CRP than stable non-drinkers, stable heavy drinkers, former drinkers and non-stable drinkers but there were no differences in the rate of change in CRP over time between groups. Stable non-drinkers had higher levels of IL-6 as did stable heavy drinkers; rates of change in IL-6 over time were also increased in the latter group. Stable non-drinkers also had higher levels of IL-1 RA. These associations were robust to adjustment for confounding factors.

Conclusion—Our novel investigation of 10-year drinking typologies shows that stable moderate alcohol consumption is associated with a long-term inflammatory marker profile that is consistent with conferring a reduced risk of developing CHD.

Correspondence: Steven Bell, Department of Public Health and Primary Care, University of Cambridge, Strangeways Research Laboratory, Worts Causeway, Cambridge CB1 8RN, UK; scb81@medschl.cam.ac.uk.
Alternative proof reader: Annie Britton (a.britton@ucl.ac.uk)

Author contributions

SB and AB conceived the research question. SB carried out the analysis. SB completed the first draft of the manuscript. AB, GM and KM provided additional intellectual content. All authors read and agreed on the final submitted manuscript.

Conflict of interest statement

None of the authors has any conflicts of interest to declare.

Keywords

alcohol; cytokines; epidemiology; inflammation; longitudinal

Introduction

Moderate alcohol consumption is thought to confer cardiometabolic protective effects [1–3] and has also been demonstrated to be related to a lower risk of a plethora other disorders of different aetiology compared to both no alcohol and heavy alcohol intake [4]. Numerous biological mechanisms have been put forward to explain the proposed cardiometabolic protection [5, 6], with favourable changes in high-density lipoprotein (HDL) cholesterol, fibrinogen and adiponectin supported by evidence from several small-scale randomised controlled feeding trials [7]. However, these factors are unlikely to entirely explain the protective effects observed for moderate consumption and cardiometabolic outcomes compared to abstinence (or the increased risk observed amongst heavier drinkers), and their causal role in the aetiology of cardiovascular disease (CVD) remains unclear [8–14].

Therefore, if the protective cardiometabolic effects observed are genuine it is likely that other biological pathways are involved, one of which may be via pro-inflammatory cytokines [15]. For example, higher levels of the acute-phase reactant C-reactive protein (CRP) are associated with an increased risk of developing CVD [16] and a variety of other disease endpoints [17]. Studies have demonstrated that higher levels of interleukin (IL)-6 are associated with an increased risk of coronary heart disease (CHD) [18], including studies examining long-term exposure to elevated IL-6 [19] or functional genetic variants of IL-6 signalling [20], suggesting that the association is causal. IL-1 is considered a master regulator of inflammation that triggers the release of a variety of inflammatory markers through activating the IL-1 receptor. The IL-1 receptor antagonist (IL-1 RA) is an endogenous inhibitor of IL-1 that prevents the activation of the IL-1 receptor by either IL-1 α or IL-1 β [21]. IL-1 RA is associated with a diverse range of diseases including CVD, type 2 diabetes, certain cancers and joint diseases such as arthritis [21–25]. In a recent large-scale Mendelian randomisation study it was found that genetically elevated levels of IL-1 RA were causally associated with an increased risk of CHD and abdominal aortic aneurysm [26].

The findings of studies investigating alcohol consumption and inflammatory markers have been mixed. There is limited and conflicting evidence from interventional studies, typically with sample sizes of less than 100 participants and over relatively short periods of time, of the association between alcohol consumption and inflammation [7, 27–29]. In terms of observational studies, some investigators have found that moderate alcohol consumption is associated with lower levels of CRP [30–32] and IL-6 [32, 33] compared to no alcohol and heavy alcohol intake, whereas others have observed no association [33, 34]. Studies of the association between alcohol intake in the general population and IL-1 RA are scarce [33].

Furthermore, most previous studies have only used a single measure of alcohol intake at baseline to define the drinking behaviour of participants, assuming that it is static thereafter. However, individuals drinking habits change over time [35, 36] and this can affect their risk

of developing disease [37]. Therefore not accounting for long-term drinking profiles or changes in alcohol consumption can introduce bias [38–40]. A classic example of such bias is the failure to separate former drinkers from never drinkers which is known to in some cases substantially impact findings and subsequent conclusions that are drawn [41, 42]. Levels of CRP, IL-6 and IL-1 RA also change over time and can similarly influence disease risk [25, 43, 44].

In summary, studies utilising repeat measures of alcohol consumption [35] or examining the influence of alcohol on the change in biomarker levels over time [45] are rare. Studies combining these two elements are non-existent. We therefore aimed to examine the association between typologies of alcohol consumption over an approximately 10-year period and markers of inflammation, including CRP, IL-6 and IL-1 RA, and their rate of change over the following 12 years.

Materials and methods

Study population

Data were obtained from the Whitehall II cohort of British civil servants [46]. The study started during the period 1985–1988 (phase 1) and included 10,308 participants (6895 men) aged 35–55 years. All civil servants within this age range in 20 London-based departments were invited by letter to participate, and 73% agreed. The first phase involved a clinical examination as well as a self-administered questionnaire to collect information on demographic characteristics, health, lifestyle factors, work characteristics, social support and life events. Subsequent phases of data collection have alternated between postal questionnaire alone and postal questionnaire accompanied by a clinical examination. Data used in this investigation were from phases 1 (1985–1988), 2 (1989–1990), 3 (1991–1994), 5 (1997–1999) and 7 (2002–2004) of the study. The maximum available sample was 8209 participants for whom there was information on alcohol consumption, age and gender, and at least one measure of CRP, IL-6 or IL-1 RA.

Ethical considerations and data access

The University College London Medical School committee on the Ethics of Human Research approved the Whitehall II study. Informed consent was obtained at baseline and renewed at each contact. Whitehall II study data, protocols and other metadata are available to *bona fide* researchers for research purposes (data sharing policy is available at <http://www.ucl.ac.uk/whitehallIII/data-sharing>).

Assessment of alcohol consumption

We used alcohol information collected at study phases 1, 2 and 3 to construct (approximately) 10-year drinking typologies. Participants were asked to report the number of alcoholic drinks they had consumed in the previous week, providing information separately for beer/cider (in pints), wine (in glasses) and spirits (in measures). Drinks were converted into grams of ethanol using a conservative estimate of 8 g for each measure of spirits and glass of wine, and 16 g for each pint of beer [47]. The sum of these converted measurements was then used to define total weekly amount of alcohol intake in grams. We

then constructed categories of alcohol consumption based on UK guidelines for sensible drinking at the time: none, moderate (within guidelines of 8–168 g ethanol per week for men and 8–112 g for women) and heavy (intake above these guidelines).

Typologies of drinkers to remain in keeping with the results over study phases 1–3 were defined as follows: stable non-drinkers, stable moderate drinkers, stable heavy drinkers, non-stable drinkers (participants who moved between categories of consumption during observation) and former drinkers [non-drinkers at study phase 3 who had previously reported consuming alcohol (at any level) at earlier study phases] Participants were permitted one missing alcohol value in the construction of the 10-year alcohol typology variable. Our reference group for analyses was stable moderate drinkers [48].

Assessment of inflammatory markers

Fasting serum samples were collected between 08.00 and 13.00, stored at -80°C and were not thawed or refrozen during storage. Although serum samples were from three different study phases, blood collection, processing and storage followed the same standard operating procedures.

CRP was measured with a high-sensitivity immunonephelometric assay in a BN ProSpec nephelometer (Dade Behring, Milton Keynes, UK). IL-6 was measured with a high-sensitivity enzyme-linked immunosorbent assay (ELISA) (R&D Systems, Oxford, UK). Values lower than the detection limit [0.154 mg/L for CRP (multiplied by 9524 in order to express the value in mmol/L) and 0.08 pg/mL for IL-6] were assigned a value equal to half the detection limit. We excluded samples with CRP concentrations suggestive of acute inflammation and related bacterial infection (>10 mg/L) [49] ($n = 242$). To measure short-term biological variation and laboratory error, a repeat sample was taken from 150 participants for CRP and 241 participants for IL-6 at phase 3 [with a mean elapsed time between samples of 32 days (SD 10.5)]. Intra-assay and inter-assay coefficients of variation were 4.7% and 8.3% for CRP and 7.5% and 8.9% for IL-6, respectively.

Serum IL-1 RA was measured in a diabetes case-cohort sample [25, 50] with the Quantikine ELISA kit (R&D Systems, Wiesbaden, Germany). All assays were performed consecutively in the same laboratory (German Diabetes Center), and samples from different study phases from the same participant were always measured using the same ELISA plate to minimize assay imprecision. Mean intra-assay and inter-assay coefficients of variation were 2.6% and 7.9%, respectively. The limit of detection was 14 pg/mL (all samples were above the limit of detection).

Covariates

Age, sex, ethnicity (white or non-white) and prevalent CHD (clinically verified events) and type 2 diabetes (cases defined by oral glucose tolerance tests and/or use of diabetes medication) at study phase 3 were entered into our statistical models as time-invariant predictors.

Time-varying covariates included in our models were socioeconomic position [defined using either current or last recorded civil service employment grade as high (unified grades 1–7),

intermediate (executive officers) or low (clerical or support staff), as previously described [51]] and health behaviours including smoking status (never, former and current) and physical activity (lowest sex-specific quartile of combined hours of moderate and vigorous physical activity defined as ‘physically inactive’). Diet quality was classified as poor or good using three questions on the type of milk and bread participants usually consumed and their frequency of fruit and vegetable intake. For each dietary component a score of one was assigned to poor diet quality indicators (whole milk, white bread, fruit and vegetable intake less than daily) and a summed score ≥ 2 was used to classify poor diet quality [52]. Body mass index (BMI) was calculated using the standard formula and participants were classified as normal weight (18.5–24.9 kg/m²), overweight/obese (≥ 25 kg/m²) or underweight (<18.5 kg/m²) using thresholds adopted by the World Health Organisation.

Statistical analysis

Differences in sample characteristics by drinking typologies were assessed using chi-squared test or one-way ANOVA. To examine the association between alcohol typologies and trajectories of inflammatory markers we used linear mixed models with time from phase 3 in years as the time metric (allowing for individually varying times of observation). We included random effects for the intercept and time and allowed for random effects to co-vary. As CRP, IL-6 and IL-1 RA were positively skewed we used natural logarithm-transformed values in all analyses (predictions were calculated using these models and then back-transformed to their original scale for graphical presentation). Preliminary analyses revealed no evidence of effect modification between drinking typologies and sex in the rate of change in any inflammatory marker. Therefore two models are presented for each outcome, one with adjustments only for age and sex and the other with adjustments for all covariates described above. These models include terms for the difference in inflammatory markers at study phase 3 by alcohol group (intercept differences) as well as interactions between drinking categories and time to describe differences in the rate of change in inflammatory markers per year of follow-up by alcohol typologies. We also include analyses with alcohol intake categories defined using data from the phase 3 assessment only, presented in the Online Supplementary Material, so that findings from the main analysis can be compared to those that would have been obtained using the conventional approach of only using alcohol intake assessed at one time point (as in the main analyses presented, these models include terms describing differences in baseline values of inflammatory markers as well as their rate of change over time). An α level of < 0.05 was considered statistically significant. All analyses were conducted using Stata 14.1.

Results

Descriptive statistics

The characteristics of participants at study phase 3 are presented in Table 1. The mean age of participants was 50 years (SD 6.1). Almost 70% of the participants were men, and the majority were of white ethnicity and high or intermediate socioeconomic position. Approximately half of the participants had never smoked whilst 14% were current smokers. The majority of participants were physically active and made good dietary choices. Half of the sample had a BMI considered to be in the normal range and 47% were considered

overweight. Prevalent CHD (3%) or type 2 diabetes (1%) was rare amongst participants. During follow-up there were 696 and 873 incident cases of CHD and type 2 diabetes, respectively. Descriptive statistics for changes in alcohol consumption and inflammatory markers are presented in Online Supplementary Table S1.

Ten-year drinking typologies

Regression coefficients of the associations between 10-year alcohol typologies and trajectories of CRP, IL-6 and IL-1 RA are shown in Table 2.

In multivariable adjusted models, all drinking typologies other than former drinkers had statistically significantly elevated levels of CRP at phase 3 in comparison to stable moderate drinkers. No drinking groups differed significantly in their rate of change in CRP over time.

All drinking typologies had elevated levels of IL-6 at phase 3 compared to moderate drinkers in the multivariable adjusted model (10.4% [95% confidence interval (95% CI) 4.9% to 16.3%] higher in stable non-drinkers, 11.5% [95% CI 6.2% to 17.0%] higher in stable heavy drinkers and 9.3% [95% CI 4.5% to 14.5%] higher in former drinkers). Stable heavy drinkers also showed increases in the rate of change in IL-6 levels over time [0.9% (95% CI 0.3% to 1.4%) per year]. Non-stable drinkers had slightly shallower increases in their rate of change in IL-6 over time [-0.4% (95% CI -0.7% to ~0.0%)] compared to moderate drinkers.

Stable non-drinkers had elevated levels of IL-1 RA compared to moderate drinkers at phase 3 [6.7% (95% CI 0.7% to 13.1%)]. No significant effects on the levels of IL-1 RA were observed for any other drinking typology, nor were any significant effects observed for any drinking typology on the rate of change in IL-1 RA over time.

The mean trajectories of each biomarker, predicted using the multivariable model, by drinking typology are presented graphically in Fig. 1.

Findings based on drinking information alone at study phase 3 were similar to those observed using 10-year drinking typologies (see Online Supplementary Table S2 and Fig. S1). However, whilst the findings were broadly concordant, those based on information on drinking behaviour over a 10-year period revealed that participants with a sustained history of heavy drinking had higher levels of inflammation as well as steeper increases in IL-6 values than were seen when using current drinking information alone. As a *post hoc* analysis for further comparison with evidence from small-scale feeding trials [28], we repeated the analysis for changes in fibrinogen during an average period of ~5 years. We found that compared to moderate drinkers, both former drinkers and stable non-drinkers had higher levels throughout follow-up. Conversely, heavy drinkers had, on average, the lowest levels of fibrinogen during observation (Online Supplementary Tables S3 and S4 and Fig. S2).

Discussion

Summary of findings

We observed that those who consistently consumed alcohol at levels considered moderate over a 10-year period had lower concentrations of CRP, IL-6 and IL-1 RA compared to non-

drinkers during the following 12 years. Former drinkers also had higher levels of each of these markers during this period, although this association was not statistically significant. Additionally, we found that heavy drinkers not only had increased levels of these cytokines compared to stable moderate drinkers, but also demonstrated a more marked rate of change in IL-6 levels over time.

Interpretation, comparison to other work and implications

Our findings are broadly consistent with the U- or J-shaped associations observed between alcohol consumption and cardiometabolic outcomes. Thus, we found that non-drinkers, former drinkers and heavy drinkers had consistently elevated levels of CRP, IL-6 and IL-1 RA over a 12-year period. These findings are supported by evidence from other studies of the effect of alcohol consumption on inflammatory markers [30, 31]. Two of these markers (IL-6 and IL-1 RA) are believed to be causally associated with CHD, providing indirect evidence that moderate alcohol consumption may confer some cardioprotective effects. The mechanisms by which alcohol achieves this are unknown, but may involve mild activation of inflammatory pathways, which confer some benefit. On the other hand these pathways are also likely to be involved in the detrimental effects of heavy drinking. These observations are concordant with large-scale Mendelian randomisation studies that have shown that inhibiting IL-6 signalling could reduce the risk of developing CHD [53] whilst inhibiting IL-1 α/β may increase the risk of CHD [26]. This highlights not only the complexity of inflammatory pathways underlying CVDs but also the complicated role of alcohol consumption in the development of such diseases. For example, moderate alcohol intake may induce oxidative stress by induction of particular heat shock proteins, which inhibit the activation of various pro-inflammatory cytokines leading to endotoxin tolerance [54]. Endotoxin tolerance is thought to be a protective mechanism against developing coronary occlusion and acute coronary syndromes [55], and may explain why moderate drinkers have a lower risk of CHD than non-drinkers. By contrast, a pattern of heavy alcohol use may lead to low-level gut bacterial translocation and consequent increased levels of pro-inflammatory cytokines. It was recently demonstrated that an acute alcohol 'binge' of 0.8 g/kg in healthy individuals led to a rapid increase in serum endotoxin and bacterial DNA, as well as increased acute-phase protein levels and pro-inflammatory cytokine responses [56]. This pattern of low-level bacterial translocation has been associated with disease progression in animal models of obesity and metabolic syndrome [57, 58].

In terms of overall effect sizes, the multivariable-adjusted effect of stable non-drinking on the intercept of IL-6 is equivalent to the per-allele effect of single-nucleotide polymorphism (SNP) rs7529229 in the *IL6R* gene (Ch1q21.3) on IL-6 levels [53] which confers an approximately 5% reduction in odds of developing CHD. CRP levels were increased by 15.7% in non-drinkers compared to stable moderate drinkers, which is similar to the per-allele difference observed for *CRP* SNP rs1130864 [59]. However, this elevated level of CRP has been shown not to confer a protective or detrimental effect for CHD [59]. The effect of non-drinking on IL-1 RA although statistically significant was relatively small, and less than the effect observed for a genetic allele count score using two common, uncorrelated, SNPs (rs6743376 and rs1542176) located upstream of *ILRN* (the gene that encodes the IL-1 RA) [26]. Given that a per-allele increase in this genetic score was only

associated with a 3% increase in the odds of developing CHD and not significantly associated with developing type 2 diabetes [26], the relative contribution of moderate drinking to a reduced risk of CHD via IL-1 α/β is likely to be minimal.

Our longitudinal approach demonstrates the importance of accounting for repeat measures of alcohol intake when estimating associations with health outcomes [35]. We found that only considering alcohol consumption at one point in time led to underestimation of the effect of heavy drinking on inflammatory markers. Furthermore, we observed that former drinkers had higher levels of cytokines than stable non-drinkers, in keeping with the idea that failing to exclude former drinkers from the group of stable non-drinkers will lead to overestimating the protective effect of moderate drinking [42]. This group may consist of former heavy drinkers (including individuals with a history of alcohol use disorders) and those with illnesses that have led to alcohol cessation [60], both of which are likely to be associated with higher levels of inflammation [61–63]. However, it is also worth noting that amongst those with existing illness, alcohol abstinence is generally associated with better long-term prognosis than continued drinking [64, 65]. Furthermore, whilst it has been shown that CVD risk factors are more common in non-drinkers [66], even when life-long non-drinkers can be separated from former and occasional drinkers, ill-health prior to the age at which alcohol consumption typically begins has been shown to be more common in those who have never consumed alcohol [67]. All of these factors may contribute to higher levels of inflammation as well as increased risk of CVD in non-drinkers. As such, the protective effect of moderate drinking on cardiometabolic outcomes is likely to be more modest than is often believed [41, 68].

Strengths and weaknesses

Our study has several strengths including the large sample size and use of repeat measures of alcohol consumption to define alcohol typologies over a 10-year period (reducing drinking category misclassification bias inherent in studies utilising only one measure of alcohol consumption) as well as repeat measures of markers of inflammation.

However, there are also several shortcomings. For example, it is known that heavy drinkers are under-represented in population level surveys; the drinkers in our sample are typically low to moderate consumers [35]. This means that the effect we observed for heavy drinking on trajectories of inflammatory markers is likely to be an under-estimate of the ‘true’ association.

Another limitation is that we used self-reported measures of alcohol consumption which have been criticised [69]. Furthermore, we focused on total weekly quantity of alcohol intake, not changes in both frequency and quantity. It is plausible that shifts in pattern of consumption may better explain differences in risk of CHD over time [70]. However, this does not affect our comparison with existing work which has also mostly focused on singular components of drinking behaviour (also largely overall weekly quantity of alcohol consumption).

Finally, participants in the Whitehall II study are not a representative sample of the general population, however it has been shown that cardiometabolic-related findings from this cohort are broadly consistent with those obtained from representative cohorts [71].

Conclusion

Stable moderate alcohol consumption is associated with a shift in the long-term inflammatory marker profile which is consistent with conferring a reduced risk of developing CHD. The implication of this is that modulation of chronic inflammation is a significant pathway through which alcohol affects overall CHD risk.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

We would like to thank all participating study members and all members of the Whitehall II study team. In addition, we thank the two anonymous referees for their careful reading of our original manuscript and helpful suggestions for improvement.

Sources of funding

This work was supported by grants from the European Research Council (ERC-StG-2012-309337_AlcoholLifecourse, PI: Annie Britton, <http://www.ucl.ac.uk/alcohol-lifecourse>) and UK Medical Research Council/Alcohol Research UK (MR/M006638/1). The Whitehall II study is supported by grants from the UK Medical Research Council (K013351), British Heart Foundation (RG/07/008/23674), Stroke Association, National Heart Lung and Blood Institute (HL036310) and National Institute on Aging (AG13196, AG034454). The funders had no role in the design of the study, data collection and analysis, or the preparation of and decision to publish the manuscript.

References

1. Ronksley PE, Brien SE, Turner BJ, Mukamal KJ, Ghali WA. Association of alcohol consumption with selected cardiovascular disease outcomes: a systematic review and meta-analysis. *BMJ*. 2011; 342:d671. [PubMed: 21343207]
2. Gémes K, Janszky I, Laugsand LE, et al. Alcohol consumption is associated with a lower incidence of acute myocardial infarction: results from a large prospective population-based study in Norway. *J Intern Med*. 2016; 279:365–375. [PubMed: 26365927]
3. Roerecke M, Rehm J. The cardioprotective association of average alcohol consumption and ischaemic heart disease: a systematic review and meta-analysis. *Addiction*. 2012; 107:1246–1260. [PubMed: 22229788]
4. Fekjær HO. Alcohol—a universal preventive agent? A critical analysis. *Addiction*. 2013; 108:2051–2057. [PubMed: 23297738]
5. Mathews M, Liebenberg L, Mathews E. The mechanism by which moderate alcohol consumption influences coronary heart disease. *Nutr J*. 2015; 14:33. [PubMed: 25889723]
6. Movva R, Figueredo VM. Alcohol and the heart: To abstain or not to abstain? *Int J Cardiol*. 2013; 164:267–276. [PubMed: 22336255]
7. Brien SE, Ronksley PE, Turner BJ, Mukamal KJ, Ghali WA. Effect of alcohol consumption on biological markers associated with risk of coronary heart disease: systematic review and meta-analysis of interventional studies. *BMJ*. 2011; 342:d636. [PubMed: 21343206]
8. Holmes MV, Asselbergs FW, Palmer TM, et al. Mendelian randomization of blood lipids for coronary heart disease. *Eur Heart J*. 2014; 36:539–550. [PubMed: 24474739]
9. Barter PJ, Caulfield M, Eriksson M, et al. Effects of Torcetrapib in Patients at High Risk for Coronary Events. *N Engl J Med*. 2007; 357:2109–2122. [PubMed: 17984165]

10. Yaghoobkar H, Lamina C, Scott RA, et al. Mendelian Randomisation Studies Do Not Support a Causal Role for Reduced Circulating Adiponectin Levels in Insulin Resistance and Type 2 Diabetes. *Diabetes*. 2013; 62:3589–3598. [PubMed: 23835345]
11. Keavney B, Danesh J, Parish S, et al. Fibrinogen and coronary heart disease: test of causality by "Mendelian randomization". *Int J Epidemiol*. 2006; 35:935–943. [PubMed: 16870675]
12. Sook Lee E, Park S, Kim E, et al. Association between adiponectin levels and coronary heart disease and mortality: a systematic review and meta-analysis. *Int J Epidemiol*. 2013; 42:1029–1039. [PubMed: 23739486]
13. Borges MC, Lawlor DA, de Oliveira C, White J, Horta B, Barros AJ. The Role of Adiponectin in Coronary Heart Disease Risk: A Mendelian Randomization Study. *Circ Res*. 2016
14. Burgess S, Harshfield E. Mendelian randomization to assess causal effects of blood lipids on coronary heart disease: lessons from the past and applications to the future. *Curr Opin Endocrinol Diabetes Obes*. 2016; 23
15. Imhof A, Koenig W. Alcohol inflammation and coronary heart disease. *Addict Biol*. 2003; 8:271–277. [PubMed: 13129828]
16. The Emerging Risk Factors Collaboration. C-Reactive Protein, Fibrinogen, and Cardiovascular Disease Prediction. *N Engl J Med*. 2012; 367:1310–1320. [PubMed: 23034020]
17. Ansar W, Ghosh S. C-reactive protein and the biology of disease. *Immunol Res*. 2013; 56:131–142. [PubMed: 23371836]
18. Kaptoge S, Seshasai SRK, Gao P, et al. Inflammatory cytokines and risk of coronary heart disease: new prospective study and updated meta-analysis. *Eur Heart J*. 2014; 35:578–589. [PubMed: 24026779]
19. Danesh J, Kaptoge S, Mann AG, et al. Long-Term Interleukin-6 Levels and Subsequent Risk of Coronary Heart Disease: Two New Prospective Studies and a Systematic Review. *PLoS Med*. 2008; 5:e78. [PubMed: 18399716]
20. IL6R Genetics Consortium Emerging Risk Factors Collaboration. Interleukin-6 receptor pathways in coronary heart disease: a collaborative meta-analysis of 82 studies. *The Lancet*. 2012; 379:1205–1213.
21. Dinarello CA, Simon A, van der Meer JWM. Treating inflammation by blocking interleukin-1 in a broad spectrum of diseases. *Nat Rev Drug Discov*. 2012; 11:633–652. [PubMed: 22850787]
22. Garlanda C, Dinarello CA, Mantovani A. The Interleukin-1 Family: Back to the Future. *Immunity*. 2013; 39:1003–1018. [PubMed: 24332029]
23. Van Tassell BW, Toldo S, Mezzaroma E, Abbate A. Targeting Interleukin-1 in Heart Disease. *Circulation*. 2013; 128:1910–1923. [PubMed: 24146121]
24. Witkin SS, Gerber S, Ledger WJ. Influence of Interleukin-1 Receptor Antagonist Gene Polymorphism on Disease. *Clin Infect Dis*. 2002; 34:204–209. [PubMed: 11740709]
25. Carstensen M, Herder C, Kivimäki M, et al. Accelerated Increase in Serum Interleukin-1 Receptor Antagonist Starts 6 Years Before Diagnosis of Type 2 Diabetes: Whitehall II Prospective Cohort Study. *Diabetes*. 2010; 59:1222–1227. [PubMed: 20185814]
26. The Interleukin 1 Genetics Consortium. Cardiometabolic effects of genetic upregulation of the interleukin 1 receptor antagonist: a Mendelian randomisation analysis. *Lancet Diabetes Endocrinol*. 2015; 3:243–253. [PubMed: 25726324]
27. Chiva-Blanch G, Magraner E, Condines X, et al. Effects of alcohol and polyphenols from beer on atherosclerotic biomarkers in high cardiovascular risk men: A randomized feeding trial. *Nutr Metab Cardiovasc Dis*. 2015; 25:36–45. [PubMed: 25183453]
28. Stote KS, Tracy RP, Taylor PR, Baer DJ. The effect of moderate alcohol consumption on biomarkers of inflammation and hemostatic factors in postmenopausal women. *Eur J Clin Nutr*. 2016; 70:470–474. [PubMed: 26554758]
29. Chiva-Blanch G, Urpi-Sarda M, Llorach R, et al. Differential effects of polyphenols and alcohol of red wine on the expression of adhesion molecules and inflammatory cytokines related to atherosclerosis: a randomized clinical trial. *Am J Clin Nutr*. 2012
30. Imhof A, Woodward M, Doering A, et al. Overall alcohol intake, beer, wine, and systemic markers of inflammation in western Europe: results from three MONICA samples (Augsburg, Glasgow, Lille). *Eur Heart J*. 2004; 25:2092–2100. [PubMed: 15571824]

31. Imhof A, Froehlich M, Brenner H, Boeing H, Pepys MB, Koenig W. Effect of alcohol consumption on systemic markers of inflammation. *The Lancet*. 2001; 357:763–767.
32. Volpato S, Pahor M, Ferrucci L, et al. Relationship of Alcohol Intake With Inflammatory Markers and Plasminogen Activator Inhibitor-1 in Well-Functioning Older Adults: The Health, Aging, and Body Composition Study. *Circulation*. 2004; 109:607–612. [PubMed: 14769682]
33. Marques-Vidal P, Bochud M, Bastardot F, et al. Associations between alcohol consumption and selected cytokines in a Swiss population-based sample (CoLaus study). *Atherosclerosis*. 2012; 222:245–250. [PubMed: 22420891]
34. Hamer M, Stamatakis E. The accumulative effects of modifiable risk factors on inflammation and haemostasis. *Brain Behav Immun*. 2008; 22:1041–1043. [PubMed: 18411023]
35. Britton A, Ben-Shlomo Y, Benzeval M, Kuh D, Bell S. Life course trajectories of alcohol consumption in the United Kingdom using longitudinal data from nine cohort studies. *BMC Med*. 2015; 13:47. [PubMed: 25858476]
36. Britton A, Bell S. Reasons why people change their alcohol consumption in later life: Findings from the Whitehall II Cohort Study. *PLoS ONE*. 2015; 10:e0119421. [PubMed: 25756213]
37. Britton A, Marmot MG, Shipley MJ. How does variability in alcohol consumption over time affect the relationship with mortality and coronary heart disease? *Addiction*. 2010; 105:639–645. [PubMed: 20148795]
38. Greenfield TK, Kerr WC. Commentary on Liang & Chikritzhs (2011): Quantifying the impacts of health problems on drinking and subsequent morbidity and mortality – life-course measures are essential. *Addiction*. 2011; 106:82–83. [PubMed: 21618726]
39. Bell S, Britton A. The Role of Alcohol Consumption in Regulating Circulating Levels of Adiponectin: A Prospective Cohort Study. *J Clin Endocrinol Metab*. 2015; 100:2763–2768. [PubMed: 26000546]
40. The Fibrinogen Studies Collaboration. Regression dilution methods for meta-analysis: assessing long-term variability in plasma fibrinogen among 27 247 adults in 15 prospective studies. *Int J Epidemiol*. 2006; 35:1570–1578. [PubMed: 17148467]
41. Knott C, Bell S, Britton A. Alcohol Consumption and the Risk of Type 2 Diabetes: A Systematic Review and Dose-Response Meta-analysis of More Than 1.9 Million Individuals From 38 Observational Studies. *Diabetes Care*. 2015; 38:1804–1812. [PubMed: 26294775]
42. Chikritzhs T, Stockwell T, Naimi T, Andreasson S, Dangardt F, Liang W. Has the leaning tower of presumed health benefits from “moderate” alcohol use finally collapsed? *Addiction*. 2015; 110:726–727. [PubMed: 25613200]
43. Tabák AG, Kivimäki M, Brunner EJ, et al. Changes in C-reactive protein levels before type 2 diabetes and cardiovascular death: the Whitehall II study. *Eur J Endocrinol*. 2010; 163:89–95. [PubMed: 20573938]
44. Metti AL, Aizenstein H, Yaffe K, et al. Trajectories of peripheral interleukin-6, structure of the hippocampus, and cognitive impairment over 14 years in older adults. *Neurobiol Aging*. 2015; 36:3038–3044. [PubMed: 26279115]
45. Janssen I, Landay AL, Ruppert K, Powell LH. Moderate wine consumption is associated with lower hemostatic and inflammatory risk factors over 8 years: The study of women’s health across the nation (SWAN). *Nutr Aging Amst Neth*. 2014; 2:91.
46. Marmot M, Brunner E. Cohort Profile: The Whitehall II study. *Int J Epidemiol*. 2005; 34:251–256. [PubMed: 15576467]
47. Britton A, O’Neill D, Bell S. Underestimating the Alcohol Content of a Glass of Wine: The Implications for Estimates of Mortality Risk. *Alcohol Alcohol*. 2016
48. Rehm J, Irving H, Ye Y, Kerr WC, Bond J, Greenfield TK. Are Lifetime Abstainers the Best Control Group in Alcohol Epidemiology? On the Stability and Validity of Reported Lifetime Abstinence. *Am J Epidemiol*. 2008; 168:866–871. [PubMed: 18701442]
49. Myers GL, Rifai N, Tracy RP, et al. CDC/AHA Workshop on Markers of Inflammation and Cardiovascular Disease: Application to Clinical and Public Health Practice: Report From the Laboratory Science Discussion Group. *Circulation*. 2004; 110:e545–e549. [PubMed: 15611379]
50. Tabák AG, Carstensen M, Witte DR, et al. Adiponectin Trajectories Before Type 2 Diabetes Diagnosis: Whitehall II study. *Diabetes Care*. 2012; 35:2540–2547. [PubMed: 22933430]

51. Britton A, Marmot M. Different measures of alcohol consumption and risk of coronary heart disease and all-cause mortality: 11-year follow-up of the Whitehall II Cohort Study. *Addiction*. 2004; 99:109–116. [PubMed: 14678069]
52. Bell S, Britton A. Drinking pattern during midlife and risk of developing depression during 28 years of follow-up: A prospective cohort study. *Drug Alcohol Depend*. 2015; 155:111–117. [PubMed: 26321670]
53. The Interleukin-6 Receptor Mendelian Randomisation Analysis (IL6R MR) Consortium. The interleukin-6 receptor as a target for prevention of coronary heart disease: a mendelian randomisation analysis. *The Lancet*. 2012; 379:1214–1224.
54. Muralidharan S, Ambade A, Fulham MA, Deshpande J, Catalano D, Mandrekar P. Moderate Alcohol Induces Stress Proteins HSF1 and hsp70 and Inhibits Proinflammatory Cytokines Resulting in Endotoxin Tolerance. *J Immunol*. 2014; 193:1975–1987. [PubMed: 25024384]
55. Biswas SK, Lopez-Collazo E. Endotoxin tolerance: new mechanisms, molecules and clinical significance. *Trends Immunol*. 2009; 30:475–487. [PubMed: 19781994]
56. Bala S, Marcos M, Gattu A, Catalano D, Szabo G. Acute Binge Drinking Increases Serum Endotoxin and Bacterial DNA Levels in Healthy Individuals. *PLoS ONE*. 2014; 9:e96864. [PubMed: 24828436]
57. Vijay-Kumar M, Aitken JD, Carvalho FA, et al. Metabolic Syndrome and Altered Gut Microbiota in Mice Lacking Toll-Like Receptor 5. *Science*. 2010; 328:228–231. [PubMed: 20203013]
58. Cani PD, Amar J, Iglesias MA, et al. Metabolic Endotoxemia Initiates Obesity and Insulin Resistance. *Diabetes*. 2007; 56:1761–1772. [PubMed: 17456850]
59. C Reactive Protein Coronary Heart Disease Genetics Collaboration. Association between C reactive protein and coronary heart disease: mendelian randomisation analysis based on individual participant data. *BMJ*. 2011; 342:d548. [PubMed: 21325005]
60. Ng Fat L, Cable N, Shelton N. Worsening of Health and a Cessation or Reduction in Alcohol Consumption to Special Occasion Drinking Across Three Decades of the Life Course. *Alcohol Clin Exp Res*. 2015; 39:166–174. [PubMed: 25623415]
61. Leclercq S, De Saeger C, Delzenne N, de Timary P, Stärkel P. Role of Inflammatory Pathways, Blood Mononuclear Cells, and Gut-Derived Bacterial Products in Alcohol Dependence. *Alcohol Smok*. 2014; 76:725–733.
62. Pawelec G, Goldeck D, Derhovanessian E. Inflammation, ageing and chronic disease. *Host Pathog Immune Senescence*. 2014; 29:23–28.
63. Dwarkasing J, Marks DL, Witkamp R, van Norren K. Hypothalamic inflammation and food intake regulation during chronic illness. *Pept Gut-Brain Mech Hunger Satiety – Full4Health Proj*. 2016; 77:60–66.
64. Verrill C, Markham H, Templeton A, Carr NJ, Sheron N. Alcohol-related cirrhosis—early abstinence is a key factor in prognosis, even in the most severe cases. *Addiction*. 2009; 104:768–774. [PubMed: 19344445]
65. Powell WJ, Klatskin G. Duration of survival in patients with Laennec’s cirrhosis: Influence of alcohol withdrawal, and possible effects of recent changes in general management of the disease. *Am J Med*. 1968; 44:406–420. [PubMed: 5641303]
66. Naimi TS, Brown DW, Brewer RD, et al. Cardiovascular risk factors and confounders among nondrinking and moderate-drinking U.S. adults. *Am J Prev Med*. 2005; 28:369–373. [PubMed: 15831343]
67. Ng Fat L, Shelton N. Associations between self-reported illness and non-drinking in young adults. *Addiction*. 2012; 107:1612–1620. [PubMed: 22404244]
68. Rehm J, Shield KD, Roerecke M, Gmel G. Modelling the impact of alcohol consumption on cardiovascular disease mortality for comparative risk assessments: an overview. *BMC Public Health*. 2016; 16:1–9. [PubMed: 26728978]
69. Bellis MA, Hughes K, Jones L, et al. Holidays, celebrations, and commiserations: Measuring drinking during feasting and fasting to improve national and individual estimates of alcohol consumption. *BMC Med*. 2015; 13:113. [PubMed: 25998218]
70. Roerecke M, Rehm J. Alcohol consumption, drinking patterns, and ischemic heart disease: a narrative review of meta-analyses and a systematic review and meta-analysis of the impact of

heavy drinking occasions on risk for moderate drinkers. *BMC Med.* 2014; 12:182. [PubMed: 25567363]

71. Batty GD, Shipley M, Tabák A, et al. Generalizability of Occupational Cohort Study Findings. *Epidemiology.* 2014; 25:932–933. [PubMed: 25265141]




Fig. 1. Model-predicted C-reactive protein, interleukin-6 and interleukin-1 receptor antagonist trajectories by 10-year drinking typologies

Multivariable adjustment for age, sex, ethnicity, prevalent coronary heart disease or type 2 diabetes at phase 3, socioeconomic position, smoking status, physical activity, diet and body mass index. Graphs are based on the back transformation of log-transformed C-reactive protein, interleukin-6 and interleukin-1 receptor antagonist predicted by linear mixed models (Table 2).

Table 1

Characteristics at phase 3 of Whitehall II study by 10-year drinking typologies

	Stable non-drinker		Stable moderate drinker		Stable heavy drinker		Non-stable drinker		Former drinker		Total		P for diff
	n	%	n	%	n	%	n	%	n	%	n	%	
Age, mean (SD) years	746	51.3 (6.1)	3909	50.3 (6.1)	671	48.8 (5.7)	2052	49.6 (6.0)	831	50.9 (6.2)	8209	50.1 (6.1)	<0.001
Sex													
Male	345	46	2885	74	570	85	1430	70	465	56	5695	69	
Female	401	54	1024	26	101	15	622	30	366	44	2514	31	<0.001
Ethnicity													
White	494	66	3642	93	658	98	1895	92	731	88	7420	90	
Non-white	252	34	267	7	13	2	157	8	100	12	789	10	<0.001
SEP													
High	106	15	1719	45	339	51	761	38	166	20	3091	38	
Intermediate	310	43	1654	43	293	44	962	48	423	51	3642	45	
Low	308	43	463	12	29	4	284	14	237	29	1321	16	<0.001
Smoking status													
Never smoker	418	63	1802	50	162	26	795	41	387	50	3564	47	
Ex-smoker	144	22	1368	38	323	51	827	43	263	34	2925	39	
Current smoker	98	15	406	11	143	23	311	16	129	17	1087	14	<0.001
Physical activity level													
Active	642	89	3649	95	629	95	1868	93	747	90	7535	94	
Inactive	81	11	183	5	31	5	137	7	81	10	513	6	<0.001
Dietary choice quality													
Good	511	71	2978	78	478	72	1532	76	612	74	6111	76	
Poor	212	29	857	22	182	28	473	24	217	26	1941	24	<0.001
BMI category													
Normal weight	342	48	2035	54	292	46	982	50	418	52	4069	52	

	Stable non-drinker		Stable moderate drinker		Stable heavy drinker		Non-stable drinker		Former drinker		Total		P for diff
	n	%	n	%	n	%	n	%	n	%	n	%	
Overweight or Obese	361	50	1691	45	347	54	964	49	382	47	3745	48	
Underweight	16	2	37	1	3	1	14	1	6	1	76	1	<0.001
Prevalent CHD at phase 3													
No	709	95	3791	97	654	98	1987	97	804	97	7945	97	
Yes	37	5	118	3	17	3	65	3	27	3	264	3	0.07
Prevalent type 2 diabetes at phase 3													
No	716	96	3868	99	660	98	2024	99	818	98	8086	99	
Yes	30	4	41	1	11	2	28	1	13	2	123	2	<0.001

BMI, body mass index; CHD, coronary heart disease; SEP, socioeconomic position; SD, standard deviation.

Table 2

Fixed-effect coefficients from linear mixed model regression for association between 10-year drinking typologies and inflammatory marker trajectories during the following 12 years

	Log _e C-reactive protein (mg/L)	Multivariable adjusted	Log _e interleukin-6 (ng/mL)	Multivariable ^d adjusted	Log _e interleukin-1 receptor antagonist (ng/mL)	Multivariable adjusted
Intercept	Age and sex adjusted -0.280 *** [-0.316,-0.244]	-0.679 *** [-0.725,-0.632]	Age and sex adjusted 0.329 *** [0.310,0.348]	0.147 *** [0.122,0.173]	Age and sex adjusted 5.554 *** [5.534,5.573]	5.408 *** [5.381,5.435]
Stable non-drinker	0.231 *** [0.142,0.320]	0.146 ** [0.057,0.236]	0.174 *** [0.124,0.224]	0.099 *** [0.048,0.151]	0.072 * [0.010,0.134]	0.065 * [0.007,0.123]
Stable moderate drinker	0.000 [ref]	0.000 [ref]	0.000 [ref]	0.000 [ref]	0.000 [ref]	0.000 [ref]
Stable heavy drinker	0.251 *** [0.160,0.343]	0.156 *** [0.067,0.244]	0.142 *** [0.094,0.191]	0.109 *** [0.060,0.157]	0.033 [-0.016,0.083]	-0.005 [-0.053,0.044]
Non-stable drinker	0.133 *** [0.074,0.192]	0.075 * [0.018,0.132]	0.064 *** [0.033,0.095]	0.039 * [0.008,0.070]	0.011 [-0.022,0.044]	-0.009 [-0.040,0.023]
Former drinker	0.121 ** [0.035,0.207]	0.071 [-0.011,-0.154]	0.118 *** [0.073,0.164]	0.089 *** [0.044,0.135]	0.045 [-0.007,0.097]	0.041 [-0.009,0.092]
Time (per year)	0.033 *** [0.030,0.037]	0.027 *** [0.024,0.031]	0.020 *** [0.018,0.022]	0.019 *** [0.017,0.021]	0.034 *** [0.032,0.036]	0.031 *** [0.028,0.033]
Stable non-drinker × time	0.004 [-0.005,0.012]	0.004 [-0.005,0.013]	-0.002 [-0.007,0.004]	-0.000 [-0.006,0.005]	0.002 [-0.005,0.009]	0.004 [-0.002,0.011]
Stable moderate drinker × time	0.000 [ref]	0.000 [ref]	0.000 [ref]	0.000 [ref]	0.000 [ref]	0.000 [ref]
Stable heavy drinker × time	-0.005 [-0.013,0.004]	-0.005 [-0.014,0.004]	0.010 *** [0.005,0.016]	0.009 ** [0.003,0.014]	0.004 [-0.002,0.010]	0.005 [-0.001,0.011]
Non-stable drinker × time	-0.003 [-0.008,0.003]	-0.003 [-0.008,0.003]	-0.004 * [-0.007,-0.000]	-0.004 * [-0.007,-0.000]	0.000 [-0.004,0.004]	0.001 [-0.003,0.005]
Former drinker × time	0.003 [-0.006,0.011]	0.003 [-0.006,0.011]	-0.004 [-0.009,0.001]	-0.004 [-0.009,0.001]	0.005 [-0.000,0.011]	0.005 [-0.001,0.010]
N (no. of observations)	8128 (19,097)	7814 (18,097)	8183 (19,182)	7868 (18,179)	3468 (9564)	3443 (9244)

*** P<0.001;

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

 $P < 0.001$;

*
 $P < 0.05$.

[†]Multivariable adjustment for age, sex, ethnicity, prevalent coronary heart disease or type 2 diabetes at phase 3, socioeconomic position, smoking status, physical activity, diet and body mass index.