



Published in final edited form as:

Kidney Int Rep. 2017 September ; 2(5): 866–873. doi:10.1016/j.ekir.2017.05.002.

Association of Sleep Duration, Symptoms, and Disorders with Mortality in Adults with Chronic Kidney Disease

Ana C. Ricardo, MD, MPH^{1,*}, Vivien Goh^{1,*}, Jinsong Chen, PhD¹, Esteban Cedillo-Couvert, MD¹, Mary Kapella, PhD, RN², Bharati Prasad, MD, MS^{1,4}, Sharmila Parvathaneni, MD², Kristen Knutson, PhD³, and James P. Lash, MD.¹

¹Department of Medicine, University of Illinois, Chicago, IL

²Department of Biobehavioral Health Science, College of Nursing, University of Illinois, Chicago, IL

³Department of Medicine, University of Chicago, Chicago, IL

⁴Jesse Brown VA Medical Center, Chicago IL

Abstract

Introduction—In general populations, short and long sleep duration, poor sleep quality and sleep disorders have been associated with increased risk of death. We evaluated these associations in individuals with chronic kidney disease (CKD).

Methods—Prospective cohort study of 1,452 National Health and Nutrition Examination Survey (NHANES) 2005-2008 participants with CKD. CKD was defined by estimated glomerular filtration rate <60 ml/min/1.73m² or urine albumin-to-creatinine ratio ≥ 30 mg/g. Sleep duration, sleep symptoms (difficulty falling asleep, difficulty staying asleep, daytime sleepiness and non-restorative sleep), and sleep disorders (restless legs syndrome and sleep apnea) were self-reported. Vital status was determined using NHANES mortality linkage through December 2011.

Results—Mean age was 61 years, 58% were women, and 75% non-Hispanic white. During 4.4 years of median follow-up, we observed 234 deaths of which 75 were due to cardiovascular causes. In multivariable analyses, compared with individuals who reported 7-8 hours of sleep, HR (95% CI) for all-cause mortality for sleep duration <7 hours and >8 hours were 1.50 (1.08-2.10) and 1.36 (0.89-2.08), respectively. The corresponding HR (95%CI) for cardiovascular mortality were 1.56 (0.72-3.37) and 1.56 (0.66-3.65). Non-restorative sleep and restless legs syndrome were associated with increased risk for all-cause mortality (1.63 [1.13-2.35], and 1.69 [1.04-275], respectively).

Corresponding Author: Ana C. Ricardo, MD, MPH, University of Illinois at Chicago, Department of Medicine, Division of Nephrology, 820 S. Wood St. 418W CSN, MC 793, Chicago, Illinois 60612-7315, United States, Phone: 1 (312) 996-8392, Fax: 1 (312) 996-7378, aricar2@uic.edu. Shared first author.

Financial Disclosure: The authors declare that they have no relevant financial interest.

Publisher's Disclaimer: This is a PDF file of an unedited manuscript that has been accepted for publication. As a service to our customers we are providing this early version of the manuscript. The manuscript will undergo copyediting, typesetting, and review of the resulting proof before it is published in its final citable form. Please note that during the production process errors may be discovered which could affect the content, and all legal disclaimers that apply to the journal pertain.

Conclusion—In adults with CKD, short sleep duration, nonrestorative sleep and restless legs syndrome are associated with increased risk of death. These findings underscore the importance of promoting adequate sleep in patients with CKD, and the need for future studies evaluating the impact of sleep interventions in this population.

Introduction

Sleep is an essential state of rest for the well-being of the mind and body, but sleep curtailment has become a common, often voluntary behavior in modern society.¹ In general populations, impaired sleep has been found to be associated with poor health outcomes including death.^{2,3} In addition, there is increasing evidence for an association between both short and long duration of habitual sleep, as well as impaired sleep quality, with prevalence and severity of major chronic diseases, including hypertension, diabetes and cardiovascular disease.⁴⁻⁷

It is estimated that among people with chronic kidney disease (CKD) the prevalence of sleep disturbances can be as high as 80%.⁸ In an analysis of the National Health and Nutrition Examination Survey (NHANES) 2005-2008, Plantinga et al found that the prevalence of inadequate sleep (defined as < 6 hours per night) was higher in individuals with mild CKD than in those with no CKD.⁹ However, the impact of sleep duration and sleep quality on clinical outcomes in individuals with CKD is not well understood. For this reason, we conducted a study to assess the association of sleep duration, sleep symptoms and disorders with all-cause and cardiovascular mortality in U.S. adults with CKD using data from NHANES 2005-2008.

Materials and Methods

Study Population

NHANES is a stratified, clustered, multistage probability sample survey of the civilian, noninstitutionalized U.S. population, conducted by the National Center for Health Statistics (NCHS) of the U.S. Centers for Disease Control and Prevention, with oversampling of non-Hispanic black and Mexican-American persons.¹⁰ The survey consists of a standardized in-home interview followed by physical examination, as well as blood and urine collection at a mobile examination center (MEC). Survey protocol was approved by the NCHS Institutional Review Board and is adherent to the Declaration of Helsinki. All participants provided informed consent. This analysis was limited to NHANES 2005-2008 participants who met the inclusion criteria (18 years or older, non-pregnant, and had available serum creatinine and urine albumin and creatinine measurements) and the study definition of CKD.

Measurements and Definitions

Chronic Kidney Disease—Serum and urine creatinine were measured using the modified kinetic Jaffé method. Urine albumin was measured using a solid-phase fluorescent immunoassay. Urine albumin and creatinine concentrations were measured in one random urine sample. CKD was defined by either an estimated glomerular filtration rate (eGFR) <60 ml/min/1.73m², using the CKD Epidemiology Collaboration (CKD-EPI) creatinine

equation,¹¹ or the presence of albuminuria (urine albumin to creatinine ratio [UACR] 30 mg/g).

Sleep—During the home interview, through a computer- assisted personal interviewing system, NHANES 2005-2008 participants answered questions regarding sleep habits and sleep-related problems from two validated instruments: the Sleep Heart Health Study Sleep Habits Questionnaire¹² and the Functional Outcomes of Sleep Questionnaire.^{13,14} For this study we used selected questions as described herein. Sleep duration was ascertained using the following question: “How much sleep do you usually get at night on weekdays or workdays?” We classified total hours of sleep as <7, 7-8 or >8.² The items used to ascertain the presence of sleep symptoms were: 1) difficulty falling asleep, “In the past month, how often did you have trouble falling asleep;” 2) difficulty staying asleep, “In the past month, how often did you wake up during the night and had trouble getting back to sleep;” 3) daytime sleepiness, “In the past month, how often did you feel excessively or overly sleepy during the day;” and 4) non-restorative sleep, “In the past month, how often did you feel unrested during the day, no matter how many hours of sleep you had.” Participants were asked to choose from one of the following options: Never, rarely (1 time a month), sometimes (2-4 times a month), often (5-15 times a month), almost always (16-30 times a month), refused or don't know. Sleep symptoms were considered to be present if reported “often” or more (at least 5 times a month). The presence of restless legs was also self-reported using the questions: “Have you ever been told by a doctor or other health professional that you have a sleep disorder?” if yes, “What was the sleep disorder?” The possible answers were “sleep apnea”, “insomnia”, “restless legs”, “other”, “refused” and “don't know”.

Covariates—Race/ethnicity was categorized as non-Hispanic white, non-Hispanic black, Mexican American, or other. In these analyses income was classified as annual family income <20,000 or 20,000 U.S. dollars, and educational attainment as less than high school or high school. Participants were considered to have health insurance if they self-reported coverage by any health insurance plan. Participants were classified as current or past/never smoker based on responses to the questions “Have you smoked at least 100 cigarettes during your entire life?” and “Do you smoke cigarettes now?” Participants had three BP measurements at the MEC in the sitting position, after 5 minutes of rest, using a standardized protocol.¹⁵ The averages of all systolic BP available readings are reported here. Hypertension was defined as systolic BP 140 mm Hg or diastolic BP 90 mm Hg or the self-reported use of antihypertensive medications. Diabetes was defined as a history of diabetes, self-reported use of insulin or other medication to treat diabetes, a fasting blood glucose 126 mg/dL, or a random blood glucose 200 mg/dL. The presence of congestive heart failure was ascertained using the following question, “has a doctor or other health professional ever told you that you had congestive heart failure”. The use of medications for sleep was ascertained using the following question: In the past month, how often did you take sleeping pills or other medication to help you sleep? The possible answers were never, rarely (1 time/month), sometimes (2-4 times/month), often (5-15 times/month) or almost always (16-30 times/month); participant who answered 5 times/month were classified as sleeping pills users. Height and weight were measured by trained NHANES staff. Body

mass index (BMI) was calculated as weight in kilograms divided by height in meters squared. The presence of depressive symptoms was defined as a Patient Health Questionnaire (PHQ-9) ≥ 10 .¹⁶

Outcome Ascertainment

We used the NHANES Linked Mortality File, which provides follow-up data on vital status from the date of the NHANES 2005-2008 survey participation through the date of death or December 31, 2011. Vital status was ascertained by the NCHS through a probabilistic match between NHANES 2005-2008 participants and National Death Index death certificate records.¹⁷ Participants who were not matched with any death records were considered to be alive through the follow-up period. Cause of death was assigned by the NCHS based on the International Classification of Diseases, 10th Revision. For this study, cardiovascular mortality was defined as death due to diseases of the heart, essential hypertension and hypertensive kidney disease, cerebrovascular disease, atherosclerosis, and other diseases/disorders of the circulatory system (codes I00-I99).¹⁸

Statistical methods

NCHS recommendations were followed to account for stratification and clustering of the survey design, as well as oversampling of ethnic minorities and elderly persons.¹⁹ Continuous variables were expressed as means (standard error) or medians (interquartile range) if not normally distributed; and categorical variables as weighted percentage. Chi-squared and student's t tests were used to compare categorical and continuous variables, respectively. Cox proportional hazards models were used to determine the association between sleep duration and symptoms and all-cause and cardiovascular mortality adjusting for important covariates (age, gender, race/ethnicity, income, education, diabetes, hypertension, congestive heart failure, sleeping pill use, smoking, eGFR, albuminuria, BMI and depression symptoms) which were chosen based on prior literature.^{3,20-22} We evaluated age, gender and diabetes as potential effect modifiers of the association between sleep variables and mortality by adding an interaction term between the corresponding sleep variable and each of the potential effect modifiers to the fully adjusted model. All tests were two-sided, and $p < 0.05$ was considered significant for hypothesis testing. The proportional hazards assumption of the Cox models was examined using Schoenfeld residuals, which showed no significant departure from proportionality over time ($p > 0.05$).²³ All statistical analyses were done using SAS 9.3 (Cary, NC).

Results

Participant Characteristics

Of the 11,791 NHANES 2005-2008 adult (age ≥ 18 years) participants who were examined, 1,395 did not have data on serum creatinine or urine albumin to creatinine ratio. Pregnant women and individuals with $eGFR < 15$ ml/min/1.73 m² were excluded. Among the remaining participants, 1,820 met our definition of CKD. Of those, we excluded participants due to missing data on sleep duration or symptoms ($n=7$), income ($n=64$), diabetes ($n=31$), smoking ($n=69$), body mass index ($n=35$), congestive heart failure ($n=12$) or other covariate ($n=150$). Therefore, our final analytic sample included 1,452 individuals. Compared with

participants who were included in analyses, those who were excluded due to missing covariate data were more likely to be younger (mean age 56.1 vs. 60.5 years, $p=0.005$); of “other” racial/ethnic background (15.6 vs. 7.5%, $p<0.001$); and to have higher eGFR (81.7 vs. 74.3 ml/min/1.73 m², $p<0.001$). There were no differences in gender or urine albumin to creatinine ratio distribution.

The mean sleep duration was 7 hours. Approximately 35.4% of individuals reported sleeping <7 hours, 54.7% 7-8 hours, and 9.9% > 8 hours. Demographic and clinical characteristics are presented overall and by sleep duration category in Table 1. Mean age was 60.4 years, 58.6% of participants were women, and 74.6% were non-Hispanic whites. Compared to individuals sleeping 7-8 hours, individuals reporting <7 hours of sleep were more likely to be younger (58.5 versus 60.5 years), non-Hispanic black (18.3% versus 7.6%), and have less than a high school education (26.9% versus 22.7%). In addition, individuals reporting <7 hours sleep were more likely to have a BMI of ≥ 30 kg/m² (46.2% versus 39.0%), have diabetes (32.5% versus 24.1%), depressive symptoms (PHQ-9 score ≥ 10 9.5% versus 4.4%), and albuminuria >300 mg/g (13.1% versus 6.9%): they were also more likely to report sleeping medication use (17.8% versus 10.0%). Compared to individuals reporting 7-8 hours of sleep, individuals who reported >8 hour sleep were older (66.7 versus 60.5 years), more likely to have <high school education (38.9% versus 22.7%), have health insurance (92.8% versus 86.4%), diabetes (29.7% versus 24.1%), congestive heart failure (11.0% versus 5.9%), lower eGFR (68.0 versus 73.1 ml/min/1.73m²), and albuminuria >300 mg/g (9.6% versus 6.9%).

In Table 2, demographic and clinical characteristics are presented by presence or absence of sleep symptoms and sleep disorders. Individuals who reported sleep symptoms were more likely to be women, current smokers, and to have a PHQ-9 score ≥ 10 .

Association of Sleep Duration with Mortality

During a median follow-up of 4.4 years, 234 deaths occurred, of which 75 were due to a cardiovascular cause. All-cause mortality rates and hazard ratios (HR) with 95% confidence intervals by category of sleep duration are summarized in Figure 1. After adjustment for sociodemographic and clinical factors, self-reported sleep duration <7 hours was associated with a 50% increased risk for all-cause mortality as compared to individuals reporting 7-8 hours of sleep (HR 1.50, 95% CI 1.08-2.10). Self-reported sleep duration >8 hours was associated with a nonsignificant increased risk for all-cause mortality compared to 7-8 hours (HR 1.36, 95% CI 0.89-2.08). Similarly, compared with self-reported sleep duration of 7-8 hours, sleep duration <7 hours and >8 hours were associated with non-statistically significant increased risk for cardiovascular death (HR [95% CI], 1.56 [0.72-3.37], and 1.56 [0.66-3.65]), respectively, Figure 2). We found no evidence of effect modification by age, gender or diabetes.

Association of Sleep-Related Symptoms with Mortality

Non-restorative sleep and restless legs syndrome were each associated with an increased risk for all-cause death (HR [95% CI], 1.63 [1.13-2.35], and 1.69 [1.04-2.75] respectively). There was no significant association between the other sleep symptoms or sleep apnea and

all-cause death (Figure 1). Furthermore, there was no significant association between sleep symptoms and cardiovascular death (Figure 2). We found no evidence of effect modification by age, gender or diabetes.

Discussion

In U.S. adults with CKD, self-reported sleep duration of <7 hours was associated with increased risk of all-cause and cardiovascular death compared with individuals reporting sleep duration of 7-8 hours. Additionally, non-restorative sleep and diagnosed restless legs syndrome were associated with increased risk of all-cause death. To our knowledge, this is the first study to examine the association of sleep duration and sleep symptoms with mortality in a representative sample of U.S. adults with non-dialysis dependent CKD.

Several population-based cohort studies have demonstrated elevated risk of all-cause and cardiovascular mortality for individuals reporting short sleep duration.^{20–22,24,25} There is a paucity of data regarding the impact of short sleep on clinical outcomes in CKD. A recent analysis from the Nurses' Health Study found that shorter sleep was associated with faster decline in kidney function but mortality was not evaluated.²⁶ Our findings suggest that the association between short sleep duration and mortality found in the general population may extend to patients with non-dialysis dependent CKD.

A number of mechanisms have been proposed to explain the association between poor sleep duration and adverse outcomes in the general population, including alterations of sympathetic nervous system activity, cortisol release, glucose intolerance, and inflammation.^{8,27–29} The associations of short sleep duration with mortality in CKD may also be reflective of disturbances in circadian rhythm. Circadian misalignment can arise when an individual's sleep is not in synchrony with their endogenous clocks.³⁰ There is growing evidence from animal and human studies suggesting that dysregulation of renal circadian rhythms may be associated with worsening blood pressure and glucose metabolism.³¹ For example, activation of the circadian clock activity in mice leads to salt-sensitive activation, whereas its suppression leads to low blood pressure.³² Future studies are needed to better understand the role of circadian rhythm in CKD outcomes.

We also found significant associations between non-restorative sleep and restless legs syndrome and all-cause mortality which have been observed in general population studies.^{33,34} In the Health Professionals Follow-Up Study, non-restorative sleep was associated with HR 1.24 (95% CI, 1.05–1.46) for all-cause mortality. The potential mechanisms linking non-restorative sleep with mortality are similar to those mentioned above for short sleep. Restless legs syndrome is common in patients with CKD.³⁵ Reasons for this increased prevalence are not understood but it has been postulated that disordered iron metabolism may be an important factor.³⁶ Multiple observational cohort studies of individuals with end-stage renal disease have shown a significant increase in the risk of death among individuals with restless leg syndrome compared with those without the disease.^{37–40} However, less is known about this association in patients with CKD who are not on dialysis.

Our findings of no association between sleep apnea and increased risk of death in individuals with CKD are in contrast with general populations studies which have shown that sleep apnea diagnosed by polysomnography or other objective measure is associated with increased mortality risk.^{41–43} The reason for this discordance might be the self-reported method of sleep apnea ascertainment in our study. It is well established that sleep apnea is often underdiagnosed in the clinical setting,⁴⁴ therefore, underestimation of its prevalence in our study might have limited the power to detect an association between sleep apnea and mortality.

Strengths of our study include the large sample size and prospective study design with median follow-up of 4.4 years and the comprehensive assessment of sleep symptoms. However, several limitations of this study should be taken into account. First, the use of self-reported questionnaires to capture sleep duration may provide an inflated estimate of sleep duration;⁴⁵ in addition, it might underestimate the prevalence of sleep disorders such as sleep apnea which are known to be common in patients with CKD undergoing objective sleep measurements.⁴⁶ Second, misclassification of CKD due to single-measurement of eGFR and UACR is possible, which may lead to misclassification of the selected study sample. Third, our study was underpowered to detect associations with cardiovascular mortality. Finally, individuals with missing serum creatinine or UACR data were excluded from analysis. Therefore, our findings may not be representative of the entire CKD population.

In conclusion, short sleep was associated with an increased risk for all-cause mortality in individuals with CKD. Non-restorative sleep and restless leg symptoms were also associated with an increased risk for all-cause mortality. These findings reinforce the importance of promoting adequate sleep in patients with CKD. Future longitudinal studies with objective measures of sleep duration and symptoms as well as randomized controlled trials investigating the effects of improving sleep duration and symptoms are needed. A better understanding of the effect of sleep duration and sleep symptoms on adverse outcomes among the CKD population could help inform CKD management and lead to improvement of health outcomes.

Acknowledgments

Support: Ana C. Ricardo is funded by the NIDDK K23DK094829 award. Esteban Cedillo-Couvert is funded by the NIDDK Research Supplement to Promote Diversity in Health-Related Research (U01DK060980). James P. Lash is funded by the NIDDK K24DK092290 award. Bharati Prasad is funded by Department of Veterans Affairs 1K2CX001026-01.

References

1. Knutson KL, Van Cauter E. Associations between sleep loss and increased risk of obesity and diabetes. *Ann N Y Acad Sci.* 2008; 1129:287–304. [PubMed: 18591489]
2. Gallicchio L, Kalesan B. Sleep duration and mortality: a systematic review and meta-analysis. *J Sleep Res.* 2009; 18:148–158. [PubMed: 19645960]
3. Empana JP, Dauvilliers Y, Dartigues JF, Ritchie K, Garipey J, Jouven X, Tzourio C, Amouyel P, Besset A, Ducimetiere P. Excessive daytime sleepiness is an independent risk indicator for cardiovascular mortality in community-dwelling elderly: the three city study. *Stroke J Cereb Circ.* 2009; 40:1219–1224.

4. Knutson KL, Van Cauter E, Rathouz PJ, Yan LL, Hulley SB, Liu K, Lauderdale DS. Association between sleep and blood pressure in midlife: the CARDIA sleep study. *Arch Intern Med.* 2009; 169:1055–1061. [PubMed: 19506175]
5. Knutson KL, Ryden AM, Mander BA, Van Cauter E. Role of sleep duration and quality in the risk and severity of type 2 diabetes mellitus. *Arch Intern Med.* 2006; 166:1768–1774. [PubMed: 16983057]
6. Meisinger C, Heier M, Löwel H, Schneider A, Döring A. Sleep duration and sleep complaints and risk of myocardial infarction in middle-aged men and women from the general population: the MONICA/KORA Augsburg cohort study. *Sleep.* 2007; 30:1121–1127. [PubMed: 17910384]
7. Cappuccio FP, D'Elia L, Strazzullo P, Miller MA. Quantity and quality of sleep and incidence of type 2 diabetes: a systematic review and meta-analysis. *Diabetes Care.* 2010; 33:414–420. [PubMed: 19910503]
8. Turek NF, Ricardo AC, Lash JP. Sleep disturbances as nontraditional risk factors for development and progression of CKD: review of the evidence. *Am J Kidney Dis Off J Natl Kidney Found.* 2012; 60:823–833.
9. Plantinga L, Lee K, Inker LA, Saran R, Yee J, Gillespie B, Rolka D, Saydah S, Powe NR. CDC CKD Surveillance Team. Association of sleep-related problems with CKD in the United States, 2005-2008. *Am J Kidney Dis Off J Natl Kidney Found.* 2011; 58:554–564.
10. Centers for Disease Control Prevention, National Center for Health Statistics. [2016 Aug 3] Natl Health Nutr Exam Surv Data 2005-2008. 2016. Internet Available from: <http://www.cdc.gov/nchs/nhanes>
11. Inker LA, Schmid CH, Tighiouart H, Eckfeldt JH, Feldman HI, Greene T, Kusek JW, Manzi J, Van Lente F, Zhang YL, Coresh J, Levey AS. CKD-EPI Investigators. Estimating glomerular filtration rate from serum creatinine and cystatin. *C N Engl J Med.* 2012; 367:20–29. [PubMed: 22762315]
12. Quan SF, Howard BV, Iber C, Kiley JP, Nieto FJ, O'Connor GT, Rapoport DM, Redline S, Robbins J, Samet JM, Wahl PW. The Sleep Heart Health Study: design, rationale, and methods. *Sleep.* 1997; 20:1077–1085. [PubMed: 9493915]
13. Chasens ER, Ratcliffe SJ, Weaver TE. Development of the FOSQ-10: a short version of the Functional Outcomes of Sleep Questionnaire. *Sleep.* 2009; 32:915–919. [PubMed: 19639754]
14. Weaver TE, Laizner AM, Evans LK, Maislin G, Chugh DK, Lyon K, Smith PL, Schwartz AR, Redline S, Pack AI, Dinges DF. An instrument to measure functional status outcomes for disorders of excessive sleepiness. *Sleep.* 1997; 20:835–843. [PubMed: 9415942]
15. NHANES. [2016 Aug 17] Survey Methods Analytic Guidelines [Internet]. Available from: http://www.cdc.gov/nchs/nhanes/survey_methods.htm
16. Kroenke K, Spitzer RL, Williams JB. The PHQ-9: validity of a brief depression severity measure. *J Gen Intern Med.* 2001; 16:606–613. [PubMed: 11556941]
17. National Center for Health Statistics, Office of Analysis and Epidemiology, NCHS. [2016 Sep 14] Linked Mortality Files Matching Methodology, September 2013. Hyattsville, Maryland. 2011. Available at the following address: [Internet]. Available from: http://www.cdc.gov/nchs/data/datalinkage/2011_linked_mortality_file_matching_methodology.pdf
18. Anderson RN, Miniño AM, Hoyert DL, Rosenberg HM. Comparability of cause of death between ICD-9 and ICD-10: preliminary estimates. *Natl Vital Stat Rep Cent Dis Control Prev Natl Cent Health Stat Natl Vital Stat Syst.* 2001; 49:1–32.
19. [2016 Aug 17] Vital and Health Statistics Reports Series 2, Number 161, September 2013 - sr02_161.pdf [Internet]. Available from: http://www.cdc.gov/nchs/data/series/sr_02/sr02_161.pdf
20. Ferrie JE, Shipley MJ, Cappuccio FP, Brunner E, Miller MA, Kumari M, Marmot MG. A prospective study of change in sleep duration: associations with mortality in the Whitehall II cohort. *Sleep.* 2007; 30:1659–1666. [PubMed: 18246975]
21. Hublin C, Partinen M, Koskenvuo M, Kaprio J. Sleep and mortality: a population-based 22-year follow-up study. *Sleep.* 2007; 30:1245–1253. [PubMed: 17969458]
22. Ikehara S, Iso H, Date C, Kikuchi S, Watanabe Y, Wada Y, Inaba Y, Tamakoshi A, JACC Study Group. Association of sleep duration with mortality from cardiovascular disease and other causes for Japanese men and women: the JACC study. *Sleep.* 2009; 32:295–301. [PubMed: 19294949]

23. Schoenfeld DA. Partial residuals for the proportional hazards regression model. *Biometrika*. 1982; 69:239–241.
24. Kripke DF, Langer RD, Elliott JA, Klauber MR, Rex KM. Mortality related to actigraphic long and short sleep. *Sleep Med*. 2011; 12:28–33. [PubMed: 20870457]
25. Vgontzas AN, Liao D, Pejovic S, Calhoun S, Karataraki M, Basta M, Fernández-Mendoza J, Bixler EO. Insomnia with short sleep duration and mortality: the Penn State cohort. *Sleep*. 2010; 33:1159–1164. [PubMed: 20857861]
26. McMullan CJ, Curhan GC, Forman JP. Association of short sleep duration and rapid decline in renal function. *Kidney Int*. 2016; 89:1324–1330. [PubMed: 27165820]
27. Charloux A, Gronfier C, Lonsdorfer-Wolf E, Piquard F, Brandenberger G. Aldosterone release during the sleep-wake cycle in humans. *Am J Physiol*. 1999; 276:E43–49. [PubMed: 9886949]
28. Knutson KL, Spiegel K, Penev P, Van Cauter E. The Metabolic Consequences of Sleep Deprivation. *Sleep Med Rev*. 2007; 11:163–178. [PubMed: 17442599]
29. Brianc¸on-Marjollet A, Weiszenstein M, Henri M, Thomas A, Godin-Ribuot D, Polak J. The impact of sleep disorders on glucose metabolism: endocrine and molecular mechanisms. *Diabetol Metab Syndr* [Internet]. 2015; 7 [2016 Sep 20] Available from: <http://www.ncbi.nlm.nih.gov/pmc/articles/PMC4381534/>.
30. Baron KG, Reid KJ. Circadian misalignment and health. *Int Rev Psychiatry Abingdon Engl*. 2014; 26:139–154.
31. Marcheva B, Ramsey KM, Peek CB, Affinati A, Maury E, Bass J. Circadian clocks and metabolism. *Handb Exp Pharmacol*. 2013:127–155.
32. Zuber AM, Centeno G, Pradervand S, Nikolaeva S, Maquelin L, Cardinaux L, Bonny O, Firsov D. Molecular clock is involved in predictive circadian adjustment of renal function. *Proc Natl Acad Sci U S A*. 2009; 106:16523–16528. [PubMed: 19805330]
33. Li Y, Zhang X, Winkelman JW, Redline S, Hu FB, Stampfer M, Ma J, Gao X. Association between insomnia symptoms and mortality: a prospective study of U.S. men. *Circulation*. 2014; 129:737–746. [PubMed: 24226807]
34. Molnar MZ, Lu JL, Kalantar-Zadeh K, Kovesdy CP. Association of incident restless legs syndrome with outcomes in a large cohort of US veterans. *J Sleep Res*. 2016; 25:47–56. [PubMed: 26377112]
35. Novak M, Winkelman JW, Unruh M. Restless Legs Syndrome in Patients With Chronic Kidney Disease. *Semin Nephrol*. 2015; 35:347–358. [PubMed: 26355253]
36. Trenkwalder C, Paulus W. Restless legs syndrome: pathophysiology, clinical presentation and management. *Nat Rev Neurol*. 2010; 6:337–346. [PubMed: 20531433]
37. Lin CH, Sy HN, Chang HW, Liou HH, Lin CY, Wu VC, Wu SL, Chang CC, Chiu PF, Li WY, Lin SY, Wu KD, Chen YM, Wu RM. Restless legs syndrome is associated with cardio/cerebrovascular events and mortality in end-stage renal disease. *Eur J Neurol*. 2015; 22:142–149. [PubMed: 25142748]
38. Winkelman JW, Chertow GM, Lazarus JM. Restless legs syndrome in end-stage renal disease. *Am J Kidney Dis Off J Natl Kidney Found*. 1996; 28:372–378.
39. Unruh ML, Levey AS, D'Ambrosio C, Fink NE, Powe NR, Meyer KB. Choices for Healthy Outcomes in Caring for End-Stage Renal Disease (CHOICE) Study: Restless legs symptoms among incident dialysis patients: association with lower quality of life and shorter survival. *Am J Kidney Dis Off J Natl Kidney Found*. 2004; 43:900–909.
40. La Manna G, Pizza F, Persici E, Baraldi O, Comai G, Cappuccilli ML, Centofanti F, Carretta E, Plazzi G, Coli L, Montagna P, Stefoni S. Restless legs syndrome enhances cardiovascular risk and mortality in patients with end-stage kidney disease undergoing long-term haemodialysis treatment. *Nephrol Dial Transplant Off Publ Eur Dial Transpl Assoc - Eur Ren Assoc*. 2011; 26:1976–1983.
41. Shah NA, Yaggi HK, Concato J, Mohsenin V. Obstructive sleep apnea as a risk factor for coronary events or cardiovascular death. *Sleep Breath Schlaf Atm*. 2010; 14:131–136.
42. Yaggi HK, Concato J, Kernan WN, Lichtman JH, Brass LM, Mohsenin V. Obstructive sleep apnea as a risk factor for stroke and death. *N Engl J Med*. 2005; 353:2034–2041. [PubMed: 16282178]

43. Marshall NS, Wong KKH, Liu PY, Cullen SRJ, Knuiaman MW, Grunstein RR. Sleep apnea as an independent risk factor for all-cause mortality: the Busselton Health Study. *Sleep*. 2008; 31:1079–1085. [PubMed: 18714779]
44. Senthilvel E, Auckley D, Dasarathy J. Evaluation of sleep disorders in the primary care setting: history taking compared to questionnaires. *J Clin Sleep Med JCSM Off Publ Am Acad Sleep Med*. 2011; 7:41–48.
45. Lauderdale DS, Knutson KL, Yan LL, Liu K, Rathouz PJ. Self-reported and measured sleep duration: how similar are they? *Epidemiol Camb Mass*. 2008; 19:838–845.
46. Sakaguchi Y, Shoji T, Kawabata H, Nihata K, Suzuki A, Kaneko T, Okada N, Isaka Y, Rakugi H, Tsubakihara Y. High prevalence of obstructive sleep apnea and its association with renal function among nondialysis chronic kidney disease patients in Japan: a cross-sectional study. *Clin J Am Soc Nephrol CJASN*. 2011; 6:995–1000. [PubMed: 21415314]




Figure 1. Event rates per 100 person-years and adjusted hazard ratios (95% confidence intervals) for all-cause mortality according to sleep duration, symptoms and disorders. Hazard ratios are adjusted for age, gender, race/ethnicity, income, education, diabetes, body mass index, hypertension, smoking, depressive symptoms, eGFR and albuminuria.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript




Figure 2. Event rates per 100 person-years and adjusted hazard ratio (95% confidence intervals) for cardiovascular mortality according to sleep duration, symptoms and disorders. Hazard ratios are adjusted for age, gender, race/ethnicity, income, education, diabetes, body mass index, hypertension, smoking, depressive symptoms, eGFR and albuminuria

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1

Characteristics of Individuals with CKD Overall and Stratified by Sleep Duration.

	Overall (N = 1,452)	<7 hours (N = 543)	7-8 hours (N = 752)	>8 hours (N = 157)
Age, years	60.4 ± 0.8	58.5 ± 1.0	60.5 ± 1.1	66.7 ± 1.9 [*]
18-44	22.2%	25.8%	21.3%	14.3% [*]
45-64	28.5%	30.9%	29.3%	16.0%
>65	49.2%	43.3%	49.4%	69.6%
Female Gender	58.6%	56.1%	59.7%	61.4%
Race				
Non-Hispanic White	74.6%	66.5%	78.6%	81.1% [*]
Non-Hispanic Black	11.3%	18.3%	7.6%	6.6%
Mexican American	6.7%	5.9%	7.3%	5.9%
Other	7.5%	9.3%	6.6%	6.4%
Household Income < \$20,000/year	23.3%	25.9%	21.4%	24.6%
Education < High School	25.8%	26.9%	22.7%	38.9% [*]
No health insurance	11.6%	9.6%	13.6%	7.2% [*]
Current Smoker	17.1%	20.7%	15.2%	14.3%
Hypertension	56.8%	60.9%	53.8%	58.9%
Diabetes	27.6%	32.5%	24.1%	29.7% [*]
Congestive heart failure	7.2%	8.3%	5.9%	11% [*]
Sleeping pill use	12.5%	17.8%	10.0%	7.8%
Depression (PHQ-9 Score ≥ 10)	6.2%	9.5%	4.4%	3.8% [*]
Systolic BP (mm Hg)	132 ± 1	133 ± 1	132 ± 1	131 ± 2
Diastolic BP (mm Hg)	69 ± 1	70 ± 1	69 ± 1	66 ± 2
BMI ≥ 30 kg/m ²	40.9%	46.2%	39.0%	32.2% [*]
HbA1C (%)	6.0 ± 0.1	6.1 ± 0.1	5.9 ± 0.1	6.0 ± 0.2
TC ≥ 200 mg/dL	44.6%	44.7%	45.8%	37.9%
LDL-C ≥ 100 mg/dL	56.5%	60.8%	55.7%	46.9%
eGFR (ml/min/1.73m ²)	74.3 ± 1.2	78.0 ± 1.7	73.1 ± 1.6	68.0 ± 2.4 [*]
Urine ACR				
<30 mg/g	33.3%	25.2%	36.9%	42.4% [*]
30-300 mg/g	57.4%	61.7%	56.3%	47.9%
>300 mg/g	9.3%	13.1%	6.9%	9.6%

^{*}
p < .05

Values are expressed as weighted mean ± standard error or weighted percentage.

Abbreviations: ACR, albumin to creatinine ratio; BMI, body mass index; BP, blood pressure; eGFR, estimated glomerular filtration rate; LDL-C, low-density lipoprotein cholesterol; PHQ-9, Patient Health Questionnaire for depression screening; TC, total cholesterol.

Table 2
Characteristics of Individuals with CKD Overall and Stratified by Sleep Symptoms and Disorders

Characteristic	Difficulty Falling Asleep		Difficulty Staying Asleep		Daytime Sleepiness		Non-Restorative Sleep		Restless Legs Syndrome		Sleep Apnea	
	Yes N = 233	No N = 1,219	Yes N = 302	No N = 1,150	Yes N = 242	No N = 1,220	Yes N = 302	No N = 1,150	Yes N = 195	No N = 1,257	Yes N = 93	No N = 1,359
Weighted %	16.9%	83.1%	21.5%	79.5%	19.4%	80.6%	25.3%	74.7%	12.5%	87.5%	7.1%	92.9%
Age, years	59.8 ± 1.4	60.5 ± 0.9	62.2 ± 1.3	59.9 ± 0.9	59.3 ± 1.4	60.7 ± 1.0	56.8 ± 1.2	61.6 ± 1.0*	62.7 ± 1.4	60.1 ± 0.8	58.9 ± 1.8	60.5 ± 0.8*
18-44	22.3%	22.2%	17.7%	23.5%	24.2%	21.7%	27.0%	20.6%	18.7%	22.7%	22.0%	22.2%*
45-64	27.5%	28.7%	29.9%	28.2%	28.9%	28.4%	32.3%	27.3%	31.9%	28.1%	37.9%	27.8%
<65	50.2%	49.0%	52.5%	48.4%	46.9%	50.0%	40.7%	52.1%	49.4%	49.2%	40.1%	49.9%
Female Gender	74.1%	55.4%*	66.5%	56.4%*	70.0%	55.8%*	68.2%	55.3%*	65.0%	57.6%	48.1%	59.1%
Race/Ethnicity												
Non-Hispanic white	75.7%	74.3%	81.8%	72.6%*	81.4%	73.0%*	79.3%	72.9%*	77.9%	74.1%	77.8%	74.3%
Non-Hispanic black	10.2%	11.5%	9.1%	11.9%	8.6%	11.9%*	8.2%	12.3%*	8.2%	11.7%	12.5%	11.2%
Mexican American	3.6%	7.3%*	5.3%	7.0%	4.6%	7.2%	4.5%	7.4%*	7.4%	6.6%	2.8%	7.0%
Other	10.5%	6.9%	3.8%	8.5%*	5.4%	8.0%	7.9%	7.4%	6.5%	7.6%	6.9%	7.5%
Annual Income <\$20,000	27.5%	22.4%	25.3%	22.7%	28.3%	22.1%*	25.6%	22.5%	32.8%	21.9%*	17.5%	23.7%*
Education <High School	29.2%	25.1%	27.8%	25.2%	29.2%	25.0%	22.6%	26.9%	32.2%	24.9%*	17.8%	26.4%*
No health insurance	12.6%	11.4%	10.7%	11.8%	10.1%	11.9%	11.9%	11.5%	12.2%	11.5%	4.4%	12.1%
Current smoker	26.2%	15.2%*	24.8%	15.0%*	23.3%	15.6%*	25.6%	14.2%*	26.3%	15.8%*	16.7%	17.1%
Hypertension	60.6%	56.1%	62.5%	55.3%*	64.5%	55.0%*	60.4%	55.6%	75.2%	54.2%*	78.7%	55.2%*
Diabetes	29.6%	27.2%	28.0%	27.5%	34.2%	26.0%*	33.4%	25.7%	35.7%	26.6%*	49.0%	26.0%*
Congestive heart failure	7.2%	7.2%	10.2%	6.4%*	9.3%	6.7%	8.5%	6.8%	12.1%	6.5%*	16.1%	6.6%*
Sleeping pill use	25.9%	8.9%*	20.3%	9.5%*	21.4%	9.4%*	22.2%	8.2%*	24.8%	9.9%*	16.2%	11.4%
Depression (PHQ-9 10)	17.8%	3.8%*	13.8%	4.0%*	17.6%	3.4%*	14.4%	3.4%*	17.0%	4.6%*	20.8%	5.0%*
Systolic BP (mm Hg)	134 ± 2	132 ± 1	135 ± 1	131.4 ± 0.7*	132 ± 1	133 ± 1	132 ± 1	132 ± 1	134 ± 2	132 ± 1	130 ± 2	133 ± 1*
Diastolic BP (mm Hg)	70 ± 2	69 ± 1	70 ± 1	69.1 ± 0.6	71 ± 1	68.9 ± 0.7	71 ± 1	69 ± 1	68 ± 1	70 ± 1	71 ± 1	69 ± 1
BMI 30 kg/m ²	42.1%	40.6%	40.4%	41.1%	44.8%	39.9%	45.4%	39.3%	51.2%	39.4%*	84.7%	37.5%*

Characteristic	Difficulty Falling /Asleep		Difficulty Staying Asleep		Daytime Sleepiness		Non-Restorative Sleep		Restless Legs Syndrome		Sleep Apnea	
	Yes N =	No N =	Yes N =	No N =	Yes N =	No N =	Yes N =	No N =	Yes N =	No N =	Yes N =	No N =
	233	1,219	302	1,150	242	1,220	302	1,150	195	1,257	93	1359
Weighted %	16.9%	83.1%	21.5%	79.5%	19.4%	80.6%	25.3%	74.7%	12.5%	87.5%	7.1%	92.9%
HbA1C (%)	6.0 ± 0.1	6.0 ± 0.1	5.9 ± 0.1	6.0 ± 0.1	6.1 ± 0.1	6.0 ± 0.1	6.1 ± 0.1	5.9 ± 0.0	6.2 ± 0.1	6.0 ± 0.0 *	6.1 ± 0.1	6.0 ± 0.0
TC 200 mg/dL	51.7%	43.2% *	46.9%	44.0%	48.2%	43.8%	48.3%	43.3%	42.6%	44.9%	27.7%	45.8%
LDL-C 100 mg/dL	63.5%	55.3%	56.1%	56.7%	56.1%	56.7%	57.6%	56.2%	53.5%	57.0%	33.3%	58.8%
eGFR (ml/min/1.73m ²)	75.7 ± 2.4	74.0 ± 1.4	71.6 ± 2.3	75.1 ± 1.3	75.1 ± 2.5	74.1 ± 1.6	78.3 ± 2.0	73.0 ± 1.5 *	70.2 ± 2.9	74.9 ± 1.2	71.0 ± 3.7	74.6 ± 1.3
Urine ACR												
<30 mg/g	29.7%	34.0%	34.8%	32.9%	31.1%	33.8%	30.0%	34.4%	37.8%	32.6%	25.3%	33.9% *
30-300 mg/g	58.5%	57.1%	54.7%	58.1%	55.9%	57.7%	56.1%	57.8%	51.6%	58.2%	59.4%	57.2%
>300 mg/g	11.8%	8.9%	10.6%	9.0%	13.1%	8.5%	13.9%	7.8% *	10.6%	9.2%	15.3%	8.9%

* p < .05

Values are expressed as weighted mean ± standard error or weighted percentage.

Abbreviations: ACR, albumin to creatinine ratio; BMI, body mass index; BP, blood pressure; CVD, cardiovascular disease; eGFR, estimated glomerular filtration rate; LDL-C, low-density lipoprotein cholesterol; PHQ-9, Patient Health Questionnaire for depression screening; TC, total cholesterol.