



Published in final edited form as:

Pediatrics. 2018 April ; 141(4): . doi:10.1542/peds.2017-3594.

Electronic Cigarette Use and Progression from Experimentation to Established Smoking

Benjamin W. Chaffee, DDS MPH PhD^{a,b}, Shannon Lea Watkins, PhD^{a,c}, and Stanton A. Glantz, PhD^{a,c,d}

^aCenter for Tobacco Control Research and Education, University of California San Francisco

^bDepartment of Preventive and Restorative Dentistry, University of California San Francisco

^cDepartment of Medicine, University of California San Francisco

^dPhilip R. Lee Institute for Health Policy Studies, University of California San Francisco

Abstract

Background—Existing longitudinal studies consistently show that never-smoking adolescents who try electronic cigarettes are at increased risk of subsequent conventional cigarette smoking. This study aimed to evaluate the association between e-cigarette use and progression to established smoking among adolescents who had already tried cigarettes at baseline.

Methods—Among youth participants (age 12–17) in the nationally representative Population Assessment of Tobacco and Health survey who had smoked a cigarette (1 puff) but not yet smoked 100 cigarettes (N=1,295), we examined three outcomes at 1-year follow-up as a function of baseline e-cigarette use: 1) having smoked 100 cigarettes (established smoking), 2) smoking during the past 30 days, and 3) both having smoked 100 cigarettes and past 30-day smoking (current established smoking). Survey-weighted multivariable logistic regression models were fitted to obtain odds ratios (OR) and 95% confidence intervals (CI) adjusted for socio-demographic variables and other smoking risk factors.

Results—Versus e-cigarette never use, having ever used e-cigarettes was positively associated with progression to established cigarette smoking (19.3% vs. 9.7%), past 30-day smoking (38.8% vs. 26.6%), and current established smoking (15.6% vs. 7.1%). In adjusted models, e-cigarette ever use positively predicted current established smoking (OR: 1.80, CI: 1.04–3.12) but did not reach statistical significance (threshold: $\alpha=0.05$) for established smoking (OR: 1.57, CI: 0.99–2.49) and past 30-day smoking (OR: 1.32, CI: 0.99–1.76).

Corresponding Author: Benjamin W. Chaffee, DDS MPH PhD, University of California San Francisco, 3333 California St., San Francisco, CA 94118, 1 (415) 476 - 9226, benjamin.chaffee@ucsf.edu.

Financial Disclosure: The authors have no financial relationships relevant to this article to disclose.

Conflicts of Interest: The authors have no conflicts of interest relevant to this article to disclose.

Contributors Statements:

Dr. Chaffee contributed to the design and conceptualization of the study and analysis plan, conducted statistical analyses, and prepared the initial manuscript draft.

Dr. Watkins contributed to the design and conceptualization of the study and analysis plan, and conducted statistical analyses.

Dr. Glantz contributed to the design and conceptualization of the study and analysis plan.

All authors revised and reviewed the manuscript and approved the final manuscript as submitted and agree to be accountable for all aspects of the work.

Conclusion—Among adolescent cigarette experimenters, using e-cigarettes was positively and independently associated with progression to current established smoking, suggesting that e-cigarettes do not divert from, and may encourage, cigarette smoking in this population.

INTRODUCTION

Electronic cigarettes (e-cigarettes) are increasingly popular among youth; from 2014 to 2016, more U.S. middle and high school students used e-cigarettes than any other tobacco product, including conventional cigarettes.¹ All currently available longitudinal studies have shown that among never-smoking adolescents and young adults, e-cigarette use is associated with subsequent cigarette smoking.^{2–6} This association was shown in studies taking place in California,³ Hawaii,² and the Mid-Atlantic region,⁷ as well as in nationally-representative U.S. samples^{6, 8} and in Canada⁴ and the United Kingdom.⁵ Seven of these studies were summarized in a recent meta-analysis, showing more than three-fold increase in the risk of cigarette smoking initiation when comparing youth e-cigarette ever users to never users.⁹ While this association between e-cigarettes and smoking initiation has been consistent across the literature and could be explained using a proposed "catalyst" model,¹⁰ some have argued that the relationship partly reflects a shared propensity for experimentation with different nicotine-containing products.¹¹

Studies of baseline cigarette never-users may include many individuals at low risk of smoking initiation. In contrast, youth who have already begun cigarette experimentation represent a population at high risk of progression to greater levels of cigarette use, later in adolescence and into adulthood. While smoking even one cigarette is concerning, becoming an established smoker in adolescence is of substantial clinical and public health concern and strongly associated with continuing to smoke regularly.¹² Therefore, the present investigation considers high-risk youth, as evident by having already tried smoking (ever smoked 1 puff) but not yet smoked 100 cigarettes, and evaluates whether e-cigarette use in this population predicts progression from experimentation to established cigarette smoking one year later.

In a previous cross-sectional analysis of the 2011 and 2012 National Youth Tobacco Surveys (NYTS), among youth who had ever smoked a cigarette, ever use of e-cigarettes was associated with being an established smoker (lifetime smoked 100 cigarettes), including after adjusting for socio-demographic variables.¹³ However, the cross-sectional design of that analysis precluded causal conclusions, due to uncertain temporal sequencing between e-cigarette use and established smoking.

The present study used the nationally representative Population Assessment of Tobacco and Health (PATH) Study¹⁴ Waves 1 (2013–14) and 2 (2014–15) to examine these same relationships prospectively. We hypothesized that among PATH youth participants who had already tried cigarette smoking but not yet smoked a total of 100 cigarettes, use of e-cigarettes would be positively associated with becoming an established cigarette smoker within 1 year.

POPULATION AND METHODS

The PATH Study selected participants using a four-stage, stratified probability design with oversampling for tobacco users, African Americans, and young adults (ages 18–24). The PATH Youth sample consisted of adolescents (up to 2 per household) whose parents were selected for the PATH Adult sample.¹⁴ The PATH Youth Study enrolled 13,651 U.S. adolescents age 12–17 years at baseline (2013–2014) with 87.9% retention (unweighted) at Wave 2 (2014–2015).

In-home, in-person computer-assisted interviews were conducted in administering the PATH questionnaire. In separate sections, participants were asked about their tobacco use (e.g., ever use, number of lifetime uses, and number of days used in the past 30 days) for 8 types of tobacco and nicotine-containing products, including cigarettes and e-cigarettes. Tobacco use questions were repeated during the Wave 2 interview, including for individuals who reached age 18 before follow-up and were therefore administered the Wave 2 Adult questionnaire.

The present analysis included youth who had smoked ≥ 1 cigarette puff but had not yet smoked 100 cigarettes at baseline (smoking experimenters) with known smoking status at follow-up (N=1,295). We examined three outcomes at follow-up as a function of baseline e-cigarette use: 1) having smoked ≥ 100 cigarettes (established smoking), 2) smoking during the past 30 days, and 3) both having smoked ≥ 100 cigarettes and past 30-day smoking (current established smoking). We categorized e-cigarette use in two ways: 1) ever use or never use, and 2) never use, non-past 30-day use (former use), or past 30-day use.

Logistic regression models (Stata 14, StataCorp LLC, College Station, TX) were used to adjust for hypothesized confounding variables in three stages. First, six separate unadjusted models were fitted to cover each combination of independent variable (Wave 1 e-cigarette never/ever use and e-cigarette never/former/past 30-day use) and dependent variable (Wave 2 established smoking, current smoking, and current established smoking). In the second stage, all models added gender, age (in years), and race/ethnicity (Hispanic/Latinx, non-Hispanic White, non-Hispanic Black, other) as covariables, matching the confounders used in a previous cross-sectional analysis of NYTS data.¹³ In the third stage, additional adjusted models also included parent education (≥ 1 parent with bachelor degree or above), urban residence (based on sampling units), household tobacco use (lives with ≥ 1 tobacco user), alcohol ever use, tobacco advertisement receptivity¹⁵ (can recall brand of favorite advertisement), sensation-seeking score (scale from 3–15), cigarette warning label exposure (Likert-type scale), interview time of year (summer vs. all other months), and ever use of any other tobacco product (i.e., cigars, pipes, hookah, bidis, kreteks, snus, dissolvable tobacco, and conventional moist snuff or chewing smokeless tobacco). Sensation-seeking score was a composite of three Likert-type items (liking frightening things, willingness to break rules, and preference for exciting and unpredictable friends) and has been shown to correlate with youth tobacco use.¹⁶ Interview time of year was included because, for youth, the scholastic calendar may play a role both in opportunity and social pressure to experiment with tobacco products.

All models were weighted for sampling design and non-response using balanced repeated replication to be representative of the Wave 1 target population.¹⁷ Multiple imputation was performed for missing observations (0.7% of data), with variance estimates adjusted accordingly.

An Institutional Review Board at the University of California San Francisco reviewed and designated the study protocol exempt for this analysis of de-identified survey data. The PATH study protocol received a National Institutes of Health Certificate of Confidentiality and approval from the Westat Institutional Review Board. Parental consent was requested on behalf of participating youth. Youth who completed the questionnaire were given \$25.

RESULTS

Among baseline cigarette experimenters (mean age: 15.5 years, 48.3% female), having ever used e-cigarettes was positively associated with progression to established cigarette smoking in Wave 2 (Table 1). Compared to e-cigarette never users, e-cigarette ever users were twice as likely to report Wave 2 established smoking (19.3% vs. 9.7%; $P<0.001$) and current established smoking (15.6% vs. 7.1%; $P<0.001$) and more likely to report past 30-day smoking (38.8% vs. 26.6%; $P<0.001$).

In models adjusted for gender, age, and race/ethnicity (Table 1), Wave 1 e-cigarette ever use (vs. never use), was associated with approximately twice the odds of progression to Wave 2 established cigarette smoking (OR: 2.23; 95% CI: 1.55, 3.21; $P<0.001$), past 30-day smoking (OR: 1.75; 95% CI: 1.35, 2.27; $P<0.001$), and current established smoking (OR: 2.43; 95% CI: 1.55, 3.80; $P<0.001$). Associations were attenuated in fully adjusted models (Table 1), but e-cigarette ever use remained a positive and statistically significant predictor of current established smoking (OR: 1.80; 95% CI: 1.04, 3.12; $P=0.035$). Associations did not reach the threshold for statistical significance for established smoking (OR: 1.57; 95% CI: 0.99, 2.49; $P=0.055$) and past 30-day smoking (OR: 1.32; 95% CI: 0.99, 1.76; $P=0.059$).

When baseline e-cigarette former use (tried but not used in past 30 days) and past 30-day use were considered separately, there was a step-wise increase in the probability of progression to future established smoking from never to former to past 30-day e-cigarette use (Table 1). For example, the probability of Wave 2 past 30-day cigarette smoking rose from baseline e-cigarette never use (26.6%) to former use (36.1%) to past 30-day use (45.3%). Both e-cigarette former use and past 30-day use remained statistically significantly associated with all three Wave 2 cigarette outcomes in models adjusted for gender, age, and race/ethnicity (Table 1). In fully adjusted models, baseline e-cigarette former use remained a statistically significant predictor of progression to current established smoking (OR: 1.85; 95% CI: 1.02, 3.36; $P=0.042$), and baseline e-cigarette past 30-day use statistically significantly predicted progression to past 30-day smoking (OR: 1.64; 95% CI: 1.12, 2.41; $P=0.010$). Adjustment variables that were consistently associated with greater progression to established smoking included household tobacco use and tobacco advertisement receptivity (Supplemental Table 1).

DISCUSSION

In this study, among youth who had experimented with cigarettes but had not progressed to established smoking, additional use of e-cigarettes was positively associated with future onset of current established smoking. Across three different definitions of established smoking and two different specifications of e-cigarette use, baseline e-cigarette users were at 1.5 to 2-times greater odds of progression to established smoking than e-cigarette never users after adjustment for confounding variables. Fully adjusted associations with e-cigarette ever use were statistically significant for one definition of established smoking (current established smoking; $P=0.035$) but fell just short of the a priori threshold for statistical significance for established smoking ($P=0.055$) and past 30-day smoking ($P=0.059$). The odds ratios in the present longitudinal analysis were in the same direction but smaller in magnitude than in the previous cross-sectional analysis of NYTS data¹³ that could have captured new trial of e-cigarettes among previously established smokers.

Regardless of how Wave 1 e-cigarette and Wave 2 smoking variables were specified, positive associations persisted after statistical adjustment for gender, age, and race/ethnicity. Adding to models the full set of confounding variables, such as household tobacco use, warning label exposure, and baseline use of other tobacco products, reduced the strength of some of the observed association to below the threshold for statistical significance. However, all associations remained positive in direction and similar in magnitude across different definitions of e-cigarette exposure and the smoking outcome.

These results suggest that e-cigarette use is more likely to encourage youth smoking than to divert youth from smoking when considering individuals who have already experimented with cigarette use. Unlike adults, particularly cigarette smokers, who commonly report a desire to quit smoking as a main motivator for e-cigarette use,¹⁸ youth are more likely to cite curiosity as a reason to try e-cigarettes.¹⁹ E-cigarette use was not associated with cigarette quit attempts or with quit contemplation among U.S. middle and high school students in any NYTS survey wave from 2011 to 2015.²⁰

In existing studies of youth who had never smoked a cigarette at baseline, those who tried e-cigarettes were more likely to initiate cigarette smoking in the future.^{2-6, 9} In addition to smoking initiation among youth never-smokers, the present study demonstrated that e-cigarette use was also associated with progression to current established smoking among youth smoking experimenters.

In a study of California 10th grade students that included never smokers and current smokers at baseline, greater frequency of e-cigarette use at baseline was associated with subsequently greater levels of smoking frequency (days smoked in past month) and heaviness (cigarettes smoked per day) six months later.²¹ Similarly, in a school-based study of adolescent never and current smokers in Canada, baseline past 30-day e-cigarette use was associated with initiation of daily smoking one year later.⁴ The results of a school-based study of baseline cigarette ever smokers in Hawaii did not yield a statistically significant change at follow-up in smoking frequency (measured as numerical categories) between baseline e-cigarette ever and never users.² However, in a school-based study of adolescents in the United Kingdom,

ever use (vs. never use) of e-cigarettes was associated with "escalation" to smoking sometimes or usually among baseline non-smokers who had used cigarettes in the past.⁵ That study reported an adjusted odds ratio that was very similar to the present study (OR: 1.89; 95% CI: 0.82, 4.33) but not statistically significant ($P=0.13$) in a smaller sample ($n=318$).⁵

The smoking outcomes evaluated in the present study represent intensity levels of clear clinical and public health concern. While, smoking as infrequently as one day in the past month in adolescence is predictive of adult smoking,²² youth who reach higher levels of smoking are even more likely to continue to smoke.¹² Additionally, while more recent (past 30-day) e-cigarette use was a stronger predictor of future established smoking than was former e-cigarette use in unadjusted models, this pattern did not necessarily persist in fully-adjusted models. This suggests that any level of e-cigarette use among adolescent cigarette experimenters may be a meaningful risk indicator of smoking progression.

Several study advantages strengthened the conclusions that can be drawn from this research. The large, prospective, and nationally representative nature of the PATH study enhanced generalizability and certainty regarding the temporal sequence between exposure and outcome. The PATH questionnaire was rigorously pilot-tested and administered under a consistent protocol.¹⁴ Furthermore, the magnitude of associations found in this study was largely consistent across different specifications of e-cigarette and cigarette use. Among other study aspects to consider, in-home administration of the PATH questionnaire could have led to differences in estimated tobacco use compared to school-based surveys. However, results of this analysis were qualitatively similar to prior work using NYTS data.¹³ As with all observational studies, residual confounding from unmeasured variables cannot be ruled out, although associations remained positive and at the threshold for statistical significance following adjustment for an extensive set of variables known to predict youth cigarette smoking.²³

In July 2017, the Food and Drug Administration announced a plan for tobacco and nicotine regulation that delayed federal e-cigarette regulation from 2018 until 2022.²⁴ However, local governments have taken regulatory action of e-cigarettes. For example, a 2017 San Francisco, California, ordinance prohibits the sale of flavored tobacco products, including e-cigarettes, with the intention of reducing the appeal of tobacco products to youth.²⁵ Our results indicate that among youth cigarette experimenters, those who have also used e-cigarettes are more likely to progress to current established smoking than those who tried cigarettes alone. As long as e-cigarettes remain attractive to youth, concern persists that these products contribute to greater combustible cigarette smoking among adolescents.

CONCLUSION

Among youth cigarette experimenters, using e-cigarettes was positively and independently associated with future current established smoking, suggesting that e-cigarettes do not divert from, and may encourage, cigarette smoking in this population. In weighing the overall public health impact of e-cigarette availability, regulation, and use, the potential to increase combustible cigarette smoking by youth deserves special consideration.

Supplementary Material

Refer to Web version on PubMed Central for supplementary material.

Acknowledgments

Funding: This work was supported in part by the US National Cancer Institute and Food and Drug Administration Center for Tobacco Products (P50CA180890), National Institute on Drug Abuse (R01DA043950), and National Cancer Institute (T32CA113710). The content is solely the responsibility of the authors and does not necessarily represent the official views of the National Institutes of Health or the FDA. The funding agencies played no role in the conduct of the research or the decision to submit the manuscript for publication.

Abbreviations

CI	confidence interval
NYTS	National Youth Tobacco Survey
OR	odds ratio
PATH	Population Assessment of Tobacco and Health

References

1. Jamal A, Gentzke A, Hu SS, et al. Tobacco use among middle and high school students - United States, 2011–2016. *MMWR Morb Mortal Wkly Rep.* 2017; 66(23):597–603. [PubMed: 28617771]
2. Wills TA, Knight R, Sargent JD, et al. Longitudinal study of e-cigarette use and onset of cigarette smoking among high school students in Hawaii. *Tob Control.* 2017; 26(1):34–39. [PubMed: 26811353]
3. Barrington-Trimis JL, Urman R, Berhane K, et al. E-cigarettes and future cigarette use. *Pediatrics.* 2016; 138(1)
4. Hammond D, Reid JL, Cole AG, Leatherdale ST. Electronic cigarette use and smoking initiation among youth: A longitudinal cohort study. *CMAJ.* 2017; 189(43):E1328–e1336. [PubMed: 29084759]
5. Conner M, Grogan S, Simms-Ellis R, et al. Do electronic cigarettes increase cigarette smoking in UK adolescents? Evidence from a 12-month prospective study. *Tob Control.* 2017
6. Watkins SL, Glantz SA, Chaffee BW. Noncigarette tobacco product use and future cigarette smoking among youth in the Population Assessment Of Tobacco and Health (PATH) study, 2013–2015. *JAMA Pediatr.* 2017 [accepted - in press].
7. Spindle TR, Hiler MM, Cooke ME, Eissenberg T, Kendler KS, Dick DM. Electronic cigarette use and uptake of cigarette smoking: A longitudinal examination of U.S. college students. *Addict Behav.* 2017; 67:66–72. [PubMed: 28038364]
8. Miech R, Patrick ME, O'Malley PM, Johnston LD. E-cigarette use as a predictor of cigarette smoking: results from a 1-year follow-up of a national sample of 12th grade students. *Tob Control.* 2017
9. Soneji S, Barrington-Trimis JL, Wills TA, et al. Association between initial use of e-cigarettes and subsequent cigarette smoking among adolescents and young adults: A systematic review and meta-analysis. *JAMA Pediatr.* 2017; 171(8):788–797. [PubMed: 28654986]
10. Schneider S, Diehl K. Vaping as a catalyst for smoking? An initial model on the initiation of electronic cigarette use and the transition to tobacco smoking among adolescents. *Nicotine Tob Res.* 2016; 18(5):647–653. [PubMed: 26386472]
11. Etter JF. Gateway effects and electronic cigarettes. *Addiction.* 2017 [Electronic publication ahead of press].

12. Sargent JD, Gabrielli J, Budney A, Soneji S, Wills TA. Adolescent smoking experimentation as a predictor of daily cigarette smoking. *Drug Alcohol Depend.* 2017; 175:55–59. [PubMed: 28391086]
13. Dutra LM, Glantz SA. Electronic cigarettes and conventional cigarette use among U.S. adolescents: A cross-sectional study. *JAMA Pediatr.* 2014; 168(7):610–617. [PubMed: 24604023]
14. Hyland A, Ambrose BK, Conway KP, et al. Design and methods of the Population Assessment of Tobacco and Health (PATH) Study. *Tob Control.* 2016; 26(4):371–378. [PubMed: 27507901]
15. Evans N, Farkas A, Gilpin E, Berry C, Pierce JP. Influence of tobacco marketing and exposure to smokers on adolescent susceptibility to smoking. *J Natl Cancer Inst.* 1995; 87(20):1538–1545. [PubMed: 7563188]
16. Stephenson MT, Hoyle RH, Palmgreen P, Slater MD. Brief measures of sensation seeking for screening and large-scale surveys. *Drug Alcohol Depend.* 2003; 72(3):279–286. [PubMed: 14643945]
17. Judkins DR. Fay's method for variance estimation. *J Official Stats.* 1990; 6(3):223.
18. Rutten LJ, Blake KD, Agunwamba AA, et al. Use of e-cigarettes among current smokers: Associations among reasons for use, quit intentions, and current tobacco use. *Nicotine Tob Res.* 2015; 17(10):1228–1234. [PubMed: 25589678]
19. Kong G, Morean ME, Cavallo DA, Camenga DR, Krishnan-Sarin S. Reasons for electronic cigarette experimentation and discontinuation among adolescents and young adults. *Nicotine Tob Res.* 2015; 17(7):847–854. [PubMed: 25481917]
20. Chaffee BW, Couch ET, Gansky SA. Trends in characteristics and multi-product use among adolescents who use electronic cigarettes, United States 2011–2015. *PLoS One.* 2017; 12(5):e0177073. [PubMed: 28475634]
21. Leventhal AM, Stone MD, Andrabi N, et al. Association of e-cigarette vaping and progression to heavier patterns of cigarette smoking. *JAMA.* 2016; 316(18):1918–1920. [PubMed: 27825000]
22. Saddleson ML, Kozlowski LT, Giovino GA, Homish GG, Mahoney MC, Goniewicz ML. Assessing 30-day quantity-frequency of U.S. adolescent cigarette smoking as a predictor of adult smoking 14 years later. *Drug Alcohol Depend.* 2016; 162:92–98. [PubMed: 26987520]
23. Wellman RJ, Dugas EN, Dutczak H, et al. Predictors of the onset of cigarette smoking: A systematic review of longitudinal population-based studies in youth. *Am J Prev Med.* 2016; 51(5): 767–778. [PubMed: 27180028]
24. FDA News Release. [Accessed November 1, 2017] FDA announces comprehensive regulatory plan to shift trajectory of tobacco-related disease, death. Published 2017. Web citation: <http://www.webcitation.org/6uf4uAFYK> Updated July 28, 2017
25. Swan, R. Group seeks referendum on flavored tobacco ban in S.F. *San Francisco Chronicle* 2017. Jul 12. <http://www.sfchronicle.com/politics/article/Group-seeks-referendum-on-flavored-tobacco-ban-in-11284771.php>

What's Known on This Subject

In prior studies of youth who have never smoked cigarettes, those who tried electronic cigarettes were more likely to initiate conventional cigarette smoking compared to e-cigarette never users. In cross-sectional studies, e-cigarette use is associated with established youth smoking.

What This Study Adds

Among youth who already experimented with cigarettes but were not yet established smokers, having used e-cigarettes was prospectively associated with future current established cigarette smoking. For these youth at high smoking risk, e-cigarettes appear to encourage progression to established smoking.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Table 1
 Progression from Cigarette Experimentation to Established Smoking According to Baseline E-Cigarette Use

	n	Weighted % with outcome	Unadjusted			Adjusted ^a			Adjusted ^b		
			OR (95% CI)	p-value	OR (95% CI)	p-value	OR (95% CI)	p-value	OR (95% CI)	p-value	
Outcome: Smoked 100 Cigarettes											
Wave 1 Predictors											
E-cigarette Never	646	9.7	reference		reference		reference		reference		
E-cigarette Ever	582	19.3	2.23 (1.55, 3.21)	<0.001	2.07 (1.41, 3.04)	<0.001	1.57 (0.99, 2.49)	0.05			
E-cigarette Never	646	9.7	reference		reference		reference		reference		
E-cigarette Former	406	18.6	2.13 (1.43, 3.18)	<0.001	2.04 (1.33, 3.12)	0.001	1.55 (0.94, 2.56)	0.09			
E-cigarette Past 30-Days	171	21.5	2.56 (1.58, 4.14)	<0.001	2.22 (1.31, 3.74)	0.003	1.69 (0.93, 3.05)	0.08			
Outcome: Smoked during Past 30 Days											
Wave 1 Predictors											
E-cigarette Never	699	26.6	reference		reference		reference		reference		
E-cigarette Ever	596	38.8	1.75 (1.35, 2.27)	<0.001	1.65 (1.26, 2.15)	<0.001	1.32 (0.99, 1.76)	0.06			
E-cigarette Never	699	26.6	reference		reference		reference		reference		
E-cigarette Former	415	36.1	1.56 (1.15, 2.12)	0.004	1.48 (1.09, 2.02)	0.01	1.20 (0.86, 1.68)	0.29			
E-cigarette Past 30-Days	176	45.3	2.29 (1.64, 3.19)	<0.001	2.10 (1.47, 2.99)	<0.001	1.64 (1.12, 2.41)	0.01			
Outcome: Smoked 100 Cigarettes and Smoked during Past 30 Days											
Wave 1 Predictors											
E-cigarette Never	644	7.1	reference		reference		reference		reference		
E-cigarette Ever	580	15.6	2.43 (1.55, 3.80)	<0.001	2.23 (1.39, 3.59)	<0.001	1.80 (1.04, 3.12)	0.03			
E-cigarette Never	644	7.1	reference		reference		reference		reference		
E-cigarette Former	406	15.5	2.41 (1.46, 3.97)	<0.001	2.29 (1.35, 3.89)	0.002	1.85 (1.02, 3.36)	0.04			
E-cigarette Past 30-Days	171	16.3	2.56 (1.52, 4.32)	<0.001	2.19 (1.24, 3.88)	0.007	1.76 (0.92, 3.37)	0.09			

^a. Model covariates include: gender, age, and race/ethnicity.

^b. Model covariates additionally include: parent education, urban residence, household tobacco use, alcohol ever use, tobacco advertisement receptivity, sensation-seeking score, cigarette warning label exposure, interview time of year, and ever use of any other tobacco product.

Author Manuscript

Author Manuscript

Author Manuscript

Author Manuscript

Abbreviations: CI = confidence interval; OR = odds ratio

Odds ratios and confidence intervals corresponding to model covariates are shown in Supplemental Table 1.